

UNIVERSIDADE ESTADUAL DE CAMPINAS
SISTEMA DE BIBLIOTECAS DA UNICAMP
REPOSITÓRIO DA PRODUÇÃO CIENTÍFICA E INTELLECTUAL DA UNICAMP

Versão do arquivo anexado / Version of attached file:

Versão do Editor / Published Version

Mais informações no site da editora / Further information on publisher's website:

<https://www.sciencedirect.com/science/article/pii/S2211601X16300050>

DOI: 10.1016/j.profoo.2016.05.005

Direitos autorais / Publisher's copyright statement:

©2016 by Elsevier. All rights reserved.

DIRETORIA DE TRATAMENTO DA INFORMAÇÃO

Cidade Universitária Zeferino Vaz Barão Geraldo

CEP 13083-970 – Campinas SP

Fone: (19) 3521-6493

<http://www.repositorio.unicamp.br>

9th International Conference on Predictive Modelling in Food

Estimation of the risk associated to marketing of swine meat contaminated with *Salmonella* spp., employing the William T. Fine method

Nataly Ruiz Quiñones^{a,b*}, Oscar Castro Aguilar^a, Martha Lily Ocampo Guerrero^a

^aResearch group GEBIUT, Tolima University, Ibagué 730006299, Colombia

^bDivision of microbial resources-Chemical, Biological and Agricultural Pluridisciplinary Research Center (CPQBA), Campinas state University UNICAMP, Paulinia 13148-218, Brazil

Abstract

Swine meat is a growing market in Colombia, reaching 3.1 million of carcasses sold in 2014. However, it is estimated that the proportion of hogs illegally marketed can be up to 50%, which exposes the population to foodborne pathogens such as Salmonella. The aim of this study was to estimate the risk associated to the marketing of swine meat contaminated with Salmonella, using the William T. Fine method in Tolima Department. The risk scenarios applied to four municipalities positive for Salmonella showed a medium and low level of risk, and an exposed population of 114,000 people, representing 8.7% of the Department's population. In the worst-case scenario, costs for hospitalization and compensation could represent more than US\$ 145 billion for the biggest municipality. It can be concluded that the William T. Fine method is applicable for biological risk analysis focused on foodborne pathogen outbreaks.

© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of Department of Food Science, Faculty of Food Engineering, University of Campinas.

Keywords: Salmonella; Swine meat; mathematic estimation; risk scenario; William T. Fine

* Corresponding author

E-mail address: nruizq@gmail.com

1. Introduction

Foodborne diseases (FBD) are a problem that affects both developed and developing countries. Salmonellosis is an infection caused by eating foods, such as pork, beef, poultry and other meat products, which are contaminated with *Salmonella* spp., causing gastroenteritis or typhoid fever in humans.^{1,2,3} In Colombia, in the period between January 2008 and August 2010, 102 FBD outbreaks were recorded, 31.7% being salmonellosis. In 2014, FBD caused by *Salmonella* spp. prevalence was 1.5%^{4,5}.

To reduce the incidence of FBD on food production chain, the implementation of control systems, like hazard analysis control point (HACCP), is necessary. Currently, different methodologies based in surveys to measure de risk in the food industry exist, but the implementations of these processes are delayed and complex. William T. Fine (1971)⁶ published a method to calculate risk and cost of the implementation of a remedial action to correct a hazard. This method combine all scenarios of consequence, exposition and probability of a possible hazard event occurrence. Currently, this method is used to estimate the degree of attention that should be paid to different risks, whether physical, chemical or biological, that occur in companies to establish priority ranges to address a risk between high, medium and low⁷.

The aim of this project was to estimate the prevalence of *Salmonella* spp. in slaughterhouses and outlets of swine meat. The analysis identified a prevalence of 5% in the six municipalities evaluated and sampled. Tolima Department have 47 municipalities and only eight slaughterhouses of swine meat. Nevertheless, only one met all requirement for pork slaughtering. Remaining seven need adjustments and the other municipalities have no interest in implementing pork slaughterhouses. Due to the circumstances exposed before, the risk associated to the marketing of swine meat contaminated with *Salmonella* was estimated using the William T. Fine method, assessing the costs for the medical attention if hypothetically a salmonellosis outbreak occurred.

2. Materials and methods

A survey of Good Practice of Manufacture (GPM) was done in six slaughterhouses and 73 outlets of swine meat, in 2010 in six municipalities of Tolima. Only four of the municipalities were salmonella positive. Estimation of risk with Fine method was performed on those four municipalities, by selecting the parameters (Table 1) and replacing them in the equation 1.

$$R = C \times E \times P \quad (1)$$

Where, **R** represents the risk; **C** are the consequences of the risk; **E** is the exposure to the risk; **P** is the probability that the risk occurs leading to the occurrence of an unwanted event.

Table 1. Descriptions and rating for the principal consequences, probability and risk exposure selected in this study⁶.

| Rating | Consequences |
|--------|---|
| 10 | Fatality: damages US\$ 5,000 to US\$ 10,000 |
| 6 | Extremely serious or disabling injuries: damages US\$ 1,000 to US\$ 5,000 |
| 4 | Injuries with temporary inability: damages US\$ 500 to US\$ 1,000 |
| 1 | Minor cuts, bruises, bumps: damage up to US\$ 500 |
| Rating | Probability |
| 7 | Is quite possible. Would not be unusual. 50% of probability. |
| Rating | Exposure |
| 6 | Frequently (approximately once daily) |
| 2 | Occasionally (from once per week to once per month) |

The **DR** (degree of risk) was calculated as the product of **R** and **HF** (Hazard factor). The HF (Table 2) was estimated as the percent of people exposed to swine meat contaminated with *Salmonella* (in concordance with the results of meat samples) in relation to the swine meat intake calculated in Colombia⁸.

Table 2. Estimation of population exposure to risk for intake of swine meat with *Salmonella*

| Municipality | Rating | Hazard Factor | Population exposure (approximation) |
|--------------|--------|---------------|--|
| Ibagué | 1 | 1 - 20% | 93172 |
| Líbano | 1 | 1 - 20% | 5154 |
| Fresno | 2 | 21 - 40% | 5707 |
| Guamo | 4 | 61 - 80% | 12930 |

3. Results and discussion

The estimation of risk by the William T. Fine method is used to calculate the degree of attention that should be paid to the different risks and it is implemented by security companies and banks^{6,7,9}. According to the data obtained in this study, the risk of contamination with *Salmonella* spp., in the four municipalities, is low and medium (Table 3). With an exposure to risk of 3% weekly, approximately.

Table 3. Results of William T. Fine estimation of risk for two scenarios and total cost for hypothetical outbreak

| Municipality | R (C x E x P) | HF (Rating of Hazard Factor) | DR (R x HF) | Cost for scenario (USD) | Total cost for outbreak (millions of USD) |
|--------------|-------------------------|--|-----------------------|----------------------------|--|
| Ibagué | 14 | 1 | 14 | 500 | \$ 241.67 |
| | 420 | 1 | 420 | 10000 | \$ 145,000.00 |
| Líbano | 14 | 1 | 14 | 500 | \$ 25.00 |
| | 420 | 1 | 420 | 10000 | \$ 15,000.00 |
| Fresno | 14 | 2 | 28 | 500 | \$ 58.33 |
| | 420 | 2 | 840 | 10000 | \$ 35,000.00 |
| Guamo | 14 | 4 | 56 | 500 | \$ 300.00 |
| | 420 | 4 | 1680 | 10000 | \$ 180,000.00 |

Unlike the HACCP system, the Fine method contemplates different risk scenarios, depending on the combination of consequences, exposure and probability, and estimates the cost that this would entail if those scenarios were presented. In USA, it is estimated that US\$ 3,000 million are spent, plus losses in productivity due to disability, for the treatment of salmonellosis. In Colombia, the San Ignacio Hospital reported that a patient with salmonellosis would entail a cost for hospitalization of US \$500 daily and a patient with salmonellosis complicated by immunosuppression hospitalized in intensive care can generate a daily cost of up to US \$ 1,200⁴. The hypothetical scenarios contemplated in this estimation reveal the importance of investing in the adequacy of facilities and training of food handlers, which would optimize the use of resources by government agencies.

4. Conclusion

The method of William T. Fine is applicable for biological risk analysis focused on the costs that are generated when an outbreak caused by food illness like Salmonella occurs. This method can be applied together with other systems of GPM in the food industry to establish management planes.

Acknowledgements

The Ministry of Agriculture and Rural Development (MADR) of the Republic of Colombia, financier agent of the project in which this work was developed.

Reference

1. World Organization for Animal Health [OIE]. Chapter 2.9.9 Salmonellosis. In: *Manual of Recommended Diagnostic Techniques and Requirements for Biological Products*. Paris: 2004. p. 1092-1107.
2. Oliveira FA d., Geimba MP, Pasqualotto AP, Brandelli A., Pasquali G., Silva W. d. & Tondo EC. Clonal relationship among Salmonella enterica serovar Enteritidis involved in foodborne outbreaks in Southern Brazil. *Food Control* 2009;**20**:606-10.
3. Mezal EH, Stefanova R. & Khan AA. Isolation and molecular characterization of Salmonella enterica serovar Javiana from food, environmental and clinical samples. *Int J Food Microbiol* 2013;**164**:113–8.
4. Ministerio de la protección social, Unidad de Evaluación de Riesgos para la Inocuidad de los Alimentos UERIA & Instituto Nacional de Salud INS. *Perfil de Riesgo Salmonella spp., (no tifoidea) en pollo entero y en piezas*. Bogotá D.C.: Imprenta Nacional de Colombia 2011 p. 102-4.
5. Sivigila. Enfermedades transmitidas por alimentos ETA. Boletín epidemiológico semanal. 2015. p. 13-6 on line. <http://www.ins.gov.co/boletin-epidemiologico/Boletn%20Epidemiolgico/2014%20Boletin%20epidemiologico%20semana%2053.pdf>
6. Fine TW. Mathematical evaluations for controlling hazards.1971. on line. <http://www.dtic.mil/dtic/tr/fulltext/u2/722011.pdf>
7. Mora VF & Aguilar ZE. Evaluación de Riesgos en Trabajos de Soldadura al Arco en el Proceso Productivo de la Empresa Astinave. (Tesis de Diplomado Superior en Seguridad y Salud Ocupacional) Universidad Estatal de Milagro. Unidad Académica Educación Continua A Distancia Y Postgrado Diplomado Superior De Seguridad Y Salud Ocupacional. Milagro, Ecuador. 2010.
8. Departamento Administrativo Nacional de Estadística (DANE). Boletín General departamento del Tolima. 2005.
9. Sociedad Colombiana de Medicina del Trabajo. Guía Técnica S. C. M. T. 005 A 03: metodología para realizar un panorama de riesgos valoración y priorización de los riesgos encontrados. *Revista de la Sociedad Colombiana de Medicina del Trabajo* 2004;**7**:41-6.

Presented at ICPMF9 (Set 8-12, 2015 – Rio de Janeiro, Brazil) as paper PMOD2015_0298.