



UNIVERSIDADE ESTADUAL DE CAMPINAS  
SISTEMA DE BIBLIOTECAS DA UNICAMP  
REPOSITÓRIO DA PRODUÇÃO CIENTÍFICA E INTELLECTUAL DA UNICAMP

**Versão do arquivo anexado / Version of attached file:**

Versão do Editor / Published Version

**Mais informações no site da editora / Further information on publisher's website:**

<https://www.sciencedirect.com/science/article/pii/S0927650506001733>

**DOI: 10.1016/j.astropartphys.2006.11.002**

**Direitos autorais / Publisher's copyright statement:**

©2007 by Elsevier. All rights reserved.

DIRETORIA DE TRATAMENTO DA INFORMAÇÃO

Cidade Universitária Zeferino Vaz Barão Geraldo

CEP 13083-970 – Campinas SP

Fone: (19) 3521-6493

<http://www.repositorio.unicamp.br>

# Anisotropy studies around the galactic centre at EeV energies with the Auger Observatory

The Pierre Auger Collaboration

J. Abraham<sup>f</sup>, M. Aglietta<sup>ao</sup>, C. Aguirre<sup>h</sup>, D. Allard<sup>bu</sup>, I. Allekotte<sup>a</sup>, P. Allison<sup>bq</sup>,  
C. Alvarez<sup>as</sup>, J. Alvarez-Muñiz<sup>bf</sup>, M. Ambrosio<sup>al</sup>, L. Anchordoqui<sup>bp,ca</sup>, J.C. Anjos<sup>j</sup>,  
C. Aramo<sup>al</sup>, K. Arisaka<sup>bt</sup>, E. Armengaud<sup>v</sup>, F. Arneodo<sup>ap</sup>, F. Arqueros<sup>bd</sup>, T. Asch<sup>ab</sup>,  
H. Asorey<sup>a</sup>, B.S. Atulugama<sup>br</sup>, J. Aublin<sup>u</sup>, M. Ave<sup>bu</sup>, G. Avila<sup>c</sup>, J. Bacelar<sup>ax</sup>, T. Backer<sup>af</sup>,  
D. Badagnani<sup>e</sup>, A.F. Barbosa<sup>j</sup>, H.M.J. Barbosa<sup>m</sup>, M. Barkhausen<sup>z</sup>, D. Barnhill<sup>bt</sup>,  
S.L.C. Barroso<sup>j</sup>, P. Bauleo<sup>bk</sup>, J. Beatty<sup>bq</sup>, T. Beau<sup>v</sup>, B.R. Becker<sup>by</sup>, K.H. Becker<sup>z</sup>,  
J.A. Bellido<sup>bz</sup>, S. BenZvi<sup>bl</sup>, C. Berat<sup>y</sup>, T. Bergmann<sup>ae</sup>, P. Bernardini<sup>aj</sup>, X. Bertou<sup>a</sup>,  
P.L. Biermann<sup>ac</sup>, P. Billoir<sup>x</sup>, O. Blanch-Bigas<sup>x</sup>, F. Blanco<sup>bd</sup>, P. Blasi<sup>ai,aq</sup>, C. Bleve<sup>bi</sup>,  
H. Blumer<sup>ae,aa</sup>, P. Boghrat<sup>bt</sup>, M. Bohacova<sup>t</sup>, C. Bonifazi<sup>j</sup>, R. Bonino<sup>ao</sup>, M. Boratav<sup>x</sup>,  
J. Brack<sup>bv</sup>, J.M. Brunet<sup>v</sup>, P. Buchholz<sup>af</sup>, N.G. Busca<sup>bu</sup>, K.S. Caballero-Mora<sup>ae</sup>, B. Cai<sup>bw</sup>,  
D.V. Camin<sup>ak</sup>, J.N. Capdevielle<sup>v</sup>, R. Caruso<sup>ar</sup>, A. Castellina<sup>ao</sup>, G. Cataldi<sup>aj</sup>, L. Cazon<sup>bu</sup>,  
R. Cester<sup>an</sup>, J. Chauvin<sup>y</sup>, A. Chiavassa<sup>ao</sup>, J.A. Chinellato<sup>m</sup>, A. Chou<sup>bm</sup>, J. Chye<sup>bo</sup>,  
D. Claes<sup>bx</sup>, P.D.J. Clark<sup>bh</sup>, R.W. Clay<sup>g</sup>, S.B. Clay<sup>g</sup>, B. Connolly<sup>bl</sup>, A. Cordier<sup>w</sup>,  
U. Cotti<sup>au</sup>, S. Coutu<sup>br</sup>, C.E. Covault<sup>bj</sup>, J. Cronin<sup>bu</sup>, S. Dagoret-Campagne<sup>w</sup>,  
T. Dang Quang<sup>cb</sup>, P. Darriulat<sup>cb</sup>, K. Daumiller<sup>aa</sup>, B.R. Dawson<sup>g</sup>, R.M. de Almeida<sup>m</sup>,  
L.A. de Carvalho<sup>m</sup>, C. De Donato<sup>ak</sup>, S.J. de Jong<sup>aw</sup>, W.J.M. de Mello Junior<sup>m</sup>,  
J.R.T. de Mello Neto<sup>q</sup>, I. De Mitri<sup>aj</sup>, M.A.L. de Oliveira<sup>o</sup>, V. de Souza<sup>l</sup>, L. del Peral<sup>be</sup>,  
O. Deligny<sup>u</sup>, A. Della Selva<sup>al</sup>, C. Delle Fratte<sup>am</sup>, H. Dembinski<sup>ad</sup>, C. Di Giulio<sup>am</sup>,  
J.C. Diaz<sup>bo</sup>, C. Dobrigkeit<sup>m</sup>, J.C. D’Oliveo<sup>av</sup>, D. Dornic<sup>u</sup>, A. Dorofeev<sup>bn</sup>, M.T. Dova<sup>e</sup>,  
D. D’Urso<sup>al</sup>, M.A. DuVernois<sup>bw</sup>, R. Engel<sup>aa</sup>, L. Epele<sup>e</sup>, M. Erdmann<sup>ad</sup>, C.O. Escobar<sup>m</sup>,  
A. Etchegoyen<sup>c</sup>, A. Ewers<sup>z</sup>, P. Facal San Luis<sup>bf</sup>, H. Falcke<sup>az,aw</sup>, A.C. Fauth<sup>m</sup>, D. Fazio<sup>ar</sup>,  
N. Fazzini<sup>bm</sup>, A. Fernandez<sup>as</sup>, F. Ferrer<sup>bj</sup>, S. Ferry<sup>bc</sup>, B. Fick<sup>bo</sup>, A. Filevich<sup>c</sup>,  
A. Filipcic<sup>bc</sup>, I. Fleck<sup>af</sup>, E. Fokitis<sup>ag</sup>, R. Fonte<sup>ar</sup>, D. Fuhrmann<sup>z</sup>, W. Fulgione<sup>ao</sup>,  
B. Garca<sup>f</sup>, D. Garcia-Pinto<sup>bd</sup>, L. Garrard<sup>bk</sup>, X. Garrido<sup>w</sup>, H. Geenen<sup>z</sup>, G. Gelmini<sup>bt</sup>,  
H. Gemmeke<sup>ab</sup>, A. Geranios<sup>ah</sup>, P.L. Ghia<sup>ao</sup>, M. Giller<sup>bb</sup>, J. Gitto<sup>f</sup>, H. Glass<sup>bm</sup>,  
F. Gobbi<sup>f</sup>, M.S. Gold<sup>by</sup>, F. Gomez Albarracin<sup>e</sup>, M. Gomez Berisso<sup>a</sup>, R. Gomez Herrero<sup>be</sup>,  
M. Gonalves do Amaral<sup>r</sup>, J.P. Gongora<sup>f</sup>, D. Gonzalez<sup>ae</sup>, J.G. Gonzalez<sup>bp</sup>, M. Gonzalez<sup>at</sup>,  
D. Gora<sup>ba,ae</sup>, A. Gorgi<sup>ao</sup>, P. Gouffon<sup>k</sup>, V. Grassi<sup>ak</sup>, A. Grillo<sup>ap</sup>, C. Grunfeld<sup>e</sup>,  
C. Grupen<sup>af</sup>, F. Guarino<sup>al</sup>, G.P. Guedes<sup>n</sup>, J. Gutierrez<sup>be</sup>, J.D. Hague<sup>by</sup>, J.C. Hamilton<sup>x</sup>,  
M.N. Harakeh<sup>ax</sup>, D. Harari<sup>a</sup>, S. Harmsma<sup>ax</sup>, S. Hartmann<sup>z</sup>, J.L. Harton<sup>bk</sup>, A. Haungs<sup>aa</sup>,  
M.D. Healy<sup>bt</sup>, T. Hebbeker<sup>ad</sup>, D. Heck<sup>aa</sup>, C. Hojvat<sup>bm</sup>, P. Homola<sup>ba</sup>, J. Horandel<sup>ae</sup>,  
A. Horneffer<sup>aw</sup>, M. Horvat<sup>bc</sup>, M. Hrabovsky<sup>t</sup>, T. Huege<sup>aa</sup>, M. Iarlori<sup>ai</sup>, A. Insolia<sup>ar</sup>,  
M. Kaducak<sup>bm</sup>, O. Kalashev<sup>bt</sup>, K.H. Kampert<sup>z</sup>, B. Keilhauer<sup>ae</sup>, E. Kemp<sup>m</sup>,

H.O. Klages<sup>aa</sup>, M. Kleifges<sup>ab</sup>, J. Kleinfeller<sup>aa</sup>, R. Knapik<sup>bk</sup>, J. Knapp<sup>bi</sup>, D.-H. Koang<sup>y</sup>,  
Y. Kolotaev<sup>af</sup>, A. Kopmann<sup>ab</sup>, O. Krömer<sup>ab</sup>, S. Kuhlman<sup>bm</sup>, J. Kuijpers<sup>aw</sup>, N. Kunka<sup>ab</sup>,  
A. Kusenko<sup>bt</sup>, C. Lachaud<sup>v</sup>, B.L. Lago<sup>q</sup>, D. Lebrun<sup>y</sup>, P. LeBrun<sup>bm</sup>, J. Lee<sup>bt</sup>,  
A. Letessier-Selvon<sup>x</sup>, M. Leuthold<sup>ad,bq</sup>, I. Lhenry-Yvon<sup>u</sup>, G. Longo<sup>al</sup>, R. López<sup>as</sup>,  
A. Lopez Agüera<sup>bf</sup>, A. Lucero<sup>f</sup>, S. Maldera<sup>ao</sup>, M. Malek<sup>bm</sup>, S. Maltezos<sup>ag</sup>,  
G. Mancarella<sup>aj</sup>, M.E. Manceñido<sup>e</sup>, D. Mandat<sup>t</sup>, P. Mantsch<sup>bm</sup>, A.G. Mariazzi<sup>bi</sup>,  
I.C. Maris<sup>ae</sup>, D. Martello<sup>aj</sup>, N. Martinez<sup>e</sup>, J. Martínez<sup>at</sup>, O. Martínez<sup>as</sup>, H.J. Mathes<sup>aa</sup>,  
J. Matthews<sup>bn,bs</sup>, J.A.J. Matthews<sup>by</sup>, G. Matthiae<sup>am</sup>, G. Maurin<sup>v</sup>, D. Maurizio<sup>an</sup>,  
P.O. Mazur<sup>bm</sup>, T. McCauley<sup>bp</sup>, M. McEwen<sup>bn</sup>, R.R. McNeil<sup>bn</sup>, G. Medina<sup>av</sup>,  
M.C. Medina<sup>c</sup>, G. Medina Tanco<sup>a</sup>, A. Meli<sup>ac</sup>, D. Melo<sup>c</sup>, E. Menichetti<sup>an</sup>, A. Menshikov<sup>ab</sup>,  
Chr. Meurer<sup>aa</sup>, R. Meyhandan<sup>bn</sup>, M.I. Micheletti<sup>c</sup>, G. Miele<sup>al</sup>, W. Miller<sup>by</sup>, S. Mollerach<sup>a</sup>,  
M. Monasor<sup>bd,be</sup>, D. Monnier Ragainé<sup>w</sup>, F. Montanet<sup>y</sup>, B. Morales<sup>av</sup>, C. Morello<sup>ao</sup>,  
E. Moreno<sup>as</sup>, C. Morris<sup>bq</sup>, M. Mostafá<sup>bz</sup>, M.A. Muller<sup>m</sup>, R. Mussa<sup>an</sup>, G. Navarra<sup>ao</sup>,  
L. Nellen<sup>av</sup>, C. Newman-Holmes<sup>bm</sup>, D. Newton<sup>bf</sup>, T. Nguyen Thi<sup>cb</sup>, R. Nichol<sup>bq</sup>,  
N. Nierstenhöfer<sup>z</sup>, D. Nitz<sup>bo</sup>, H. Nogima<sup>m</sup>, D. Nosek<sup>s</sup>, L. Nožka<sup>t</sup>, J. Oehlschläger<sup>aa</sup>,  
T. Ohnuki<sup>bt</sup>, A. Olinto<sup>bu</sup>, L.F.A. Oliveira<sup>q</sup>, V.M. Olmos-Gilbaja<sup>bf</sup>, M. Ortiz<sup>bd</sup>,  
S. Ostapchenko<sup>aa</sup>, L. Otero<sup>f</sup>, M. Palatka<sup>t</sup>, J. Pallotta<sup>f</sup>, G. Parente<sup>bf</sup>, E. Parizot<sup>u</sup>,  
S. Parlati<sup>ap</sup>, M. Patel<sup>bi</sup>, T. Paul<sup>bp</sup>, K. Payet<sup>y</sup>, M. Pech<sup>t</sup>, J. Pękala<sup>ba</sup>, R. Pelayo<sup>at</sup>,  
I.M. Pepe<sup>p</sup>, L. Perrone<sup>aj</sup>, S. Petrera<sup>ai</sup>, P. Petrinca<sup>am</sup>, Y. Petrov<sup>bk</sup>, D. Pham Ngoc<sup>cb</sup>,  
T.N. Pham Thi<sup>cb</sup>, R. Piegai<sup>e</sup>, T. Pierog<sup>aa</sup>, O. Pisanti<sup>al</sup>, T.A. Porter<sup>bn</sup>, J. Pouryamout<sup>z</sup>,  
L. Prado Junior<sup>m</sup>, P. Privitera<sup>am</sup>, M. Prouza<sup>bl</sup>, E.J. Quel<sup>f</sup>, J. Rautenberg<sup>z</sup>, H.C. Reis<sup>l</sup>,  
S. Reucroft<sup>bp</sup>, B. Revenu<sup>v</sup>, J. Rídký<sup>t</sup>, A. Risi<sup>f</sup>, M. Risse<sup>aa</sup>, C. Rivière<sup>y</sup>, V. Rizi<sup>ai</sup>,  
S. Robbins<sup>z</sup>, M. Roberts<sup>bs</sup>, C. Robledo<sup>as</sup>, G. Rodriguez<sup>bf</sup>, D. Rodríguez Frías<sup>be</sup>,  
J. Rodriguez Martino<sup>am</sup>, J. Rodriguez Rojo<sup>am</sup>, G. Ros<sup>bd,be</sup>, J. Rosado<sup>bd</sup>, M. Roth<sup>aa</sup>,  
C. Roucelle<sup>x</sup>, B. Rouillé-d'Orfeuil<sup>x</sup>, E. Roulet<sup>a</sup>, A.C. Rovero<sup>b</sup>, F. Salamida<sup>ai</sup>,  
H. Salazar<sup>as</sup>, G. Salina<sup>am</sup>, F. Sánchez<sup>c</sup>, M. Santander<sup>d</sup>, E.M. Santos<sup>j</sup>, S. Sarkar<sup>bg</sup>,  
R. Sato<sup>d</sup>, V. Scherini<sup>z</sup>, H. Schieler<sup>aa</sup>, T. Schmidt<sup>ae</sup>, O. Scholten<sup>ax</sup>, P. Schovánek<sup>t</sup>,  
F. Schüssler<sup>aa</sup>, S.J. Sciutto<sup>e</sup>, M. Scuderi<sup>ar</sup>, D. Semikoz<sup>v</sup>, G. Sequeiros<sup>an</sup>, R.C. Shellard<sup>j</sup>,  
B.B. Siffert<sup>q</sup>, G. Sigl<sup>v</sup>, P. Skelton<sup>bi</sup>, W. Slater<sup>bt</sup>, N. Smetniansky De Grande<sup>c</sup>,  
A. Smiałkowski<sup>bb</sup>, R. Šmída<sup>t</sup>, B.E. Smith<sup>bi</sup>, G.R. Snow<sup>bx</sup>, P. Sokolsky<sup>bz</sup>, P. Sommers<sup>br</sup>,  
J. Sorokin<sup>g</sup>, H. Spinka<sup>bm</sup>, E. Strazzeri<sup>am</sup>, A. Stutz<sup>y</sup>, F. Suarez<sup>ap</sup>, T. Suomijärvi<sup>u</sup>,  
A.D. Supanitsky<sup>c</sup>, J. Swain<sup>bp</sup>, Z. Szadkowski<sup>z,bb</sup>, A. Tamashiro<sup>b</sup>, A. Tamburro<sup>ae</sup>,  
O. Tascau<sup>z</sup>, R. Ticona<sup>i</sup>, C. Timmermans<sup>aw,ay</sup>, W. Tkaczyk<sup>bb</sup>, C.J. Todero Peixoto<sup>m</sup>,  
A. Tonachini<sup>an</sup>, D. Torresi<sup>ar</sup>, P. Travnicek<sup>t</sup>, A. Tripathi<sup>bt</sup>, G. Tristram<sup>v</sup>,  
D. Tscherniakhovski<sup>ab</sup>, M. Tueros<sup>e</sup>, V. Tunnicliffe<sup>bh</sup>, R. Ulrich<sup>aa</sup>, M. Unger<sup>aa</sup>,  
M. Urban<sup>w</sup>, J.F. Valdés Galicia<sup>av</sup>, I. Valiño<sup>bf</sup>, L. Valore<sup>al</sup>, A.M. van den Berg<sup>ax</sup>,  
V. van Elewyck<sup>u</sup>, R.A. Vazquez<sup>bf</sup>, D. Veberič<sup>bc</sup>, A. Veiga<sup>e</sup>, A. Velarde<sup>i</sup>, T. Venters<sup>bu</sup>,  
V. Verzi<sup>am</sup>, M. Videla<sup>f</sup>, L. Villaseñor<sup>au</sup>, T. Vo Van<sup>cb</sup>, S. Vorobiov<sup>v</sup>, L. Voyvodic<sup>bm</sup>,  
H. Wahlberg<sup>e</sup>, O. Wainberg<sup>c</sup>, T. Waldenmaier<sup>ae</sup>, P. Walker<sup>bh</sup>, D. Warner<sup>bk</sup>,  
A.A. Watson<sup>bi</sup>, S. Westerhoff<sup>bl</sup>, C. Wiebusch<sup>z</sup>, G. Wiczorek<sup>bb</sup>, L. Wiencke<sup>bz</sup>,  
B. Wilczyńska<sup>ba</sup>, H. Wilczyński<sup>ba</sup>, C. Wileman<sup>bi</sup>, M.G. Winnick<sup>g</sup>, J. Xu<sup>ab</sup>,  
T. Yamamoto<sup>bu</sup>, P. Younk<sup>bo</sup>, E. Zas<sup>bf</sup>, D. Zavrtnik<sup>bc</sup>, M. Zavrtnik<sup>bc</sup>, A. Zech<sup>x</sup>,  
A. Zepeda<sup>at</sup>, M. Zha<sup>bi</sup>, M. Ziolkowski<sup>af</sup>

<sup>a</sup> Centro Atómico Bariloche (CNEA), Instituto Balseiro (CNEA and UNCuyo), CONICET, 8400 San Carlos de Bariloche, Río Negro, Argentina<sup>b</sup> Instituto de Astronomía y Física del Espacio (CONICET), CC 67, Suc. 28 (1428) Buenos Aires, Argentina

- <sup>c</sup> Laboratorio Tandem (CNEA), CONICET, Univ. Tec. Nac. (Reg. Buenos Aires), Av. Gral. Paz 1499, (1650) San Martín, Buenos Aires, Argentina
- <sup>d</sup> Pierre Auger Southern Observatory, Av. San Martín Norte 304, (5613) Malargüe, Prov. De Mendoza, Argentina
- <sup>e</sup> Universidad Nacional de la Plata, Facultad de Ciencias Exactas, Departamento de Física and IFLP/CONICET, Univ. Nac. de Buenos Aires, FCEyN, Departamento de Física, C.C. 67, (1900) La Plata, Argentina
- <sup>f</sup> Universidad Tecnológica Nacional, Regionales Mendoza y San Rafael, CONICET, CEILAP-CITEFA, Rodríguez 273 Mendoza, Argentina
- <sup>g</sup> University of Adelaide, Department of Physics, Adelaide, SA 5005, Australia
- <sup>h</sup> Universidad Católica de Bolivia, Av. 16 Julio 1732, POB 5829, La Paz, Bolivia
- <sup>i</sup> Universidad Mayor de San Andrés, Av. Villazón N 1995, Monoblock Central, Bolivia
- <sup>j</sup> Centro Brasileiro de Pesquisas Físicas, Rua Dr. Xavier Sigaud, 150, CEP 22290-180 Rio de Janeiro, RJ, Brazil
- <sup>k</sup> Universidade de São Paulo, Inst. de Física, Cidade Universitária Caixa Postal 66318, Caixa Postal 66318, 05315-970 São Paulo, SP, Brazil
- <sup>l</sup> Universidade de São Paulo, Instituto Astronômico e Geofísico, Cidade Universitária, Rua do Matao 1226, 05508-900 São Paulo, SP, Brazil
- <sup>m</sup> Universidade Estadual de Campinas, Gleb Wataghin Physics Institute (IFGW), Departamento de Raios Cósmicos e Cronologia, CP 6165, 13083-970 Campinas, SP, Brazil
- <sup>n</sup> Univ. Estadual de Feira de Santana, Departamento de Física, Campus Universitário, BR 116, KM 03, 44031-460 Feira de Santana, Brazil
- <sup>o</sup> Universidade Estadual do Sudoeste da Bahia (UESB), Dep. Ciências Exatas, Estrada do Bem-Querer km4, 45083-900 Vitória da Conquista, BA, Brazil
- <sup>p</sup> Universidade Federal da Bahia, Campus da Ondina, 40210-340 Salvador, BA, Brazil
- <sup>q</sup> Univ. Federal do Rio de Janeiro (UFRJ), Instituto de Física, Cidade Universitária, Caixa Postal 68528, 21945-970 Rio de Janeiro, RJ, Brazil
- <sup>r</sup> Univ. Federal Fluminense, Inst. de Física, Campus da Praia Vermelha, 24210-340 Niterói, RJ, Brazil
- <sup>s</sup> Charles University, Institute of Particle and Nuclear Physics, Faculty of Mathematics and Physics, V Holesovickach 2, CZ-18000 Prague 8, Czech Republic
- <sup>t</sup> Institute of Physics of the Academy of Sciences of the Czech Republic, Na Slovance 2, CZ-182 21 Praha 8, Czech Republic
- <sup>u</sup> Institut de Physique Nucléaire, Université Paris-Sud 11 and IN2P3/CNRS, 15, rue Georges Clemenceau, 91400 Orsay, France
- <sup>v</sup> Laboratoire AstroParticule et Cosmologie, Université Paris VII, 11, Place Marcelin Berthelot, F-75231 Paris Cedex 05, France
- <sup>w</sup> Laboratoire de l'Accélérateur Linéaire, Université Paris-Sud 11 and IN2P3/CNRS, BP 34, Batiment 200, F-91898 Orsay cedex, France
- <sup>x</sup> Laboratoire de Physique Nucléaire et de Hautes Energies, Université Paris 6 & 7 and IN2P3/CNRS, 4 place Jussieu, 75252 Paris Cedex 05, France
- <sup>y</sup> Laboratoire de Physique Subatomique et de Cosmologie (LPSC), IN2P3/CNRS, Université Joseph-Fourier (Grenoble 1), 53, ave. des Martyrs, F-38026 Grenoble CEDEX, France
- <sup>z</sup> Bergische Universität Wuppertal, Fachbereich C – Physik, Gauß-Str. 20, D-42097 Wuppertal, Germany
- <sup>aa</sup> Forschungszentrum Karlsruhe, Institut für Kernphysik, Postfach 3640, D-76021 Karlsruhe, Germany
- <sup>ab</sup> Forschungszentrum Karlsruhe, Institut für Prozessdatenverarbeitung und Elektronik, Postfach 3640, D-76021 Karlsruhe, Germany
- <sup>ac</sup> Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany
- <sup>ad</sup> RWTH Aachen, III. Physikalisches Institut A, Physikzentrum, Huyskensweg, D-52056 Aachen, Germany
- <sup>ae</sup> Universität Karlsruhe (TH), Institut für Experimentelle Kernphysik (IEKP), Postfach 6980, D-76128 Karlsruhe, Germany
- <sup>af</sup> Universität Siegen, Fachbereich 7 Physik – Experimentelle Teilchenphysik, Emmy Noether-Campus, Walter-Flex-Str. 3, D-57068 Siegen, Germany
- <sup>ag</sup> Physics Department, School of Applied Sciences, National Technical University of Athens, Zografou 15780, Greece
- <sup>ah</sup> Physics Department, Nuclear and Particle Physics Section, University of Athens, Ilissia 15771, Greece
- <sup>ai</sup> Dipartimento di Fisica dell'Università de l'Aquila and INFN, Via Vetoio, I-67010 Coppito, Aquila, Italy
- <sup>aj</sup> Dipartimento di Fisica dell'Università di Lecce and Sezione INFN, via Arnesano, I-73100 Lecce, Italy
- <sup>ak</sup> Dipartimento di Fisica dell'Università di Milano and Sezione INFN, via Celoria 16, I-20133 Milan, Italy
- <sup>al</sup> Dipartimento di Fisica dell'Università di Napoli and Sezione INFN, Via Cintia 2, 80123 Napoli, Italy
- <sup>am</sup> Dipartimento di Fisica dell'Università di Roma II "Tor Vergata" and Sezione INFN, Via della Ricerca Scientifica, I-00133 Roma, Italy
- <sup>an</sup> Dipartimento di Fisica Sperimentale dell'Università di Torino and Sezione INFN, Via Pietro Giuria, 1, I-10125 Torino, Italy
- <sup>ao</sup> Istituto di Fisica dello Spazio Interplanetario (INAF), sezione di Torino and Dipartimento di Fisica Generale dell'Università and INFN Torino, Via P. Giuria 1, 10125 Torino, Italy
- <sup>ap</sup> INFN, Laboratori Nazionali del Gran Sasso, Strada Statale 17/bis Km 18+910, I-67010 Assergi (L'Aquila), Italy
- <sup>aq</sup> Osservatorio Astrofisico di Arcetri, Largo E. Fermi 5, I-50125 Florence, Italy
- <sup>ar</sup> Dipartimento di Fisica dell'Università di Catania and Sezione INFN, Corso Italia, 57, I-95129 Catania, Italy
- <sup>as</sup> Benemérita Universidad Autónoma de Puebla (BUAP), Ap. Postal J – 48, 72500 Puebla, Puebla, Mexico
- <sup>at</sup> Centro de Investigación y de Estudios Avanzados del IPN (CINVESTAV), Apartado Postal 14-740, 07000 México, D.F., Mexico
- <sup>au</sup> Universidad Michoacana de San Nicolás de Hidalgo (UMSNH), Edificio C-3 Cd Universitaria, C.P. 58040 Morelia, Michoacan, Mexico
- <sup>av</sup> Universidad Nacional Autónoma de México (UNAM), Apdo. Postal 20-364, 01000 México, D.F., México
- <sup>aw</sup> Department of Astrophysics, IMAPP, Radboud University, 6500 GL Nijmegen, Netherlands
- <sup>ax</sup> Kernfysisch Versneller Instituut (KVI), Rijksuniversiteit Groningen, Zernikelaan 25, NL-9747 AA Groningen, Netherlands
- <sup>ay</sup> NIKHEF, POB 41882, NL-1009 DB Amsterdam, Netherlands
- <sup>az</sup> ASTRON, PO Box 2, 7990 AA Dwingeloo, Netherlands
- <sup>ba</sup> Institute of Nuclear Physics PAN, Radzikowskiego 52, 31-342 Cracow, Poland
- <sup>bb</sup> University of Łódź, Pomorska 149/153, 90 236 Łódź, Poland
- <sup>bc</sup> University of Nova Gorica, Laboratory for Astroparticle Physics, Vipavska 13, POB 301, SI-5000 Nova Gorica, Slovenia
- <sup>bd</sup> Departamento de Física Atómica, Molecular y Nuclear, Facultad de Ciencias Físicas, Universidad Complutense de Madrid, E-28040 Madrid, Spain
- <sup>be</sup> Space Plasmas and Astroparticle Group, Universidad de Alcalá, Pza. San Diego, s/n, 28801 Alcalá de Henares (Madrid), Spain
- <sup>bf</sup> Departamento de Física de Partículas, Campus Sur, Universidad, E-15782 Santiago de Compostela, Spain
- <sup>bg</sup> Rudolf Peierls Centre for Theoretical Physics, University of Oxford, Oxford OX1 3NP, United Kingdom
- <sup>bh</sup> Institute of Integrated Information Systems, School of Electronic Engineering, University of Leeds, Leeds LS2 9JT, United Kingdom
- <sup>bi</sup> School of Physics and Astronomy, University of Leeds, Leeds LS2 9JT, United Kingdom
- <sup>bj</sup> Case Western Reserve University, Department of Physics, Cleveland, OH 44106, United States
- <sup>bk</sup> Colorado State University, Department of Physics, Fort Collins, CO 80523, United States
- <sup>bl</sup> Columbia University, Department of Physics, New York, NY 10027, United States

- <sup>bm</sup> Fermilab, MS367, POB 500, Batavia, IL 60510-0500, United States  
<sup>bn</sup> Louisiana State University, Department of Physics and Astronomy, Baton Rouge, LA 70803-4001, United States  
<sup>bo</sup> Michigan Technological University, Physics Department, 1400 Townsend Drive, Houghton, MI 49931-1295, United States  
<sup>bp</sup> Northeastern University, Department of Physics, 110 Forsyth Street, Boston, MA 02115-5096, United States  
<sup>bq</sup> Ohio State University, 2400 Olentangy River Road, Columbus, OH 43210-1061, United States  
<sup>br</sup> Pennsylvania State University, Department of Physics, 104 Davey Lab, University Park, PA 16802-6300, United States  
<sup>bs</sup> Southern University, Department of Physics, Baton Rouge, LA 70813-0400, United States  
<sup>bt</sup> University of California, Los Angeles (UCLA), Department of Physics and Astronomy, Los Angeles, CA 90095, United States  
<sup>bu</sup> University of Chicago, Enrico Fermi Institute, 5640 S. Ellis Avenue, Chicago, IL 60637, United States  
<sup>bv</sup> University of Colorado, Physics Department, Boulder, CO 80309-0446, United States  
<sup>bw</sup> University of Minnesota, School of Physics and Astronomy, 116 Church St. SE, Minneapolis, MN 55455, United States  
<sup>bx</sup> University of Nebraska, Department of Physics and Astronomy, 116 Brace Lab, Lincoln, NE 68588-0111, United States  
<sup>by</sup> University of New Mexico, Department of Physics and Astronomy, 800 Yale, Albuquerque, NM 87131, United States  
<sup>bz</sup> University of Utah, 115 S. 1400 East # 201, Salt Lake City, UT 84112-0830, United States  
<sup>ca</sup> University of Wisconsin-Milwaukee, Department of Physics, Milwaukee, WI 53201, United States  
<sup>cb</sup> Institute for Nuclear Science and Technology (INST), 5T-160 Hoang Quoc Viet Street, Nghia Do, Cau Giay, Hanoi, Viet Nam

Received 17 July 2006; received in revised form 6 November 2006; accepted 6 November 2006

Available online 18 December 2006

## Abstract

Data from the Pierre Auger Observatory are analyzed to search for anisotropies near the direction of the Galactic Centre at EeV energies. The exposure of the surface array in this part of the sky is already significantly larger than that of the fore-runner experiments. Our results do not support previous findings of localized excesses in the AGASA and SUGAR data. We set an upper bound on a point-like flux of cosmic rays arriving from the Galactic Centre which excludes several scenarios predicting sources of EeV neutrons from Sagittarius A. Also the events detected simultaneously by the surface and fluorescence detectors (the ‘hybrid’ data set), which have better pointing accuracy but are less numerous than those of the surface array alone, do not show any significant localized excess from this direction.

© 2006 Elsevier B.V. All rights reserved.

**Keywords:** Cosmic rays; Anisotropies; Galactic centre

## 1. Introduction

The Galactic Centre region constitutes an attractive target for cosmic ray (CR) anisotropy studies at EeV energies, where  $1 \text{ EeV} = 10^{18} \text{ eV}$ . These may be the highest energies for which the galactic component of the cosmic rays is still dominant. Moreover, since the Galactic Centre (GC) harbors the very massive black hole associated with the radio source Sagittarius A\*, as well as the expanding supernova remnant Sagittarius A East, it contains objects that might be candidates for powerful CR accelerators. The recent high significance observation by H.E.S.S. of a TeV  $\gamma$  ray source near the location of Sagittarius A\* [1], together with the discovery of a region of extended emission from giant molecular clouds in the central 200 pc of the Milky Way [2], further motivates the search for excesses in this direction. The location of the Pierre Auger Observatory in the southern hemisphere makes it particularly suitable for anisotropy studies in this region since the GC, passing only  $6^\circ$  from the zenith at the site, lies well within the field of view of the experiment. The number of CRs of EeV energies accumulated so far at the Pierre Auger Observatory from this part of the sky greatly exceeds that from previous

observations, allowing several interesting searches to be made.

There have been reports by the AGASA experiment [3,4] indicating a  $4.5\sigma$  excess of cosmic rays with energies in the range  $10^{18}$ – $10^{18.4}$  eV in a  $20^\circ$  radius region centred at right ascension and declination coordinates  $(\alpha, \delta) \simeq (280^\circ, -17^\circ)$ , in which the number of observed and expected events [4] are  $n_{\text{obs}}/n_{\text{exp}} = 506/413.6 = 1.22 \pm 0.05$ , where the error quoted is the one associated with Poisson background fluctuations. Note that the GC itself, for which we will adopt hereafter the Sagittarius A\* J2000.0 coordinates,  $(\alpha, \delta) = (266.3^\circ, -29.0^\circ)$ , lies outside the AGASA field of view ( $\delta > -24.2^\circ$ ). Later searches near this region with a reanalysis of SUGAR data [5], though with smaller statistics, failed to confirm these findings, but reported a  $2.9\sigma$  excess flux of CRs with energies in the range  $10^{17.9}$ – $10^{18.5}$  eV in a region of  $5.5^\circ$  radius centred at  $(\alpha, \delta) = (274^\circ, -22^\circ)$ , for which they obtained  $n_{\text{obs}}/n_{\text{exp}} = 21.8/11.8 = 1.85 \pm 0.29$ .

It is also sensible to search for a point-like excess from the GC. Due to the imperfect reconstruction of the arrival directions, the point source would be smeared on the angular scale of the resolution of the experiment. In

particular, EeV neutrons emitted by one of the possible energetic sources in the centre of the Galaxy may reach the Earth before decaying, and they would not be deflected by galactic magnetic fields. It is interesting to note that several scenarios predicting neutron fluxes from the GC detectable by Auger have been put forward in recent years [6–11].

In this work we use Auger data from the on-going construction phase to test the previous reports of localized excesses obtained with AGASA and SUGAR data, and to set limits on a CR flux from the GC direction in a window matched to the angular resolution of the experiment at EeV energies. A preliminary analysis of this kind was presented in [12].

The AGASA experiment has also reported a large scale anisotropy at EeV energies corresponding to a dipole-like modulation in right ascension of  $\sim 4\%$  amplitude, with a maximum near the GC and a deficit in the anti-centre direction. We defer the analysis of such large scale signatures for future work. This will require, in particular, control of the systematic uncertainty of the modulation of the exposure in right ascension induced by weather effects, which for the present Auger data set is estimated to be at a level of 1%. Uncertainties in the background estimates at this level do not affect the conclusions reached in the search for localized excesses performed in the present work.

## 2. Data set

The Auger surface detector [13], located in Malargüe, Argentina (latitude  $-35.2^\circ$ , longitude  $69.5^\circ$  W and mean altitude 1400 m a.s.l.), has been growing in size during the data taking period considered in this work, which goes from January 1st 2004 (when 154 detectors had been deployed) to March 30th 2006 (when 930 detectors were already deployed). The surface detectors consist of plastic tanks filled with 12,000 l of ultra-pure water in which the charged particles from the air showers produce Cherenkov light, which is reflected by the Tyvek™ liners and collected by three photomultipliers. The basic cell of the array is triangular, with separations of 1.5 km between detector units, and hence the complete array with 1600 detectors will cover an area of 3000 km<sup>2</sup>.

We consider the events from the surface detector (SD) array with three or more tanks triggered in a compact configuration. The events have to satisfy the level 5 quality trigger condition, which requires that the detector with the highest signal be surrounded by a hexagon of working detectors, since this ensures that the event is well reconstructed. We also restrict the events to zenith angles  $\theta < 60^\circ$ .

The energies are obtained using the inferred signal size at 1000 m from the reconstructed shower core,  $S(1000)$ , adopting a conversion that leads to a constant flux in different sky directions above 3 EeV, where the acceptance is saturated. This is the so-called Constant Intensity Cut cri-

terion implemented in [14]. A calibration of the energies is performed using clean fluorescence data, i.e. hybrid events that were recorded when there were contemporaneous aerosol measurements, whose longitudinal profiles include the shower maximum in a measured range of at least  $350 \text{ g cm}^{-2}$  and in which there is less than 10% Cherenkov contamination. The estimated systematic uncertainty in the reconstructed shower energy with the fluorescence technique is currently 25% [17]. For the hybrid events measured with both techniques the dispersion between SD and FD energy assignments is at the level of 35% in this energy range. From the uncertainty in the measurements of the signals from the Cherenkov tanks [15] the statistical uncertainty in the energy determination which results from the fitting procedure is about 20% for the energy range considered in this work, i.e.  $10^{17.9} \text{ eV} < E < 10^{18.5} \text{ eV}$ . Notice that in this energy range 48% of the events involve just three tanks, 34% involve 4 tanks and only 18% more than 4 tanks. For three tank events the 68% quantile angular resolution is about  $2.2^\circ$  and the resolution improves for events with 4 tanks or more [16].

Regarding the hybrid events, i.e. those with signal from both the fluorescence detectors (FD) and surface array, the angular resolution achieved is much smaller, typically below 1 degree [16]. Also, given that hybrid events may trigger with just one surface detector, the associated energy threshold ( $\sim 10^{17} \text{ eV}$ ) is lower, and events up to zenith angles of  $75^\circ$  are included in the data set. However, the statistics accumulated are significantly less, in part due to the  $\sim 15\%$  duty cycle of the fluorescence telescopes and also because at EeV energies the FD is not fully efficient at detecting showers over the full SD array. There are for instance 79265 SD events in the data set considered with energies  $10^{17.9} \text{ eV} < E < 10^{18.5} \text{ eV}$ , while the corresponding number of well reconstructed hybrid events in the same energy range is just 3439. Note that  $\sim 25\%$  of the hybrid events in this energy range involve less than three surface detectors, and are hence not included in the SD only data set.

## 3. Results

To study the possible presence of anisotropies, one needs first to obtain the background expectations for the different sky directions under the assumption of an isotropic CR distribution. This is a delicate issue since modulations of the exposure in right ascension are induced by the dead time of the detectors and the constantly growing array size. Also the effects of weather variations, especially near the energy threshold of the detector, may be non-negligible since they may affect the shower development in the atmosphere and/or the response of the electronics. Preliminary studies of these effects indicate that the possible weather-induced background modulations for the present data set are at a level of 1%, and are

hence below the Poisson noise for the angular windows considered.<sup>1</sup>

We have followed two different approaches [18] to estimate the isotropic expectations for the SD analysis:

- *The semi-analytic technique:* At EeV energies the zenith angle dependence of the exposure differs from the geometric one corresponding to full acceptance,  $dN \propto \sin \theta \cos \theta d\theta$ , mainly due to the attenuation in the atmosphere which affects large zenith angle showers. We therefore perform an analytic fit to the  $\theta$  distribution of the observed events in the energy range under study and then make a convolution with the number of hexagons with active detectors (which gives a measure of the aperture for events satisfying the quality trigger criterion) as a function of time, assuming a uniform response in azimuth. Through this procedure one obtains an exposure which accounts for the non-saturated acceptance effects and for the non-uniform running times and array growth. This technique allows to recover the detector's acceptance with negligible biases even in the case in which a large scale pattern is present in the CRs arrival distribution (see Ref. [18] for details).
- *The shuffling technique:* Here the expected number of events in any direction is obtained by averaging many data sets obtained by shuffling the observed events in the energy range of interest so that the arrival times are exchanged among them and the azimuths are drawn uniformly. The shuffling can be performed in separate zenith angle bins or by just mixing them all, and we found no significant difference between these two possibilities. By construction, this exposure preserves exactly the  $\theta$  distribution of the events and accounts for the detector dead times, array growth and even in principle for weather-induced modulations. It might however partially absorb modulations induced by large scale intrinsic anisotropies present in the CR flux, such as those due to a global dipole.

As implemented in the current analysis, the two techniques differ essentially in the treatment of the time dependence of the detectors acceptance. With shuffling we follow the detected rates while with the semi-analytic technique we assume a dependence only proportional to the detector size, and these two quantities differ only slightly.

The background estimate obtained with the shuffling technique in the GC region turns out to be about 0.5% larger than the one obtained with the semi-analytic method. Since this difference is much smaller than the size of the excesses that we are testing and is also below the level of the Poisson fluctuations, we will hence mainly quote in the following the values obtained using the semi-analytic technique.

### 3.1. Testing the AGASA and SUGAR excesses

In Fig. 1 we show a map of the GC region depicting the Li-Ma significances<sup>2</sup> [19] of overdensities in circular windows of  $5^\circ$  radius, for SD data with energies in the range  $10^{17.9}$ – $10^{18.5}$  eV. This angular scale is convenient to visualize the distribution of overdensities in the windows explored by SUGAR and AGASA. The galactic plane is represented with a solid line and the location of the Galactic Centre is indicated with a cross. The region in which AGASA reported an excess (in a slightly narrower energy range) is the big circle in the neighborhood of the GC, with the dashed line indicating the lower boundary of the region observed by AGASA. The smaller circle indicates the region where an excess in the SUGAR data was reported.

The size of the overdensities present in this map is consistent with what would be expected as a result of statistical fluctuations of an isotropic sky. Indeed, Fig. 2 depicts the distribution of these overdensities together with the expectations from an isotropic flux (average and  $2\sigma$  bounds obtained from Monte Carlo simulations), and no significant departure from isotropy is observed.

For the  $20^\circ$  circle centred at the AGASA location and for  $10^{18}$  eV  $< E < 10^{18.4}$  eV, 2116 events are observed while 2159.6 are expected using the semi-analytic technique, while 2169.7 are expected using the shuffling technique. It is clear that no significant excess is observed. Note that the number of events is more than four times that collected by AGASA in this region, in part due to the fact that the GC lies well within the field of view of Auger, and in part due to the fact that the total exposure of Auger is already double that achieved by AGASA.

It must be borne in mind that there may be systematic differences in the energy calibration of the two experiments. To test whether these differences could have possibly masked the AGASA reported excess, we show in Table 1 the observed and expected rates for different energy ranges, offset by 0.1 decade in energy (i.e. by about 25%), keeping  $E_{\max}/E_{\min}$  fixed. We have added a systematic error of 1% to the expected rates to account for the effects of possible weather induced modulations. These results show that no significant excesses are seen in the AGASA region for any of these cases. In particular, at the  $2\sigma$  level the excess in this region is always less than 6%, well below the 22% excess reported by AGASA.

Since it is conceivable that particles leading to a localized excess are different from the bulk of the CRs (e.g. if they are nucleons and the bulk of the CRs in this energy range are heavier nuclei), one may also wonder if the Auger sensitivity to these particles could be reduced. In particular, since for Auger the acceptance in this energy range is not yet saturated, it will be larger for heavy nuclei than for

<sup>1</sup> A detailed account of weather effects is certainly necessary to test large scale patterns at the few percent level. Relevant studies are in progress.

<sup>2</sup> For the  $\alpha$  parameter in the expression of the Li-Ma significance we use  $\alpha = n_{\text{exp}}/n_t$ , with  $n_t$  the total number of events in the energy range considered and  $n_{\text{exp}}$  the background expected in the angular region searched.




Fig. 1. Map of CR overdensity significances near the GC region on top-hat windows of  $5^\circ$  radius. The GC location is indicated with a cross, lying along the galactic plane (solid line). Also the regions where the AGASA experiment found their largest excess as well as the region of the SUGAR excess are indicated.




Fig. 2. Histogram of overdensities on  $5^\circ$  radius windows and for  $10^{17.9} \text{ eV} < E < 10^{18.5} \text{ eV}$ , together with isotropic expectations (average and  $2\sigma$  bounds). Overdensities are computed on a grid of  $3^\circ$  spacing for the patch of the sky depicted in Fig. 1.

Table 1  
Events in the AGASA region for different shifted energy intervals

$E_{\min}$ [eV]	$E_{\max}$ [eV]	$n_{\text{obs}}/n_{\text{exp}}$
$10^{17.9}$	$10^{18.3}$	$3179/3153.5 = 1.01 \pm 0.02(\text{stat}) \pm 0.01(\text{syst})$
$10^{18}$	$10^{18.4}$	$2116/2159.5 = 0.98 \pm 0.02(\text{stat}) \pm 0.01(\text{syst})$
$10^{18.1}$	$10^{18.5}$	$1375/1394.5 = 0.99 \pm 0.03(\text{stat}) \pm 0.01(\text{syst})$

protons because showers initiated by heavier primaries develop earlier and are hence more spread out at ground level. Using the estimates in [20] for the acceptance of p

and Fe primaries, we find that the sensitivity to protons is about  $\sim 30\%$  smaller than to Fe in the energy range studied (assuming an  $E^{-3}$  spectrum). In the case in which the 22% excess reported by AGASA (which had full efficiency at EeV energies) was due to nucleons while the background was due to heavy nuclei, at least a 15% excess should have been expected in Auger data. This is much larger than the upper limit we are obtaining.

Regarding the localized excess observed in SUGAR data, we find in the same angular window and energy range that  $n_{\text{obs}}/n_{\text{exp}} = 286/289.7 = 0.98 \pm 0.06$ , and hence with more than an order of magnitude larger statistics no significant excess is seen in this window. Shifting the energy range to account for possible offsets also resulted in no significant excess.

### 3.2. Bounds on a point-like neutron source at the GC

#### 3.2.1. The surface detector results

The optimal search for a point-like source is best done using a Gaussian filter matching the angular resolution of the experiment [21]. For this we can assume that the reconstructed directions are distributed with respect to the true direction (separated by an angle  $\beta$ ) according to  $\exp(-\beta^2/2\sigma^2)$  per unit solid angle, where  $\sigma \simeq 1.5^\circ$  at EeV energies, corresponding to a 68% quantile of  $2.25^\circ$ , where we have ignored a mild zenith angle dependence for simplicity.

We use for this search an energy range between  $E_{\min} = 10^{17.9} \text{ eV}$  and  $E_{\max} = 10^{18.5} \text{ eV}$ . Below  $E_{\min}$  the Auger SD acceptance is very suppressed. Note also that most neutrons from a source at the GC would have decayed in flight

before reaching the Earth for lower energies. On the other hand, energies above  $E_{\max}$  may be hard to achieve for galactic sources.

For the Gaussian window centred in the Sagittarius A\* direction we get  $n_{\text{obs}}/n_{\text{exp}} = 53.8/45.8$ . This corresponds to a ratio of  $1.17 \pm 0.10$ , where the estimate of the uncertainty takes into account that the window is Gaussian. Applying the results of [21], we get a 95% CL upper bound on the number of events from the source of  $n_s^{95} = 18.5$ . To translate this into a bound on the source flux we make two assumptions:

- We assume that the spectrum of the source is similar to that of the CRs, which is approximately  $\propto E^{-3.3}$  in this energy range. If the source spectrum were actually harder, the bound we obtain would be a conservative one.
- We assume that the composition of the CRs in this energy range is similar to that of the source, i.e. proton-like. We will then discuss how the limit is modified if the CRs were heavier, in which case the detector acceptance would be different for the bulk of the CRs and for the neutron source.

Under these assumptions, the energy dependent acceptance of the detector has the same effect upon the source flux and the background flux, so that one can relate the ratio between the CR flux and the expected number of background events in this window, with the ratio between the source flux upper limit and the bound obtained for  $n_s^{95}$ .

We take for the differential CR spectrum flux the expression

$$\Phi_{\text{CR}}(E) \simeq \kappa 50 \left( \frac{E}{\text{EeV}} \right)^{-3.3} \text{EeV}^{-1} \text{km}^{-2} \text{yr}^{-1} \text{sr}^{-1}, \quad (1)$$

which has an  $E^{-3.3}$  dependence (consistent with the value found e.g. by HiRes [22] in the energy range  $10^{17.5} \text{eV} < E < 10^{18.5} \text{eV}$ ), and is a smooth extrapolation of the spectrum measured at the Auger Observatory<sup>3</sup> at  $E > 3 \text{EeV}$ .

The factor  $\kappa$  is introduced to parameterise our limited knowledge of the true CR flux and it should be of order unity according to the existing measurements of the spectrum at EeV energies. Note that at  $\sim 3 \text{EeV}$  the normalisation of the HiRes and AGASA spectra are above the one reported by Auger. In particular, the HiRes normalisation would correspond to adopting  $\kappa = 1.2$  while the AGASA normalisation would correspond to a value for  $\kappa$  of about 2.

Consider a Gaussian filter matching the angular resolution characterized by  $\sigma$

$$W(\beta) \equiv \exp\left(-\frac{\beta^2}{2\sigma^2}\right), \quad (2)$$

where  $\beta$  is the angle from the direction of Sagittarius A\*. Then the expected number of events in the specified energy range is

$$n_{\text{exp}} = 2\pi \int_0^\pi d\beta \sin\beta W(\beta) \int_{E_{\min}}^{E_{\max}} dE A(E) \Phi_{\text{CR}}(E), \quad (3)$$

where  $A(E)$  is the energy dependent exposure of the experiment. Similarly, the number of events expected to be observed from the point-like source will be

$$n_s = \int_0^\pi \frac{d\beta \sin\beta}{\sigma^2} W(\beta)^2 \int_{E_{\min}}^{E_{\max}} dE A(E) \Phi_s(E), \quad (4)$$

where we take into account that, due to the finite angular resolution of the experiment, the arrival directions of the observed source events are expected to be distributed according to

$$\frac{d\Phi_s}{d\Omega}(\beta, E) = \frac{\exp(-\beta^2/2\sigma^2)}{2\pi\sigma^2} \Phi_s(E). \quad (5)$$

Using the assumptions noted above, we then get an expression for the source flux integrated over the energy range considered,

$$\Phi_s \equiv \int_{E_{\min}}^{E_{\max}} dE \Phi_s(E) \quad (6)$$

with a 95% CL upper bound of

$$\Phi_s^{95} = \frac{n_s^{95}}{n_{\text{exp}}} 4\pi\sigma^2 \int_{E_{\min}}^{E_{\max}} dE \Phi_{\text{CR}}(E) = \kappa 0.13 \text{km}^{-2} \text{yr}^{-1}. \quad (7)$$

Note that the bound on the source flux just scales with the parameter  $\kappa$ , because what is constrained is the ratio between the source and background fluxes.

Let us now discuss how the bound would change if the bulk of the CRs were heavy nuclei in this energy range. Following the discussion in the previous Section, we conclude that the upper limit to the flux from the putative source will have to be scaled by a factor  $\sim 1.3$  under the assumption that the CRs are iron nuclei and that the source is a source of neutrons. We thus see that the bound on the neutron flux could be up to  $\sim 30\%$  higher if the CR composition at EeV energies were heavy.

Due to the steeply falling CR spectrum, the bound in Eq. (7) also holds for  $E_{\max} \rightarrow \infty$ , i.e. in the inclusive range  $E > 10^{17.9} \text{eV}$ . Setting instead  $E_{\min} = 1 \text{EeV}$ , the corresponding bound is  $\Phi_s^{95} = \kappa 0.06 \text{km}^{-2} \text{yr}^{-1}$ .

We point out that some of the theoretical predictions for neutron fluxes (those associated with the AGASA claim, but not those associated with the TeV results) are based on the AGASA normalization for the CR flux, which is about a factor of 3 larger than the Auger flux normalization. The earlier predictions must thus be reduced by this factor to be compared with the flux bounds obtained here. The predictions of Refs. [7], [8] and [9], which exceed the upper-bound obtained by more than one order of magnitude, are already excluded, and that of [10] is at the level of the present Auger sensitivity.

<sup>3</sup> A power law fit to the Auger Observatory measurements [14] leads to  $\Phi_{\text{CR}}(E) = (30.9 \pm 1.7) \times (E/\text{EeV})^{-2.84 \pm 0.03} \text{EeV}^{-1} \text{km}^{-2} \text{yr}^{-1} \text{sr}^{-1}$  (statistical error only).

### 3.2.2. The hybrid results

We have also studied the GC region as observed with hybrid events, detected by both the FD and SD. These events have a better angular resolution [16] ( $0.7^\circ$  at 68% C.L. in the energy range studied).

Considering the events with  $10^{17.9} \text{ eV} < E < 10^{18.5} \text{ eV}$ , no significant excess is seen in the GC direction. For instance, in an optimal top-hat window of  $1.59\sigma \simeq 0.75^\circ$  radius, 0.3 events are expected (as estimated using a shuffling method) while no single event direction falls within that circle. This leads to a source flux upper-bound at 95% CL of

$$\Phi_s^{95} = \kappa 0.24 \text{ km}^{-2} \text{ yr}^{-1}, \quad (8)$$

which is about a factor of 2 weaker than the SD flux bound. Note that the energy assignments of the FD apply regardless of the assumed CR composition (except for a small correction to account for the missing energy), be they protons or heavy nuclei. However, the acceptance has a dependence on composition because different primaries develop at different depths in the atmosphere. Since a quality requirement for hybrid events is to have the maximum of the shower development inside the field of view of the telescopes, this affects the sensitivity to different primaries. The bound obtained is indeed a conservative one if the bulk of the CRs are heavy nuclei.

### 3.2.3. Relation to a point-like photon source

In [1] the H.E.S.S. collaboration has reported a remarkably flat spectrum of gamma rays above 165 GeV (and up to 10 TeV) from the direction of Sagittarius A\*. A naive extrapolation of this spectrum would lead to a flux of gamma rays above 1 EeV of  $0.04 \text{ km}^{-2} \text{ yr}^{-1}$ . Note however that the bound obtained by us for a neutron source (which is comparable to this extrapolation) does not apply straightforwardly for photon primaries, since the acceptance (and energy assignments) are modified.

The spectrum of photons reported from the GC ridge [2] is also remarkably flat so that this region too merits future study. The Galactic Centre may house sources of very high-energy cosmic rays detectable through gamma radiation. It is clear then that further exposure with the Auger Observatory of this region and a dedicated analysis will be of interest. Also an exploration down to the FD threshold will be important for the search of photon sources.

## 4. Conclusions

Using the first 2.3 years of Auger data we have searched for localized anisotropies near the direction of the Galactic Centre, which is well within the field of view of the Observatory. With statistics much greater than those of previous experiments, we have looked for a point-like source in the direction of Sagittarius A, without finding a significant excess. This excludes several scenarios of neutron sources in the GC suggested recently. Our searches on larger angular windows in the neighborhood of the GC do not show

abnormally over-dense regions. In particular, they do not support the large excesses reported in AGASA data (of 22% on  $20^\circ$  scales) and SUGAR data (of 85% on  $5.5^\circ$  scales).

## Acknowledgements

We are very grateful to the following agencies and organizations for financial support: Gobierno de Mendoza, Comisión Nacional de Energía Atómica y Municipalidad de Malargüe, Argentina; the Australian Research Council; Fundacao de Amparo a Pesquisa do Estado de Sao Paulo, Conselho Nacional de Desenvolvimento Científico e Tecnológico and Fundacao de Amparo a Pesquisa do Estado de Rio de Janeiro, Brasil; National Science Foundation of China; Ministry of Education of the Czech Republic (projects LA134 and LN00A006); Centre National de la Recherche Scientifique, Institut National de Physique Nucléaire et Physique des Particules (IN2P3/CNRS), Institut National des Sciences de l'Univers (INSU/CNRS) et Conseil Régional Ile de France, France; German Ministry for Education and Research and Forschungszentrum Karlsruhe, Germany; Istituto Nazionale di Fisica Nucleare, Italy; Consejo Nacional de Ciencia y Tecnología, Mexico; the Polish State Committee for Scientific Research (grant numbers 1P03D 01430, 2P03B 11024 and 2PO3D 01124), Poland; Slovenian Research Agency; Ministerio de Educación y Ciencia (FPA2003-08733-C02, 2004-01198), Xunta de Galicia (2003 PXIC20612PN, 2005 PXIC20604PN) and Feder Funds, Spain; Particle Physics and Astronomy Research Council, UK; the US Department of Energy, the US National Science Foundation, USA; UNESCO; and ALFA-EC funds, in the framework of the HELEN Project.

## References

- [1] F. Aharonian et al., *Astron. Astrophys.* 425 (2004) L13. astro-ph/0408145.
- [2] F. Aharonian et al., H.E.S.S. Collaboration, *Nature* 439 (2006) 695. arXiv:astro-ph/0603021.
- [3] N. Hayashida et al., AGASA Collaboration, *Astropart. Phys.* 10 (1999) 303. arXiv:astro-ph/9807045.
- [4] (a) N. Hayashida et al. (AGASA Collaboration), in: *Proceedings of the 26th ICRC, Salt Lake City, vol. 3, 1999, p. 256*;  
(b) M. Teshima et al. (AGASA Collaboration), in: *Proceedings of the 27th ICRC, Hamburg, vol. 1, 2001, p. 337*.
- [5] J.A. Bellido, R.W. Clay, B.R. Dawson, M. Johnston-Hollitt, *Astropart. Phys.* 15 (2001) 167. arXiv:astro-ph/0009039.
- [6] G.A. Medina Tanco, A.A. Watson, in: *Proceedings of the 27th ICRC, Hamburg, 2001, p. 531*.
- [7] M. Bossa, S. Mollerach, E. Roulet, *J. Phys. G: Nucl. Part. Phys.* 29 (2003) 1409. arXiv:astro-ph/0304023.
- [8] F. Aharonian, A. Neronov, *Astrophys. J.* 619 (2005) 306. arXiv:astro-ph/0408303.
- [9] R.M. Crocker, M. Fatuzzo, J.R. Jokipii, F. Melia, R.R. Volkas, *Astrophys. J.* 622 (2005) 892. arXiv:astro-ph/0408183.
- [10] D. Grasso, L. Maccione, *Astropart. Phys.* 24 (2005) 273. arXiv:astro-ph/0504323.
- [11] P.L. Biermann, G.A. Medina-Tanco, R. Engel, G. Pugliese, *Astrophys. J.* 604 (2004) L29. arXiv:astro-ph/0401150.

- [12] The Pierre Auger Collaboration, in: Proceedings of the 29th ICRC, Pune, V7, p. 67. arXiv:astro-ph/0507331.
- [13] The Pierre Auger Collaboration, Nucl. Instr. Meth. A 523 (2004) 50.
- [14] The Pierre Auger Collaboration, in: Proceedings of the 29th ICRC, Pune, V7, p. 387. arXiv:astro-ph/0507150.
- [15] The Pierre Auger Collaboration, in: Proceedings of the 29th ICRC, Pune, V7, p. 167.
- [16] The Pierre Auger Collaboration, in: Proceedings of the 29th ICRC, Pune, V7, p. 17.
- [17] The Pierre Auger Collaboration, in: Proceedings of the 29th ICRC, Pune, V7, p. 13.
- [18] The Pierre Auger Collaboration, in: Proceedings of the 29th ICRC, Pune, V7, p. 63. arXiv:astro-ph/0507517.
- [19] T.-P. Li, Y.-Q. Ma, Astrophys. J. 272 (1983) 317.
- [20] D. Allard et al., for the Pierre Auger Collaboration, in: Proceedings of the 29th ICRC, Pune, V7, p. 71. arXiv:astro-ph/0511104.
- [21] P. Billoir, A. Letessier-Selvon, arXiv:astro-ph/0507538.
- [22] D.R. Bergman, for the HiRes Collaboration, in: Proceedings of the 29th ICRC, Pune, V7, p. 101. arXiv:astro-ph/0507484.