



FACULDADE DE ODONTOLOGIA DE PIRACICABA
UNIVERSIDADE ESTADUAL DE CAMPINAS
DEPARTAMENTO DE ODONTOLOGIA RESTAURADORA



Área de Dentística

Monografia de Especialização

Versatilidade das técnicas de clareamento dental:

Combinação de técnicas de clareamento caseiro e de consultório- relato de casos

Aluna: Marcela Alvarez Ferretti

Orientador: Prof. Flávio Henrique Baggio Aguiar

Piracicaba
2021

Marcela Alvarez Ferretti

*Versatilidade das técnicas de clareamento dental:
combinação de técnicas de clareamento caseiro e consultório- relato de casos*

Monografia apresentada à Faculdade de Odontologia de Piracicaba da Universidade Estadual de Campinas como parte dos requisitos exigidos para a obtenção do título de Especialista em Dentística.

Orientador: Prof. Flávio Henrique Baggio Aguiar

ESTE EXEMPLAR CORRESPONDE À
VERSÃO FINAL DA MONOGRAFIA
APRESENTADA PELO ALUNO
MARCELA ALVAREZ FERRETTI, E
ORIENTADA PELO PROF. DR.
FLÁVIO HENRIQUE BAGGIO
AGUIAR.

Piracicaba
2021

Ficha catalográfica
Universidade Estadual de Campinas
Biblioteca da Faculdade de Odontologia de Piracicaba
Marilene Girello - CRB 8/6159

F415v Ferretti, Marcela Alvarez, 1991-
Versatilidade das técnicas de clareamento dental : combinação de técnicas de clareamento caseiro e de consultório - relato de casos / Marcela Alvarez Ferretti. – Piracicaba, SP : [s.n.], 2021.

Orientador: Flávio Henrique Baggio Aguiar.
Trabalho de Conclusão de Curso (especialização) – Universidade Estadual de Campinas, Faculdade de Odontologia de Piracicaba.

1. Dentes - Clareamento. 2. Peróxido de hidrogênio. 3. Peróxido de carbamida. 4. Combinação de medicamentos. I. Aguiar, Flavio Henrique Baggio, 1977-. II. Universidade Estadual de Campinas. Faculdade de Odontologia de Piracicaba. III. Título.

Informações adicionais, complementares

Título em outro idioma: Combination of at-home and in-office bleaching techniques - case series: bleaching combination techniques

Palavras-chave em inglês:

Teeth - Bleaching

Hydrogen peroxide

Carbamide peroxide

Drug combinations

Área de concentração: Dentística

Titulação: Especialista

Data de entrega do trabalho definitivo: 28-09-2021

Agradecimentos

Primeiramente, agradeço à Deus por todas as oportunidades que Ele vem colocando em minha vida, e por me permitir ter saúde e disposição para realizá-las. Agradeço à minha família que esteve comigo em todos os momentos, me apoiando e me dando todo o suporte que necessitei para que chegasse até aqui, por mais difíceis que estes fossem. Eu amo vocês!

Agradeço também ao diretor da Faculdade de Odontologia de Piracicaba da Universidade de Campinas, na pessoa de seu diretor, Prof. Dr. Francisco Haiter Neto.

Um agradecimento especial ao meu orientador Prof. Flávio H.B. Aguiar que sempre se fez presente durante a minha caminhada desde a graduação até a pós-graduação, compartilhando muito conhecimento e acreditando no meu potencial. Te admiro muito pelo profissional que és.

Aos demais professores da Área de Dentística, Luis Roberto Marcondes Martins, Marcelo Giannini, Giselle Maria Marchi, Vanessa Cavalli e Débora Alves Nunes Leite Lima, pela convivência diária e pelo conhecimento transmitido durante o curso de especialização.

Dedicatória

Dedico este trabalho a todos os colaboradores que juntamente a mim se dedicaram ao atendimento dos pacientes e elaboração do artigo. Matheus Kury, Beatriz Mendonça, Vanessa Cavalli, Marcelo Giannini e Flávio Aguiar, obrigada por dividirem seus conhecimentos comigo, pela parceria de trabalho e por tornarem a publicação dos casos possível.

Resumo: O clareamento dental é uma das técnicas estéticas mais conservadoras no âmbito odontológico. Técnicas de clareamento caseira ou de consultório, ou a combinação de ambas, podem ser realizadas para alcançar o prognóstico desejado. Esta combinação das técnicas pode ser realizada por meio de diferentes abordagens, como apresentado neste relato de casos. Um paciente do gênero masculino, 24 anos de idade, cujos incisivo central superior e caninos apresentaram cor A2 e A3, respectivamente, foi inicialmente submetido à técnica de clareamento de consultório de duração por sessão de 40 min com gel de peróxido de hidrogênio (PH) 40%, em duas sessões. Posteriormente, a técnica de clareamento caseira foi realizada durante duas semanas com gel de peróxido de carbamida (PC) a uma concentração de 10%. A cor final dos incisivos centrais superiores foi alterada para 1M1, enquanto a dos caninos para B1. Por outro lado, uma paciente do gênero feminino, 30 anos de idade, foi inicialmente submetida ao clareamento caseiro (durante a noite, por 4 semanas- PC 10%), e subsequentemente, com aplicação pontual de agente clareador (PH 35%) em consultório, em sessão de 45 min. A cor do incisivo central superior da paciente, mudou de A1 para 0,5M1 e do canino superior de A3 para B1. As escalas de cor VITA Classic (A1-D4), e VITA Bleachedguided 3D-MASTER foram utilizadas nesta série de casos para determinar a coloração dos dentes durante os tratamentos. Em ambos os tratamentos, os pacientes não reportaram sensibilidade dental significativa, e suas expectativas finais foram atendidas.

Palavras-chave: Clareamento dental, peróxido de hidrogênio, peróxido de carbamida, combinação.

Summary: Tooth bleaching is one of the most conservative aesthetic techniques. At-home and in-office bleaching may be performed, as well as the combination of both techniques. As this combination may be done in different manners, distinct orders of combinations were proposed in this report. A 24 years-old man, whose upper central incisor and canine shades were A2 and A3 respectively, was initially treated by chair-side with 40-min application of 40% hydrogen peroxide (HP) for two sessions. Afterwards, an overnight two-weeks at-home bleaching was performed with 10% carbamide peroxide (CP). The shade of upper central incisors changed to 1M1 and canine was B1. Conversely, a 30 years-old woman was firstly conducted with at-home bleaching (overnight 4-weeks 10% CP) and, subsequently, with a single appointment of in-office bleaching for 45 minutes (35% HP). The shade of her upper central incisor changed from A1 to 0.5M1 and the upper canine from A3 to B1. The VITA classical (A1-D4) shade guide and VITA Bleached guide 3D-MASTER were used to determine the tooth color during the treatments. In both bleaching treatments, patients reported no significant tooth sensitivity and the final outcomes met their expectation.

Key words: Tooth Bleaching, Hydrogen Peroxide, Carbamide Peroxide, Combination.

Sumário:

1. Introdução.....	9
2. Artigo.....	10
2.1 Combination of at-home and in-office bleaching techniques: case series/ Combinação das técnicas de clareamento dental caseira e de consultório: relato de casos.....	10
3. Conclusão.....	24
Referências.....	24
Anexos.....	28
Anexo 1	28
Anexo 2	29
Anexo 3.....	30

1. Introdução

Um aumento da apreciação da Odontologia estética pelos pacientes vem sendo muito observada nos dias atuais, estendendo-se desde a reabilitação oral, até por pequenos tratamentos estéticos que utilizam técnicas minimamente invasivas como o clareamento dental. Este último, considerado uma técnica extremamente conservativa e segura (1). O clareamento dental em dentes vitais pode ser realizado tanto no consultório odontológico, como pelo próprio paciente em casa. Mesmo assim, a combinação de ambas as técnicas é uma medida rotineiramente adotada por dentistas, e comumente utilizada com a finalidade de aumentar a efetividade clareadora (2).

Enquanto os géis clareadores de peróxidos de carbamida e hidrogênio em baixas concentrações são utilizados no período noturno por várias semanas e aplicado pelo próprio paciente em casa, altas concentrações desses agentes clareadores nos géis permitem que as aplicações realizadas no consultório odontológico pelo cirurgião- dentista apresentem resultados finais satisfatórios, em menor tempo (3). Mesmo assim, o clareamento caseiro, o qual também deve ser supervisionado pelo dentista, tende a causar uma menor sensibilidade dental, visto sua baixa concentração do agente ativo. De qualquer forma, a técnica utilizada em consultório é amplamente utilizada na rotina clínica, de modo que promove uma alteração de cor dental de forma acelerada (4). Além disso, o tratamento em consultório permite ao paciente que não se adapta tão facilmente com a moldeira permanecer com uma técnica odontológica conservadora, ao invés de ser diretamente submetido a tratamentos mais invasivos, como restaurações diretas e indiretas (5).

Embora alguns estudos reportem a eficácia de ambas as técnicas clareadoras utilizadas de forma combinada e separada (4,5,6), a literatura aponta que o tratamento clareador utilizando a combinação das duas técnicas existentes- caseira e de consultório- pode aumentar a eficácia clareadora (6), como também aumentar a sensibilidade dental pós tratamento (6,8). Mesmo assim, os estudos são inconclusivos, e avanços sobre a combinação das técnicas são baseados no aumento da eficácia, como também na estabilidade de cor. Visto por outro ângulo, a motivação do paciente também pode aumentar, já que a mudança de cor pode ser vista mais rapidamente (9). Apesar disso, fatores como a severidade e origem da pigmentação

dental, espessura dentinária, idade, comprometimento e expectativa dos pacientes devem ser levado em conta na decisão para a escolha das técnicas combinadas (3).

Em vista destes fatos, como também na heterogeneidade de aspectos clínicos observados nos pacientes que procuram procedimentos estéticos, a combinação das técnicas clareadoras se mostra como uma alternativa viável quando o tratamento inicial proposto não atende as expectativas finais do paciente. Desta forma, a comparação da combinação das técnicas em diferentes sequências é reportada nestes relatos de casos.

2. Artigo: Trabalho aceito para a publicação na Revista Gaúcha de Odontologia

2.1. Title: Combination of at-home and in-office bleaching techniques: case series/ Combinação das técnicas de clareamento dental caseira e de consultório: relato de casos.

Short Title: Bleaching combination techniques/ Combinação de técnicas de clareamento dental

Resumo: O clareamento dental é uma das técnicas estéticas mais conservadoras no âmbito odontológico. Técnicas de clareamento caseira ou de consultório, ou a combinação de ambas, podem ser realizadas para alcançar o prognóstico desejado. Esta combinação das técnicas pode ser realizada por meio de diferentes abordagens, como apresentado neste relato de casos. Um paciente do gênero masculino, 24 anos de idade, cujos incisivo central superior e caninos apresentaram cor A2 e A3, respectivamente, foi inicialmente submetido à técnica de clareamento de consultório de duração por sessão de 40 min com gel de peróxido de hidrogênio (PH) 40%, em duas sessões. Posteriormente, a técnica de clareamento caseira foi realizada durante duas semanas com gel de peróxido de carbamida (PC) a uma concentração de 10%. A cor final dos incisivos centrais superiores foi alterada para 1M1, enquanto a dos caninos para B1. Por outro lado, uma paciente do gênero feminino, 30 anos de idade, foi inicialmente submetida ao clareamento caseiro (durante a noite, por 4 semanas- PC 10%), e subsequentemente, com aplicação pontual de agente clareador (PH 35%) em consultório, em sessão de 45 min. A cor do incisivo central superior da paciente, mudou de A1 para 0,5M1 e do canino superior de A3 para B1. As escalas de cor VITA Classic (A1-D4), e VITA Bleachedguided 3D-MASTER foram utilizadas nesta série de casos para determinar a coloração dos dentes durante os tratamentos. Em ambos os

tratamentos, os pacientes não reportaram sensibilidade dental significativa, e suas expectativas finais foram atendidas.

Palavras-chave: Clareamento dental, peróxido de hidrogênio, peróxido de carbamida, combinação.

Summary: Tooth bleaching is one of the most conservative aesthetic techniques. At-home and in-office bleaching may be performed, as well as the combination of both techniques. As this combination may be done in different manners, distinct orders of combinations were proposed in this report. A 24 years-old man, whose upper central incisor and canine shades were A2 and A3 respectively, was initially treated by chair-side with 40-min application of 40% hydrogen peroxide (HP) for two sessions. Afterwards, an overnight two-weeks at-home bleaching was performed with 10% carbamide peroxide (CP). The shade of upper central incisors changed to 1M1 and canine was B1. Conversely, a 30 years-old woman was firstly conducted with at-home bleaching (overnight 4-weeks 10% CP) and, subsequently, with a single appointment of in-office bleaching for 45 minutes (35% HP). The shade of her upper central incisor changed from A1 to 0.5M1 and the upper canine from A3 to B1. The VITA classical (A1-D4) shade guide and VITA Bleached guide 3D-MASTER were used to determine the tooth color during the treatments. In both bleaching treatments, patients reported no significant tooth sensitivity and the final outcomes met their expectation.

Key words: Tooth Bleaching, Hydrogen Peroxide, Carbamide Peroxide, Combination.

Introduction

As an overwhelming appreciation of Cosmetic Dentistry is observed among patients in recent days, starting oral rehabilitation or minor aesthetic treatment by a minimally invasive procedure as tooth bleaching is considered a suitable conservative approach [1]. Tooth bleaching in vital teeth may be performed either at home or in dental office. However, combination of both techniques is a prevalent measure taken by the dentists, commonly used as an attempt to enhance the bleaching effectiveness [2].

While carbamide and hydrogen peroxides under low concentrations are normally tray-delivered overnight during few weeks, higher concentrated bleaching agents are applied by the dentist for a shorter period of time for some appointments [3]. Thus, at-home bleaching, which must be supervised by the dentist, is prone to cause less tooth

sensitivity. However, in-office therapy has been largely used in clinical routine since it provides faster color alteration [4] . Furthermore, it provides an opportunity to the patient not comfortable to use the tray to still be treated with a conservative approach instead of being directly submitted to more invasive treatments, e.g. direct and indirect restorations [5] .

Although some studies reported the efficacy of both at-home and in-office techniques separately or combined [4,6,7], literature has pointed out that combined bleaching may enhance bleaching efficacy [6] and increase tooth sensitivity [6,8]. Even though studies are inconclusive, advantages of the combined techniques are based on the boost of the bleaching efficacy and the improvement of color stability. Also, motivation of patients might increase since the color change would be visible more immediately [9]. Nevertheless, factors such as severity and origin of tooth staining, dentin thickness, age, commitment and expectation of the patients should also be taken into account in the decision for combined bleaching [3] .

In view of these facts, and also of the heterogeneous clinical aspects that may be found among patients looking for aesthetic procedures, combination of techniques appears as a feasible alternative when the initial proposed treatment does not meet the expected outcome. Hence, a comparison of bleaching combination in different approaches is reported in a case series.

Cases Report

Case 1

A 24 years-old man sought for the clinical facilities of Piracicaba Dental School with the main complaint of discolored teeth. The patient had not previously undergone to tooth bleaching. After dental prophylaxis and clinical examination, patient was considered eligible for bleaching therapy. Considering VITA classical (A1-D4) shade guide (VITA Zahnfabrik, Germany), initial shade of upper central incisor and canine showed shade A2 (Figure 1A) and A3 (Figure 1B), respectively. After two sessions with two 20-min 40% hydrogen peroxide (HP) (Opalescence Boost 40% - Ultradent, USA) application (Figure 2A), shades of upper central incisor and canine diminished to A1 (Figure 1C) and A2 (Figure 1D), respectively. As the patient's expectation of canine shades was not met, at-home bleaching was performed with 10% carbamide peroxide (CP) (Whiteness Perfect 10%, FGM, Joinville, SC, Brazil) for 4 hours a day during 2 weeks. An individual acetate tray was made and instructions of gel application

from right to left premolar in both arches were prescribed (Figure 2B). Seven days elapsed from the end of the treatment, upper central incisor (Figure 1E) showed shade 1M1 and upper canine, B1 (Figure 1F).




Figure 1: Treatment starting by in-office technique followed by at-home bleaching.

A- Initial color aspect, in which upper central incisor showed shade A2; **B-** The upper canine showed shade A3; **C-** Following the in-office bleaching procedures, upper central incisor shade changed to A1; **D-** Upper canine showed shade A2 after in-office bleaching; **E-** At the end of the combined bleaching, central incisor showed shade 1M1; **F-** The canine showed shade B1 following the combined technique.



Figure 2: Bleaching procedures performed during the first case. **A-** Application of the in-office bleaching gel with an adequate gingival barrier; **B-** Individual acetate tray made for at-home bleaching technique.

The shade of incisor was measured by means of a bleached shade guide that ranges from 0M1 to 5M3 (VITA Bleached guide 3D-MASTER, VITA Zahnfabrik, Germany). The Figure 3 showed the initial and final patients aspect, respectively (Figure 3A-B).



Figure 3: Clinical picture of the first patient before and after combined bleaching started by in-office technique. **A-** Initial color aspect of the patient showing good oral health, but teeth with yellow appearance. **B-** final clinical aspect, in which the health of the patient was maintained and whiter teeth, was detected.

Case 2

The shades of upper central incisor and canine of a 30 years-old woman, who has also never undergone tooth bleaching, showed A1 (Figure 4A) and A3 (Figure 4B). Initially, 10% CP at-home bleaching was indicated for 2 weeks with supervision from operators. After seven days, shades A1 and A2 were shown for upper incisor and canine (Figure 4C and 4D). Finally, after two weeks of at-home bleaching, upper incisor and canine showed 1M1 and B1, respectively (Figure 4E and 4F).




Figure 4; Clinical evaluation of the second patient at the first phase of the bleaching treatment (at-home technique). **A-** Initial upper central incisor showing shade A1; **B-** Initially, upper canine showed shade A3; **C-** Shade A1 was shown for upper central incisor after 1 week; **D-** Upper canine showed shade A2 after 1 week; **E-** After 2 weeks, central incisor showed shade 1M1; **F-** The canine showing shade B1;

The patient reported that the higher discoloration observed in upper canine (B1) and lower incisors (A1) in comparison to upper incisors (1M1) after the at-home bleaching did not meet her expectations (Figure 5A-C). In view of these facts, in-office application of 35% HP [3 x 15 min (Whiteness HP, FGM, Joinville, SC, Brazil)] was performed only on the teeth considered discolored by the patient (Figure 5D). Seven days elapsed from professional HP application, shades of incisor and canine showed 0,5M1 and B1, respectively (Figure 5E and 5F), and the lower incisors showed B1 (Figure 5G).




Figure 5: The second phase of the bleaching therapy applied in the second patient (in-office technique). **A** – Upper central incisor showing shade 1M1 after 2-week at-home bleaching; **B**- Shade B1 was shown for upper canine; **C**- Lower central incisors showed shade A1 after at-home bleaching during 2 weeks; **D**- In-office bleaching technique only on upper Canines and lower teeth; **E**- Following the in-office bleaching, central incisor reduced to 0.5M1; **F**- Final upper canine was maintained in B1; **G**- Lower incisors showed shade B1 after the in-office bleaching appointment.

As in the previous case, the shade of incisor was measured by means of a bleached shade guide (VITA Bleachedguide 3D-MASTER, VITA Zahnfabrik, Germany). The initial and final patients aspect is observed at Figure 6, respectively (Figure 6A and 6B).



Figure 6: Clinical picture of the second patient before and after combined bleaching started by at-home technique. **A-** Starting aspect of the patient presenting acceptable oral health and yellow appearance of the teeth. The colors of upper and lower arches were unpaired. **B-** Clinical aspect after the bleaching therapy with whiter teeth and improved color harmony among the arches.

Discussion

In these case series, bleaching was planned based on the information collected during initial interview of both patients. Upon the first appointment, the male patient reported a very important academic presentation due the following two weeks. Since the patient desired whiter teeth for this upcoming event and short time was available, two in-office bleaching appointment were performed [4]. As a further in-office session was not possible, two-week at-home bleaching was determined so that the expectations could be meet. Even though studies in literature are controversial regarding additional color change promoted by combination of bleaching techniques, Vaez et al. (2018) reported that a single preliminary session of in-office bleaching with 35% HP enhanced color change for patients submitted to at-home therapy with 10% CP [10]. However, it is important to take into account that CP's application regimen for our patient was four times greater, which not ensures that combination would be beneficial for color change

in other patients. Although the variety of concentration and time of gel's applications hampers the interpretation of bleaching combination efficacy [11], the first clinical case showed an adequate final result.

Conversely, bleaching on the female patient started by at-home treatment due to her preference. However, after 4 weeks of CP overnight application, both patient and professionals noticed that lower arch and upper canines did not follow the same color change pattern as upper incisors. Therefore, an additional in-office session with high-concentrated HP was performed to provide faster result than at-home bleaching [4]. There is no data in the literature investigating the difference among upper and lower teeth on bleaching efficacy. Effect of peroxides is believed to be a result of HP's free radicals interaction with dentin chromogens [3]. However, the enamel/dentin penetration of hydrogen peroxide is inversely proportional to the diffusion distance [12]. Therefore, it could be hypothesized that free radicals of HP pass through lower dentin faster than in the upper, thereby diminishing the breakdown of staining molecules of dentin. As other factors could have influenced this clinical aspect such as periodicity of CP's application by the patient or the stability of the bleaching gel, research on this topic are paramount to determine the influence of teeth's thickness for bleaching.

While bleaching combination presents long-term clinical stability [13], a recent clinical trial revealed no difference among at-home or in-office and combined techniques [8]. A recent systematic review with meta-analysis revealed that there was no statistical difference between the techniques regarding color change. Nevertheless, the authors pointed out that these data should be cautiously evaluated due to the wide variation of gels application times both chair-side and at-home [11]. Furthermore, there is little information regarding which sequence of bleaching combination would produce higher color change. Most studies performed an initial in-office session followed by at-home therapy for few weeks, which is known as "jump start" technique [14]. However, the opposite way could also be investigated once clinical situation such as in the second case might overcome color discrepancy among arches faster due to higher concentration of in-office bleaching gels [3].

Initial color evaluation of both cases was performed using a classical visual guide shade. However, the final color measurement of central incisor was determined by means of a bleached-guide once the tooth was incompatible with B1, the highest value found in classical guides [7]. Although the same was not observed for upper canine, B1 was reached in both case series. Even though randomized controlled clinical trials are

more indicated to determine comparison of both combinations' efficacy [4,7], the two techniques herein adopted were efficient to produce color change on upper canines, regardless of the age of patients. As luminosity and yellowness of teeth are lower and higher, respectively, in older patients [15], longer regimen would be necessary to breakdown the chromospheres.

Given the combination of in-office and at-home bleaching, an *in vivo* study showed that this combination is beneficial for the patient and the dentist owing to the synergy among both techniques that potentiates shade alteration and, consequently, shorten the time of treatment [11]. Nonetheless, individual risk of tooth sensitivity must be considered when combining in-office and at-home bleaching. Zhao et al. (2013) indicated *in vivo* that at-home application of low-concentrated gel is likely to enhance tooth permeability, which, in turn, increased high-concentrated HP penetration towards dental tissues. Thus, the authors stated that such sequence of bleaching combination should be avoided in patients presenting teeth's morphology alteration and hypersensitivity [15]. As the volunteers of our cases presented teeth with no major implications, no contraindication was found for the combination of at-home and in-office bleaching whatsoever. Indeed, the patients reported a minor tooth sensitivity only during the day of HP application but the pain did not last any longer, which is in accordance with the literature [16].

Tooth sensitivity as a response of tooth bleaching may be related to the presence of hydrogen peroxide presence in pulp tissue after gel's application [14,17]. It has been recently demonstrated that the pH of high-concentrated HP gels plays a role in peroxide penetration through tooth structures towards pulp chamber [18]. Balladares et al. (2019) showed that high-concentrated HP, which was used in the first case, led to the same intrapulpal concentration of HP under different application techniques (1 x 45 min or 3 x 15 min). However, three 15-min application of the same 35% HP used in the second case resulted in significant lower penetration. The pH of both bleaching gels decreased only during 1 x 45 min application [19]. Even though availability of hydrogen peroxide is still high even after a single 45-min gel application [20], caution is still necessary when deciding to replenish or not peroxide gels since a variety of products with different and unknown pH values are commercialized [21]. To date, there is a lack of data regarding the intrapulpal penetration of low-concentrated agents and combined techniques.

Despite the fact that information in the literature is missing to indicate the sequence of bleaching techniques combination, the combinations presented in these

case series were efficient to meet the expectations of the patients. According to the previously evidences discussed, combination of techniques should be indicated when supported by an adequate investigation of color aspect and expectations of each patient.

Acknowledgements:

The authors thank the patients for giving permission for publishing this case, and thank also the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior-Brasil (CAPES- Finance code 001) for the support in this case report.

REFERENCES

1. Silva FBD, Chisini LA, Demarco FF, Horta BL, Correa MB. Desire for tooth bleaching and treatment performed in Brazilian adults: findings from a birth cohort. *Braz Oral Res.* 2018 Mar 8;32:e12. doi:10.1590/1807-3107bor-2018.vol32.0012.
2. Carey CM. Tooth whitening: What we now know. *J Evid Based Dent Pract.* 2014;14(SUPPL.):70–6.
3. Kwon SR, Wertz PW. Review of the mechanism of tooth whitening. *J Esthet Restor Dent.* 2015 Sep-Oct; 27(5):240-57
4. Lima SNL, Ribeiro IS, Grisotto MA, Fernandes ES, Hass V, deJesus Tavares RR, Pinto SCS, Lima DM, Loguercio AD, Bandeca MC. Evaluation of several clinical parameters after bleaching with hydrogen peroxide at different concentrations: A randomized clinical trial. *J Dent.* 2018 Jan; 68:91-97. doi: 10.1016/j.jdent.2017.11.008.
5. Baratieri L, Monteiro Jr. S. Clareamento Dental. In: Guanabara Koogan, editor. *Odontologia Restauradora: Fundamentos e Possibilidades.* São Paulo; 2ª Ed. 2015. p. 832.
6. Machado LS, Anchieta RB, dos Santos PH, Briso AL, Tovar N, Janal MN, Coelho PG, Sundfeld RH. Clinical Comparison of At-Home and In-Office Dental Bleaching Procedures: A Randomized Trial of a Split-Mouth Design. *Int J Periodontics Restorative Dent.* 2016 Mar-Apr;36(2):251-60. doi: 10.11607/prd.2383

7. Loguercio AD, Servat F, Stanislawczuk R, Rezende M. Effect of acidity of in-office bleaching gels on tooth sensitivity and whitening: a two-center double-blind randomized clinical trial. *Clin Oral Investig*. 2017 Dec; 21(9):2811-2818. doi: 10.1007/s00784-017-2083-5.
8. Dourado Pinto AV, Carlos NR, Amaral FLBD, França FMG, Turssi CP, Basting RT. At-home, in-office and combined dental bleaching techniques using hydrogen peroxide: Randomized clinical trial evaluation of effectiveness, clinical parameters and enamel mineral content. *Am J Dent*. 2019 Jun; 32(3):124-132.
9. Loguercio AD, Martins LM, Silva LM, Reis A. In-Office Whitening. In: Perdigão J. *Tooth Whitening An Evidence-Based Perspective* 2016; 1a ed. Switzerland: 145-67
10. Vaez SC, Correia A, Santana TR, Santana M, Peixoto AC, Leal PC, Faria-E-Silva AL. Is a Single Preliminary Session of In-office Bleaching Beneficial for the Effectiveness of At-home Tooth Bleaching? A Randomized Controlled Clinical Trial. *Oper Dent*. 2019 Jul/Aug; 44(4):E180-E189. doi: 10.2341/18-196-C.
11. de Geus J, Wambier L, Kossatz S, Loguercio A, Reis A. At-home vs In-office Bleaching: A Systematic Review and Meta-analysis. *Oper Dent*. 2016; 41(4):341–56. doi: 10.2341/15-287-LIT.
12. de Oliveira Duque CC, Soares DG, Basso FG, Hebling J, de Souza Costa CA. Influence of enamel/dentin thickness on the toxic and esthetic effects of experimental in-office bleaching protocols. *Clin Oral Investig*. 2017 Nov; 21(8):2509-2520. doi: 10.1007/s00784-017-2049-7. Epub 2017 Jan 14.
13. Rezende M, Ferri L, Kossatz S, Loguercio AD, Reis A. Combined Bleaching Technique Using Low and High Hydrogen Peroxide In-Office Bleaching Gel. *Oper Dent*. 2016 Jul-Aug; 41(4): 388-96. doi: 10.2341/15-266-C.
14. Briso ALF, Rahal V, Galinari MO, Soares DG, Souza Costa CG. Complication from the Use of Peroxides. In: Perdigão J. *Tooth Whitening An Evidence-Based Perspective*

2016; 1a ed. Switzerland: 45-80.

15. Zhao K, Zong L, Zhang Q, Att W. Clinical comparison between two bleaching techniques: a 180-day follow-up study. *Quintessence Int.* 2013;44(8):601-7. doi:10.3290/j.qi.a29702.

16. Kwon SR. Tooth Whitening: How Does It Work. In: Perdigão J. *Tooth Whitening-a Evidence-Based Perspective* 2016; 1a ed. Switzerland: 21-34

17. Cavalli V, Silva BG, Berger SB, Abuna G, Marson FC, Tabchoury C, Giannini M. Effect of Adhesive Restoration and Bleaching Technique on the Concentration of Hydrogen Peroxide In the Pulp Chamber. *Oper Dent.* 2017 Mar/Apr;42(2):E44-E54. doi: 10.2341/16-079-L.

18. Acuña ED, Parreiras SO, Favoreto MW, Cruz GP, Gomes A, Borges CPF, Loguercio AD, Reis A. In-office bleaching with a commercial 40% hydrogen peroxide gel modified to have different pHs: Color change, surface morphology, and penetration of hydrogen peroxide into the pulp chamber. *J Esthet Restor Dent.* 2019 Feb 20. doi: 10.1111/jerd.12453. [Epub ahead of print].

19. Balladares L, Alegría-Acevedo LF, Montenegro-Arana A, Arana-Gordillo LA, Pulido C, Salazar-Gracez MT, Reis A, Loguercio AD. Effects of pH and Application Technique of In-office Bleaching Gels on Hydrogen Peroxide Penetration into the Pulp Chamber. *Oper Dent.* 2019 Apr 22. doi: 10.2341/18-148-L. [Epub ahead of print]

20. Marson FC, Gonçalves RS, Silva CO, Cintra LT, Pascotto RC, Santos PH, Briso AL. Penetration of hydrogen peroxide and degradation rate of different bleaching products. *Operative Dentistry* 2015; 40(1): 72-9. doi: 10.2341/13-270-L.

21. Cavalli V, Silva BGD, Berger SB, Marson FC, Tabchoury CPM, Giannini M. Decomposition Rate, pH, and Enamel Color Alteration of At-Home and In-Office Bleaching Agents. *Braz Dent J.* 2019 Jul 22;30(4):385-396. doi:10.1590/0103-6440201902484.

3. Conclusão

Apesar das lacunas presentes na literatura sobre a combinação das técnicas de clareamento dental utilizando diferentes sequências, os relatos de casos presente trabalho demonstra que a combinação de técnicas em diferentes sequências foram de encontro com as expectativas dos pacientes. De acordo com as evidências discutidas anteriormente, a combinação de técnicas de clareamento dental podem ser indicadas, visto que são suportadas por investigação adequada tanto para o aspecto de cor dental, como em relação expectativa de ambos pacientes.

Referências:

1. Silva FBD, Chisini LA, Demarco FF, Horta BL, Correa MB. Desire for tooth bleaching and treatment performed in Brazilian adults: findings from a birth cohort. *Braz Oral Res.* 2018 Mar 8;32:e12. doi:10.1590/1807-3107bor-2018.vol32.0012.
2. Carey CM. Tooth whitening: What we now know. *J Evid Based Dent Pract.* 2014;14(SUPPL.):70–6.
3. Kwon SR, Wertz PW. Review of the mechanism of tooth whitening. *J Esthet Restor Dent.* 2015 Sep-Oct; 27(5):240-57
4. Lima SNL, Ribeiro IS, Grisotto MA, Fernandes ES, Hass V, deJesus Tavares RR, Pinto SCS, Lima DM, Loguercio AD, Bandeca MC. Evaluation of several clinical parameters after bleaching with hydrogen peroxide at different concentrations: A randomized clinical trial. *J Dent.* 2018 Jan; 68:91-97. doi: 10.1016/j.jdent.2017.11.008.
5. Baratieri L, Monteiro Jr. S. Clareamento Dental. In: Guanabara Koogan, editor. *Odontologia Restauradora: Fundamentos e Possibilidades.* São Paulo; 2ª Ed. 2015. p. 832.
6. Machado LS, Anchieta RB, dos Santos PH, Briso AL, Tovar N, Janal MN, Coelho PG, Sundfeld RH. Clinical Comparison of At-Home and In-Office Dental Bleaching Procedures: A Randomized Trial of a Split-Mouth Design. *Int J Periodontics Restorative Dent.* 2016 Mar-Apr;36(2):251-60. doi: 10.11607/prd.2383

7. Loguercio AD, Servat F, Stanislawczuk R, Rezende M. Effect of acidity of in-office bleaching gels on tooth sensitivity and whitening: a two-center double-blind randomized clinical trial. *Clin Oral Investig*. 2017 Dec; 21(9):2811-2818. doi: 10.1007/s00784-017-2083-5.

8. Dourado Pinto AV, Carlos NR, Amaral FLBD, França FMG, Turssi CP, Basting RT. At-home, in-office and combined dental bleaching techniques using hydrogen peroxide: Randomized clinical trial evaluation of effectiveness, clinical parameters and enamel mineral content. *Am J Dent*. 2019 Jun; 32(3):124-132.

9. Loguercio AD, Martins LM, Silva LM, Reis A. In-Office Whitening. In: Perdigão J. *Tooth Whitening An Evidence-Based Perspective* 2016; 1a ed. Switzerland: 145-67

10. Vaez SC, Correia A, Santana TR, Santana M, Peixoto AC, Leal PC, Faria-E-Silva AL. Is a Single Preliminary Session of In-office Bleaching Beneficial for the Effectiveness of At-home Tooth Bleaching? A Randomized Controlled Clinical Trial. *Oper Dent*. 2019 Jul/Aug; 44(4):E180-E189. doi: 10.2341/18-196-C.

11. de Geus J, Wambier L, Kossatz S, Loguercio A, Reis A. At-home vs In-office Bleaching: A Systematic Review and Meta-analysis. *Oper Dent*. 2016; 41(4):341–56. doi: 10.2341/15-287-LIT.

12. de Oliveira Duque CC, Soares DG, Basso FG, Hebling J, de Souza Costa CA. Influence of enamel/dentin thickness on the toxic and esthetic effects of experimental in-office bleaching protocols. *Clin Oral Investig*. 2017 Nov; 21(8):2509-2520. doi: 10.1007/s00784-017-2049-7. Epub 2017 Jan 14.

13. Rezende M, Ferri L, Kossatz S, Loguercio AD, Reis A. Combined Bleaching Technique Using Low and High Hydrogen Peroxide In-Office Bleaching Gel. *Oper Dent*. 2016 Jul-Aug; 41(4): 388-96. doi: 10.2341/15-266-C.

* De acordo com as normas da UNICAMP/FOP, baseadas na padronização do International Committee of Medical Journal Editors - Vancouver Group. Abreviatura dos periódicos em conformidade com o PubMed.

14. Briso ALF, Rahal V, Galinari MO, Soares DG, Souza Costa CG .Complication from the Use of Peroxides. In: Perdigão J. Tooth Whitening An Evidence-Based Perspective 2016; 1a ed. Switzerland: 45-80.
15. Zhao K, Zong L, Zhang Q, Att W. Clinical comparison between two bleaching techniques: a 180-day follow-up study. Quintessence Int. 2013;44(8):601-7. doi:10.3290/j.qi.a29702.
16. Kwon SR. Tooth Whitening: How Does It Work. In: Perdigão J. Tooth Whitening-a Evidence-Based Perspective 2016; 1a ed. Switzerland: 21-34
17. Cavalli V, Silva BG, Berger SB, Abuna G, Marson FC, Tabchoury C, Giannini M. Effect of Adhesive Restoration and Bleaching Technique on the Concentration of Hydrogen Peroxide In the Pulp Chamber. Oper Dent. 2017 Mar/Apr;42(2):E44-E54.doi: 10.2341/16-079-L.
18. Acuña ED, Parreiras SO, Favoreto MW, Cruz GP, Gomes A, Borges CPF, Loguercio AD, Reis A. In-office bleaching with a commercial 40% hydrogen peroxide gel modified to have different pHs: Color change, surface morphology, and penetration of hydrogen peroxide into the pulp chamber. J Esthet Restor Dent. 2019 Feb 20. doi: 10.1111/jerd.12453. [Epub ahead of print].
19. Balladares L, Alegría-Acevedo LF, Montenegro-Arana A, Arana-Gordillo LA, Pulido C, Salazar-Gracez MT, Reis A, Loguercio AD. Effects of pH and Application Technique of In-office Bleaching Gels on Hydrogen Peroxide Penetration into the Pulp Chamber. Oper Dent. 2019 Apr 22. doi: 10.2341/18-148-L. [Epub ahead of print]
20. Marson FC, Gonçalves RS, Silva CO, Cintra LT, Pascotto RC, Santos PH, Briso AL. Penetration of hydrogen peroxide and degradation rate of different bleaching products. Operative Dentistry 2015; 40(1): 72-9. doi: 10.2341/13-270-L.
21. Cavalli V, Silva BGD, Berger SB, Marson FC, Tabchoury CPM, Giannini M. Decomposition Rate, pH, and Enamel Color Alteration of At-Home and In-Office Bleaching Agents. Braz Dent J. 2019 Jul 22;30(4):385-396. doi:10.1590/0103-

6440201902484.

Anexos:

1. Carta de aceite do artigo para a publicação na Revista Gaúcha de Odontologia.

RGO - Revista Gaúcha de Odontologia

Decision Letter (RGO-2019-0149)

From: biagomes@yahoo.com

To: marcela.a.ferretti@gmail.com, baguiar@unicamp.br

CC:

Subject: RGO - Revista Gaúcha de Odontologia - Decision on Manuscript ID RGO-2019-0149

Body: 19-Feb-2020

Dear Dr. Ferretti:

Manuscript ID RGO-2019-0149 entitled "Combination of at-home and in-office bleaching techniques: case series" which you submitted to the RGO - Revista Gaúcha de Odontologia, has been reviewed. The comments of the reviewer(s) are included at the bottom of this letter.

Based on the evaluation of our reviewer(s) and on the recommendation of our Associate Editor, the manuscript received the Final Decision "[Major Review]".

This means that the manuscript has potential for publication, but some corrections should be made. This decision still does not qualify the manuscript as "Accepted". Therefore, I invite you to respond to the reviewer(s)' comments and revise your manuscript.

To revise your manuscript, log into <https://mc04.manuscriptcentral.com/rgo-scielo> and enter your Author Center, where you will find your manuscript title listed under "Manuscripts with Decisions." Under "Actions," click on "Create a Revision." Your manuscript number has been appended to denote a revision.

You may also click the below link to start the revision process (or continue the process if you have already started your revision) for your manuscript. If you use the below link you will not be required to login to ScholarOne Manuscripts.

*** PLEASE NOTE: This is a two-step process. After clicking on the link, you will be directed to a webpage to confirm. ***

https://mc04.manuscriptcentral.com/rgo-scielo?URL_MASK=4c7331e5933d40d290f142beef070878

You will be unable to make your revisions on the originally submitted version of the manuscript. Instead, revise your manuscript using a word processing program and save it on your computer. Please also highlight the changes to your manuscript within the document by using the track changes mode in MS Word or by using bold or colored text.

2. Termo de Consentimento Livre e Esclarecido dos pacientes dos relatos de casos.

FORMULÁRIO DE CONSENTIMENTO DO PACIENTE

Nome da pessoa descrita no artigo ou mostrada na fotografia: _____

Assunto da fotografia ou do manuscrito: Procedimento clínico de clareamento dental

Título do artigo: **Combination of at-home and in-office bleaching techniques: case series**

Autor para correspondência: **Marcela Alvarez Ferretti**

Eu, Lucas Vinícius Domingues, RG nº 489502131-3, residente
à Rua Cardel Arcobrende nº 230,
Complemento: Ap 105, Bairro: Pinheiros, na cidade de
São Paulo, paciente (ou responsável legal de: _____), por meio
deste **Termo de Consentimento Livre e Esclarecido**, consinto que o Dr.
Marcela Alvarez Ferretti tire fotografias, faça vídeos e outros tipos
de imagens minhas, sobre o meu caso clínico. Consinto que estas imagens sejam
utilizadas para finalidade didática e científica, divulgadas em aulas, palestras,
conferências, cursos, congressos etc., e publicadas em livros, artigos, portais de
internet, revistas científicas e similares, podendo inclusive ser mostrado o meu rosto, o
que pode fazer com que eu (ou ele) seja reconhecido.

Consinto também que sejam utilizadas e divulgadas as imagens de meus exames, como
radiografias, tomografias computadorizadas, ressonâncias magnéticas, ultrassons,
eletromiografias, histopatológicos (exame no microscópio da peça cirúrgica retirada) e
outros.

Este consentimento pode ser revogado, sem qualquer ônus ou prejuízo à minha pessoa,
a meu pedido ou solicitação, desde que a revogação ocorra antes da publicação.

Fui esclarecido de que não receberei nenhum ressarcimento ou pagamento pelo uso
das minhas imagens e também compreendi que o Dr. **Marcela Alvarez Ferretti** e a
equipe de profissionais que me atende e atenderá durante todo o tratamento não terá
qualquer tipo de ganho financeiro com a exposição da minha imagem nas referidas
publicações.

Assinatura do paciente ou responsável: Lucas Vinícius Domingues
Data: 29/10/2019

FORMULÁRIO DE CONSENTIMENTO DO PACIENTE

Nome da pessoa descrita no artigo ou mostrada na fotografia:

Ana Paula Bettoni Teles

Assunto da fotografia ou do manuscrito: Procedimento clínico de clareamento dental

Título do artigo: **Combination of at-home and in-office bleaching techniques: case series**

Autor para correspondência: _____

Eu, Ana Paula Bettoni Teles, RG nº 1738353-6, residente à Rua Fernando Febeliano da Costa nº 1818, Complemento: _____, Bairro: Alemães, na cidade de Piracicaba, paciente (ou responsável legal de: _____), por meio

deste **Termo de Consentimento Livre e Esclarecido**, consinto que o Dr.

_____ tire fotografias, faça vídeos e outros tipos de imagens minhas, sobre o meu caso clínico. Consinto que estas imagens sejam utilizadas para finalidade didática e científica, divulgadas em aulas, palestras, conferências, cursos, congressos etc., e publicadas em livros, artigos, portais de internet, revistas científicas e similares, podendo inclusive ser mostrado o meu rosto, o que pode fazer com que eu (ou ele) seja reconhecido.

Consinto também que sejam utilizadas e divulgadas as imagens de meus exames, como radiografias, tomografias computadorizadas, ressonâncias magnéticas, ultrassons, eletromiografias, histopatológicos (exame no microscópio da peça cirúrgica retirada) e outros.

Este consentimento pode ser revogado, sem qualquer ônus ou prejuízo à minha pessoa, a meu pedido ou solicitação, desde que a revogação ocorra antes da publicação.

Fui esclarecido de que não receberei nenhum ressarcimento ou pagamento pelo uso das minhas imagens e também compreendi que o Dr. **Matheus Kury Rodrigues** e a equipe de profissionais que me atende e atenderá durante todo o tratamento não terá qualquer tipo de ganho financeiro com a exposição da minha imagem nas referidas publicações.

Assinatura do paciente ou responsável: Ana Paula Bettoni Teles

Data: 22/10/2019

Versatilidade das técnicas de clareamento: Combinação de técnicas de clareamento caseiro e de consultório- relato de casos

RELATÓRIO DE ORIGINALIDADE

10%	9%	8%	%
ÍNDICE DE SEMELHANÇA	FONTES DA INTERNET	PUBLICAÇÕES	DOCUMENTOS DOS ALUNOS

FONTES PRIMÁRIAS

1	eprints.unm.ac.id Fonte da Internet	4%
2	repositorio.unicamp.br Fonte da Internet	2%
3	link.springer.com Fonte da Internet	1%
4	www.sbpqo.org.br Fonte da Internet	1%
5	"Tooth Whitening", Springer Science and Business Media LLC, 2016 Publicação	1%
6	"Full Issue PDF", Operative Dentistry, 2017 Publicação	1%

Excluir citações

Desligado

Excluir

< 1%

Excluir bibliografia

Em

correspondências