



UNIVERSIDADE ESTADUAL DE CAMPINAS
FACULDADE DE ODONTOLOGIA DE PIRACICABA

DAYANE JAQUELINE GROSS

**ESTUDO MORFOLÓGICO E MORFOMÉTRICO DAS
VARIAÇÕES ANATÔMICAS DO VENTRE ANTERIOR DO
MÚSCULO DIGÁSTRICO**

MORPHOLOGICAL AND MORPHOMETRIC STUDY OF THE
ANATOMICAL VARIATIONS OF ANTERIOR BELLY OF
DIGASTRIC MUSCLE

PIRACICABA

2022

DAYANE JAQUELINE GROSS

**ESTUDO MORFOLÓGICO E MORFOMÉTRICO DAS
VARIAÇÕES ANATÔMICAS DO VENTRE ANTERIOR DO
MÚSCULO DIGÁSTRICO**

**MORPHOLOGICAL AND MORPHOMETRIC STUDY OF THE
ANATOMICAL VARIATIONS OF ANTERIOR BELLY OF
DIGASTRIC MUSCLE**

Dissertação apresentada à Faculdade de Odontologia de Piracicaba da Universidade Estadual de Campinas como parte dos requisitos exigidos para obtenção do título de Mestra em Biologia Buco-Dental, na Área de Anatomia

Dissertation presented to the Piracicaba Dental School of the University of Campinas in partial fulfillment of the requirements for the degree of MSc in Dental Biology in the Anatomy area.

Orientador: Prof. Dr. Alexandre Rodrigues Freire

ESTE EXEMPLAR CORRESPONDE À VERSÃO FINAL DA DISSERTAÇÃO DEFENDIDA PELA ALUNA DAYANE JAQUELINE GROSS E ORIENTADA PELO PROF. DR. ALEXANDRE RODRIGUES FREIRE.

PIRACICABA

2022

Ficha catalográfica
Universidade Estadual de Campinas
Biblioteca da Faculdade de Odontologia de Piracicaba
Marilene Girello - CRB 8/6159

G914e Gross, Dayane Jaqueline, 1991-
Estudo morfológico e morfométrico das variações anatômicas do centro anterior do músculo digástrico / Dayane Jaqueline Gross. – Piracicaba, SP : [s.n.], 2022.

Orientador: Alexandre Rodrigues Freire.
Dissertação (mestrado) – Universidade Estadual de Campinas, Faculdade de Odontologia de Piracicaba.

1. Variação anatômica. 2. Músculos. I. Freire, Alexandre Rodrigues, 1985-. II. Universidade Estadual de Campinas. Faculdade de Odontologia de Piracicaba. III. Título.

Informações Complementares

Título em outro idioma: Morphological and morphometric study of the anatomic variations of anterior belly of digastric muscle

Palavras-chave em inglês:

Anatomic variation

Muscles

Área de concentração: Anatomia

Titulação: Mestra em Biologia Buco-Dental

Banca examinadora:

Alexandre Rodrigues Freire [Orientador]

Cristhiane Martins Schmidt

Eduardo Daruge Junior

Data de defesa: 27-09-2022

Programa de Pós-Graduação: Biologia Buco-Dental

Identificação e informações acadêmicas do(a) aluno(a)

- ORCID do autor: <https://orcid.org/0000-0001-6642-4672>

- Currículo Lattes do autor: <http://lattes.cnpq.br/2260347245738959>



UNIVERSIDADE ESTADUAL DE CAMPINAS
Faculdade de Odontologia de Piracicaba

A Comissão Julgadora dos trabalhos de Defesa de Dissertação de Mestrado, em sessão pública realizada em 27 de setembro de 2022, considerou a candidata DAYANE JAQUELINE GROSS aprovada.

PROF. DR. ALEXANDRE RODRIGUES FREIRE

PROF^a. DR^a. CRISTHIANE MARTINS SCHMIDT

PROF. DR. EDUARDO DARUGE JUNIOR

A Ata da defesa, assinada pelos membros da Comissão Examinadora, consta no SIGA/Sistema de Fluxo de Dissertação/Tese e na Secretaria do Programa da Unidade.

DEDICATÓRIA

A todos professores com os quais aprendi a nobreza do que é o ensino.
À minha família que é a base da minha vida.
Ao meu noivo que sempre esteve ao meu lado.

AGRADECIMENTOS

À Universidade Estadual de Campinas, na pessoa do Magnífico Reitor Prof. Dr. Antônio José de Almeida Meirelles.

À Faculdade de Odontologia de Piracicaba, na pessoa do Senhor Diretor, Prof. Dr. Flávio Henrique Baggio Aguiar.

A Coordenadoria de Pós-Graduação, na pessoa do Senhor Coordenador Prof. Dr. Valentim Adelino Ricardo Barão.

À Equipe Técnica da Coordenadoria de Pós-Graduação, em nome da senhora Ana Paula Carone Gonzales, Érica A. Pinho Sinhoreti, Raquel Quintana Marcondes Cesar, Claudinéia Prata Pradella e do senhor Leandro Viganó, sempre dispostos a ajudar.

Ao programa de pós-graduação em Biologia Buco-Dental, na figura do coordenador Prof. Dr. Marcelo Rocha Marques.

Ao meu orientador Prof. Dr Alexandre Rodrigues Freire, que em nenhum momento me deixou desamparada e trouxe seu ensino e ajuda com maestria e zelo, dando assistência e ajuda maior do que eu pudesse imaginar, meu muito obrigada.

À Prof^a Dr.^a Ana Cláudia Rossi, que me acompanhou durante todo o mestrado, onde muitos momentos com toda a sua ajuda, me fez parecer que fosse a minha orientadora. Sua ajuda fez muita diferença.

Aos meus pais, por acreditarem sempre em mim e me incentivarem ao estudo.

Ao meu companheiro Rudgero, que sempre foi paciente e amoroso em todos os momentos que precisei estar ausente.

À minha amiga Patrícia que sempre me incentivou a buscar a melhor formação.

À Beatriz e a Luciane, que ajudaram em toda pesquisa, agradeço a toda dedicação.

A todos familiares e amigos que de forma direta e indireta colaboraram para a realização deste estudo.

RESUMO

O músculo digástrico pertence a um grupo de músculos na região submental denominados supra-hioideos. Estudos têm relatado variabilidade sem tamanho, forma, envolvendo principalmente seu ventre anterior. Entretanto, tais características não têm sido detalhadamente descritas e, devido às diferenças em suas variações, é necessária sua classificação. O objetivo do trabalho foi avaliar a morfologia e obter dados morfométricos das variações anatômicas do ventre anterior do músculo digástrico. Foram analisadas 31 peças cadavéricas (cabeças humanas) do laboratório de Anatomia da FOP-UNICAMP. A morfologia do ventre anterior do músculo digástrico foi avaliada de forma observacional, identificando as possíveis variações anatômicas que foram caracterizadas e classificadas quanto a quantidade de ventres musculares, direcionamento das fibras, local de origem e local de inserção. As medidas morfométricas foram verificadas utilizando um paquímetro digital, em que foram obtidas as medidas de largura e comprimento do ventre muscular. Foi realizada a incidência das variações anatômicas através do cálculo da porcentagem das peças que apresentarem determinada variação anatômica. A análise estatística foi realizada no software GraphPAD Prism v.8 (San Diego, CA, EUA). Foi considerado um nível de significância de 5%. Foi realizada uma estatística descritiva (em %) para determinar a incidência das variações anatômicas do ventre anterior do músculo digástrico. Dos 31 cadáveres avaliados, a variação anatômica do ventre anterior do músculo digástrico esteve presente em 6 (19, 31%; 1 feminino e 5 masculinos). Todas as variações anatômicas apresentaram um ventre acessório ao ventre anterior do músculo digástrico. Entretanto, estes ventres acessórios se configuraram de forma diferente na localização, direção das fibras musculares e nas suas dimensões (comprimento e largura). Conclui-se que das variações encontradas, mais presente é o ventre acessório. A presença e o conhecimento anatômico dessa e das demais variações se faz importante, para o auxílio em procedimentos cirúrgicos; patológicos ou de função diagnóstica.

Palavras-chave: variação anatômica; músculo digástrico; ventre muscular.

ABSTRACT

The digastric muscle belongs to a group of muscles in the submental region called suprahyoid muscles. Studies have reported variability in size, shape, mainly involving its anterior belly. However, such characteristics have not been described in detail and, due to the differences in their variations, their classification is necessary. The objective of this work was to evaluate the morphology and obtain morphometric data of the anatomical variations of the anterior belly of the digastric muscle. Thirty-one cadaveric parts (human heads) from the Anatomy laboratory at FOP-UNICAMP were analyzed. The morphology of the anterior belly of the digastric muscle was evaluated observationally, identifying the possible anatomical variations that were characterized and classified according to the amount of muscle bellies, fiber direction, place of origin and insertion site. The morphometric measurements were verified using a digital caliper, in which the measurements of width and length of the muscle belly were obtained. The incidence of anatomical variations was performed by calculating the percentage of pieces that presented a certain anatomical variation. Statistical analysis was performed using GraphPAD Prism v.8 software (San Diego, CA, USA). A significance level of 5% was considered. Descriptive statistics (in %) were performed to determine the incidence of anatomical variations of the anterior belly of the digastric muscle. Of the 31 cadavers evaluated, the anatomical variation of the anterior belly of the DM was present in 6 (19, 31%; 1 female and 5 male). All anatomical variations presented an accessory belly to the anterior belly of the MD. However, these accessory bellies were configured differently in the location, direction of muscle fibers and in their dimensions (length and width). It is concluded that the incidence of the variations found is the presence of an accessory belly. The presence and anatomical knowledge of this and other variations is important to assist in surgical procedures, pathological or diagnostic function.

Keywords: anatomical variation; digastric muscle; muscle belly.

SUMÁRIO

1.INTRODUÇÃO	10
2.ARTIGO	12
3.CONCLUSÃO	24
REFERÊNCIAS	25
ANEXOS	27
ANEXO 1: Certificação do Comitê de Ética em Pesquisa	27
ANEXO 2: Comprovante de Aceite no Periódico	28
ANEXO 3: Comprovante de originalidade e anti-plágio	29
ANEXO 4: Comprovante de autorização para publicação do artigo	30

1. INTRODUÇÃO

O músculo digástrico (MD) localiza-se na região submentual e é um músculo que possui dois ventres, denominados ventre anterior e ventre posterior, separados por um tendão intermediário, o qual forma uma ancoragem no osso hioide. Sua origem caracteriza-se pela fixação do ventre posterior na incisura mastoidea, na base do crânio, e sua inserção se localiza na fossa digástrica, na base da mandíbula na região do mento. Por possuírem origens embriológicas distintas, seus ventres possuem inervação e vascularização diferentes, sendo que o ventre anterior é innervado pelo nervo milohioideo, ramo do nervo mandibular. A artéria e a veia submentuais, ramos da artéria e veia faciais, são responsáveis por sua vascularização (Aszmann et al., 1998). O ventre anterior situa-se superficialmente ao músculo milohioideo (Hollinshead, 1961) e sua função é elevar o osso hioide quando os músculos infra-hioideos estão relaxados e abaixar a mandíbula quando o osso hioide está fixo (Reyes et al., 2007). Além disso, deprime a mandíbula para auxiliar a mastigação e a fala, e para estabilizar o osso hioide durante a deglutição e a fala (Coquerelle et al., 2017).

O MD apresenta uma grande variabilidade de tamanho e forma. Essas variações envolvem principalmente seu ventre anterior, caracterizadas pelas alterações no número de ventres anteriores ou pela presença de feixes acessórios originários de um tendão comum ou do tendão intermediário (Celik et al., 2002; Mangalagiri & Razvi, 2009). As variações do ventre anterior podem apresentar-se uni ou bilaterais e mais raramente sendo simétricas, estendendo-se para a mandíbula ou para o músculo milohioideo ou ainda cruzando a linha média para o músculo milohioideo oposto ou para a fossa digástrica oposta. Esses feixes ou deslizamentos musculares unilaterais ou bilaterais podem ocorrer como resultado de uma redução filogenética do MD ou de um desenvolvimento incomum do material ontogenético (Atamaz & Govsa, 2005).

Na literatura, algumas variações já foram descritas. Turan-ozdemir et al. (2004) observaram um ventre anterior acessório bilateral do digástrico. Aktekin et al. (2003) relataram uma variação bilateral e simétrica do ventre anterior do digástrico em que os feixes acessórios estavam dispostos em cruz, superficiais ao músculo milohioideo. Traini (1983) relatou um caso de músculos digástrico acessórios bilaterais. Reyes et al. (2007) observaram ventres anteriores acessórios bilaterais formando uma

configuração triangular, onde ambos os ventres correm do tendão intermediário e se inserem sobre a rafe milohioidea. Zuhail et al. (2007) relataram um caso de quadrificação bilateral do ventre anterior do músculo digástrico.

O conhecimento da variabilidade do MD é de suma importância durante a mobilização do retalho miocutâneo do músculo platísmo em procedimentos reconstrutivos (Turan-ozdemir et al., 2004). O conhecimento aprofundado da anatomia típica e variável dos ventres anteriores é de suma importância devido ao uso muscular após procedimentos de paralisia do nervo facial e retalho da artéria submentual (Uzun et al, 2001). Além disso, o ventre anterior é utilizado como ponto de referência durante a identificação do nervo lingual ou do ducto da glândula submandibular (Norton, 1991).

Outras implicações cirúrgicas desta variante podem ser encontradas na cirurgia plástica e reconstrutiva. Atamaz-Pinar et al. (2005), descreveram sobre o uso de retalho submental para reconstrução de defeitos intraorais ou na face. O MD também pode ser usado para restaurar a função depressora do lábio inferior resultante de uma paralisia do ramo marginal mandibular do nervo facial, onde o tendão intermediário do MD poderia ser suturado ao ângulo lateral do lábio inferior para restaurar a simetria perdida após a paralisia. Um músculo anormal ou com variações nessa região pode dificultar a retirada do retalho e a transferência do feixe muscular (Tan, 2002).

As áreas submentual e submandibular são muito bem vascularizadas, e o conhecimento de suas anormalidades musculares é essencial quando um retalho miocutâneo é mobilizado em cirurgia plástica reconstrutiva. Variações vasculares podem coexistir e, para manter e aumentar a viabilidade do retalho, o padrão de ramificação dos vasos deve ser bem preservado no intraoperatório (Notsis et al., 2020). Assim casos de variações nessa região devem ser cuidadosamente identificadas para evitar interpretações errôneas e auxiliar no correto procedimento cirúrgico ou estético, além de contribuir para o ensino da Anatomia.

Nesta dissertação, o objetivo do estudo realizado foi identificar os tipos de variações anatômicas do ventre anterior do músculo digástrico e as caracterizá-las morfometricamente.

2. ARTIGO

Title: Morphological study of the anatomical variations of anterior belly of digastric muscle in Brazilian cadavers*

Running head: Anatomical variations of digastric muscle

***O presente estudo foi aceito para publicação no periódico internacional “*Folia Morphologica*” (ANEXO 2).**

Authors: Dayane Jaqueline Gross¹, Ana Cláudia Rossi¹, Beatriz Carmona Ferreira-Pileggi¹, Luciane Naomi Oguma Watanabe¹, Paulo Roberto Botacin², Felipe Bevilacqua Prado¹, Alexandre Rodrigues Freire¹

1- Department of Biosciences, Anatomy Division, Piracicaba Dental School, University of Campinas, Piracicaba, São Paulo, Brazil.

2- Basic Sciences Department, Araçatuba Dental School, Paulista State University – UNESP, Araçatuba, São Paulo, Brazil.

Address for correspondence to:

Prof. Alexandre Rodrigues Freire

PhD, Researcher, Department of Biosciences, Anatomy Division, Piracicaba Dental School, University of Campinas, Piracicaba, São Paulo, Brazil.

Av. Limeira, 901- Areião, Piracicaba - SP – Brazil 13.414-903

E-mail: alerfreire@gmail.com

+55 19 21065721

ABSTRACT

Background: Cases of variations in anterior belly of the digastric muscle must be carefully identified to avoid misinterpretations and assist in the correct surgical or aesthetic procedure and help in the teaching of Anatomy. The aim of this study was to describe the anatomical variations of anterior belly of digastric muscle in Brazilian cadavers. **Materials and methods:** Thirty-one human heads were selected, from adult cadavers (18 to 80 years, 29 males and 2 females). The morphology of the anterior belly of the digastric muscle was observed, identifying the possible anatomical variations that were characterized and classified according to the amount of muscle bellies, fiber direction and place of origin and insertion. The morphometric measurements were performed using a digital caliper. To analyze the data obtained, photographic documentation, anatomical description and individual morphometric description of each muscle belly were performed. The incidence of anatomical variation was obtained in percentage (%). **Results:** The anatomical variation of the anterior belly of the digastric muscle was present in 6 cadavers (19.31%; 1 female and 5 male). All anatomical variations presented an accessory belly to the anterior belly. However, these accessory bellies were configured differently in the location, direction of muscle fibers and in their dimensions (length and width). **Conclusions:** The gross anatomy of the anterior belly of the DM and their variations are important to assist in surgical procedures, pathological or diagnostic function. In addition, asymmetrical variations in the submental region must be carefully identified to avoid misinterpretations.

Keywords: anatomy, digastric muscle, anatomical variations, morphology.

INTRODUCTION

The digastric muscle (DM) is in the submental region and is a muscle that has two bellies, called the anterior belly and the posterior belly, separated by an intermediate tendon, which forms an anchorage in the hyoid bone. Its origin is characterized by the attachment of the posterior belly to the mastoid notch, at the base of the skull, and its insertion is in the digastric fossa, at the base of the mandible in the chin region. Because they have different embryological origins, their bellies have different innervation and vascularization, and the anterior belly is innervated by the mylohyoid nerve, a branch of the mandibular nerve. The submental artery and vein, branches of the facial artery and vein, are responsible for its vascularization [2]. The anterior belly lies superficial to the mylohyoid muscle [6] and its function is to elevate the hyoid bone when the infrahyoid muscles are relaxed and to lower the mandible when the hyoid bone is fixed [13].

The DM shows great variability in size, shape, and number. These variations mainly involve the anterior belly of the muscle and the number of anterior bellies or the presence of accessory bundles originating from a common tendon or the intermediate tendon [4, 9]. The anterior belly variations can be unilateral or bilateral and more rarely being symmetrical, extending to the mandible or the mylohyoid muscle or even crossing the midline to the opposite mylohyoid or the opposite digastric fossa [3].

DM variability is of paramount importance during mobilization of the platysma myocutaneous flap in reconstructive procedures [15]. In-depth knowledge of the typical and variable anatomy of the anterior bellies is of paramount importance due to muscle use after facial nerve paralysis and submental artery flap procedures [16]. In addition, the anterior belly is used as a reference point when identifying the lingual nerve or submandibular gland duct [12].

The submental and submandibular areas are very well vascularized, and knowledge of their muscle abnormalities is essential when a myocutaneous flap is mobilized in reconstructive plastic surgery. Vascular variations can coexist and, to maintain and increase the viability of the flap, the branching pattern of the vessels must be well preserved intraoperatively [10]. Thus, cases of variations in this region must be carefully identified to avoid misinterpretations and assist in the correct surgical or aesthetic procedure and help in the teaching of Anatomy.

The aim of this study was to describe the anatomical variations of anterior belly of DM in Brazilian cadavers.

MATERIALS AND METHODS

Ethics approval

The research was analyzed and approved by the Committee of Research Ethics of the University of Campinas (Protocol number CEP-FOP-UNICAMP-CAAE 59065622.6.0000.5418) (ANEXO 1).

Sample

Thirty-one human heads were selected, embedded cadaver, belonging to the Laboratory of Anatomy of the Piracicaba Dental School from University of Campinas (FOP-UNICAMP). The heads were from adult cadavers, selected at random ranging in age from 18 to 80 years and of both sexes (29 males and 2 females).

The inclusion criteria were that cadaveric specimens from adult individuals were used, showing the preservation of anatomical structures in the submental region, especially the suprahyoid muscles, among which the DM is included. The exclusion criteria were that cadaveric specimens that do not show the preservation of the suprahyoid muscles in the submental region, or that present any pathology that have the presence of a foreign body in the submental region, interfering with the anatomy of the suprahyoid muscles, were excluded.

Calibration

Initially, an anatomical study of the submental region was performed to identify the anatomy of the suprahyoid muscles and adjacent anatomical structures. After this study, the examiner identified the anterior belly of the DM and calibrated the measurements to be performed using a digital caliper (Mitutoyo®).

Evaluation of the Morphology and the Morphometry of the anterior belly of the DM

The morphology of the anterior belly of the DM was observed, identifying the possible anatomical variations that were characterized and classified according to the amount of muscle bellies, fiber direction and place of origin and insertion.

The morphometric measurements were performed using a digital caliper (Mitutoyo®), in which the measurements of the width (mm) and length (mm) of the anterior belly were obtained to add to the morphological descriptions.

Data analysis

To analyze the data obtained, photographic documentation, anatomical description and individual morphometric description of each muscle belly were performed. The incidence of anatomical variation was obtained in percentage (%).

RESULTS

Of the 31 cadavers evaluated, the anatomical variation of the anterior belly of the DM was present in 6 (19.31%; 1 female and 5 male). All anatomical variations presented an accessory belly to the anterior belly of the DM. However, these accessory bellies were configured differently in the location, direction of muscle fibers and in their dimensions (length and width). Anatomical variations are described below. The occurrence of each type can be seen in Figure 1.




Figure 1. Photographs of the submental region with emphasis on the location of the anatomical variation of the anterior bellies of digastric muscle. Different colors delimit the belly area. 1: main anterior belly. Lowercase letters: accessory bellies.

In Figure 1, A and B show an anatomical variation of the anterior belly of the DM with the appearance of a unilateral accessory minor belly, in which its muscle fibers originate from the intermediate tendon of the DM and are obliquely directed to insert into the mylohyoid muscle raphe together with mylohyoid muscle fibers. In the total of variations found, 3 heads presented this configuration. In the heads that were found in this configuration, the accessory belly (1a) on the left side (figure 1A) had a length equal to 26.36 mm and a width equal to 6 mm. As for the main bellies, the DM anterior belly on the right side had a length of 52.62 mm and a width of 16.22 mm. And the DM anterior belly on the left side had a length equal to 54.27 mm and width equal to 20.94 mm. In figure 1-B, the accessory belly (1a) was located on the right side, with a length equal to 26.70 mm and width equal to 12.47 mm. As for the main bellies, the DM anterior belly on the right side had a length equal to 43.43 mm and a width equal to 11.42 mm. And the DM anterior belly on the left side had a length equal to 33.35 mm and width equal to 15.08 mm.

In the figure 1C, a more complex configuration can be seen, in which the variation occurred bilaterally. The accessory bellies in this configuration originate in the intermediate tendon of the DM and their fibers are directed obliquely across the midsagittal plane until they reach their point of insertion at the base of the mandible on the opposite side in an extension of the digastric fossa. Both accessory bellies are superficial in relation to the mylohyoid muscle. The accessory belly 1a has a length equal to 48.16 mm and a width equal to 12.54 mm. And the accessory belly 1b has a length equal to 39.16 mm and a width equal to 8.63 mm. As for the main bellies, the DM anterior belly on the right side had a length equal to 40.23 mm and a width equal to 14.26 mm. And the DM anterior belly on the left side had a length equal to 41.16 mm and width equal to 16.11 mm.

In the figure 1D, the variation was presented with the presence of accessory bellies on the right and left sides. Muscle fibers originate from the intermediate tendon of the DM and insert into the extension of the digastric fossa ipsilaterally. However, a small number of fibers can be observed originating from the intermediate tendon on the opposite side, crossing the midline in a similar way to that found in item C of figure 1. The accessory bellies are found superficially to the mylohyoid muscle. The accessory belly 1a is 26.53 mm (length) and 6.92 mm (width). And the accessory belly 1b has a length equal to 23.49 mm and a width equal to 4.99 mm. As for the main

bellies, the DM anterior belly on the right side had a length equal to 30.17 mm and a width equal to 11.58 mm. And the DM anterior belly on the left side had a length equal to 35.68 mm and a width equal to 8.96 mm.

Considering the variations found, the configuration found in item E of figure 1 was the most complex. In this situation, the variation appeared bilaterally, with different accessory bellies on each side. In addition, the main anterior belly on the left side also showed an anatomical variation.

On the left side, the main anterior belly originates from the intermediate tendon, but its fibers are directed obliquely to insert into the base of the mandible, right in the midsagittal plane. In addition, part of its fibers has insertion in the mylohyoid muscle raphe. The accessory belly on the left, at its origin, presents fibers together with the main anterior belly. However, when approaching the midsagittal plane, the group of fibers separates to insert into the mylohyoid muscle raphe. Thus, the accessory belly on the left is located posterior to the main anterior belly. On the right side, the main anterior belly did not show variation as on the left side. The accessory belly presented a different configuration from the other variations that were found. Its fibers originate in the intermediate tendon and are directed anteriorly to insert into the digastric fossa together with the main anterior belly. This accessory belly is located laterally in relation to the main anterior belly and is separated from it by a fascia. It is also observed that at its insertion, the accessory belly on this side presented a longer tendon in relation to the main anterior belly.

In the case E of figure 1, the anterior belly 1a had a length equal to 24.79 mm and width equal to 12.94 mm. And the anterior belly 1b had a length equal to 33.03mm and width equal to 12.94mm. As for the main bellies, the DM anterior belly on the right side (blue) had a length equal to 34.02 mm and a width equal to 14.13 mm. And the MD anterior belly on the left side (red) had a length equal to 40.94 mm and width equal to 12.20 mm.

DISCUSSION

DM variability is of paramount importance during mobilization of the platysma myocutaneous flap in reconstructive procedures [15]. In-depth knowledge of the typical and variable anatomy of the anterior bellies is of paramount importance due to muscle use after facial nerve paralysis and submental artery flap procedures [16]. In addition,

the anterior belly is used as a reference point when identifying the lingual nerve or submandibular gland duct [12].

The knowledge of the anatomical variations of the anterior belly of the DM is also necessary for professionals who perform conservative and surgical treatments in the submental region, as well as in teaching the anatomy of this structure. Therefore, knowing the anatomical variations can facilitate diagnosis and prevent possible complications during surgical procedures [13].

The variations may also be responsible for esthetic alterations in the submental region. For example, a bulky anterior belly of the digastric muscle can contribute to an unsightly contour of the neck by giving a bulge in the submental area. By removing a good portion of the anterior belly muscle (i.e., making the anterior belly thinner), the bulge in the submental area can be minimized, thus giving the patient a better contour of the neck [5].

Regarding the variations found, ethnicity as the most frequent factor in the occurrence of these variations. Literary data show a higher prevalence of these variations in the Asian population, especially the Japanese. While in the general population a rate of 31.4% was found, in the Asian population the rate was 51.7% for the occurrence of anatomical variation of the DM [1].

Another factor responsible for variations, such as bilateral duplication of the anterior belly, may be the embryological origin, which is explained by a deficiency in the differentiation of the mesoderm of the first pharyngeal arch on both sides or by abnormal migration of the cells of the pharyngeal arch neural crest. In relation to the crossing of muscle fibers in the midline, it may be due to the proximity of the pharyngeal arches [8].

The DM is an important muscle in the neck, which pulls the mandible down to open the mouth and elevates the hyoid bone for stabilization during swallowing and is also involved in chewing and speaking. In the functional aspect, the changes may affect some DM functions. For example, in crossed anterior bellies, the difficulty of opening the mouth is suggestive. Probably, the crossing of the bellies makes it difficult for the mandible to be moved downwards.

In the supernumerary bundles, due to their proximity to the anterior belly of the digastric and mylohyoid muscles, additional force is likely to be provided to these muscles during mastication [14]. In the case of the additional belly or unilateral

accessory, an imbalance can occur when elevating the hyoid bone or lowering the mandible [11]. These unilateral or bilateral muscle bundles or slips may occur because of a phylogenetic reduction of the DM or an unusual development of ontogenetic material [3].

Notably, knowing the variations found, clinicians and surgeons can make use of and must be very careful in interpreting radiographs and CT scans, as the variations may appear like soft tissue masses, especially in the unilateral type [13].

CONCLUSION

The gross anatomy of the anterior belly of the DM and their variations is important to assist in surgical procedures, pathological or diagnostic function. In addition, asymmetrical variations in the submental region must be carefully identified to avoid misinterpretations.

ACKNOWLEDGMENTS

The authors sincerely thank those who donated their bodies to science so that anatomical research could be performed. Results from such research can potentially increase mankind's overall knowledge that can then improve patient care. Therefore, these donors and their families deserve our highest gratitude [7].

The authors are grateful to the technician of the Anatomy Laboratory, sr. Cristiano Manoel, for performing the gross anatomic dissections.

REFERENCES

1. Arayapisit T, Vorakulpipat C, Srimaneekarn N et al. Anatomical Variations of the Anterior Belly of the Digastric Muscle in Thai Cadavers: A Cross-Sectional Study. *J Int Soc Prev Community Dent.* 2022 Apr 8;12(2):171-177. doi: 10.4103/jispcd.JISPCD_188_21.
2. Aszmann OC, Ebmer JM, Dellon AL. The anatomic basis for the innervated mylohyoid/digastric flap in facial reanimation. *Plast Reconstr Surg.* 1998;102(2):369-72. doi: 10.1097/00006534-199808000-00010
3. Atamaz Pinar Y, Govsa F, Bilge O. The anatomical features and surgical usage of the submental artery. *Surg Radiol Anat.* 2005 Aug;27(3):201-5. doi: 10.1007/s00276-005-0317-8.

4. Celik HH, Aldur MM, Ozdemir B, Akşit MD. Abnormal digastric muscle with unilateral quadrification of the anterior belly. *Clin Anat.* 2002 Jan;15(1):32-4. doi: 10.1002/ca.1088.
5. Connell BF, Shamoun JM. The significance of digastric muscle contouring for rejuvenation of the submental area of the face. *Plast Reconstr Surg.* 1997 May;99(6):1586-90.
6. Hollinshead WH. The Jaws, Palate and Tongue, In: *Anatomy for Surgeons.* vol 1, 3rd print, A Hoeber – Harper Book, New York. 1961:340-400.
7. Iwanaga J, Singh V, Ohtsuka A, Hwang Y, Kim HJ, Morys J, Ravi KS, Ribatti D, Trainor PA, Sañudo JR, Apaydin N, Şengül G, Albertine KH, Walocha JA, Loukas M, Duparc F, Paulsen F, Del Sol M, Addis P, Hegazy A, Tubbs RS. Acknowledging the use of human cadaveric tissues in research papers: Recommendations from anatomical journal editors. *Clin Anat.* 2021 Jan;34(1):2-4. doi: 10.1002/ca.23671.
8. Kim SD, Loukas M. Anatomy and variations of digastric muscle. *Anat Cell Biol.* 2019 Mar;52(1):1-11. doi: 10.5115/acb.2019.52.1.1.
9. Mangalagiri AS, Razvi MR. Variations in the anterior belly of digastric. *Int J Health Sci (Qassim).* 2009; 3: 257–62.
10. Natsis K, Piagkou M, Lazaridis N, Anastasopoulos N. Asymmetry of the Accessory Anterior Digastric Muscle Bellies: The Clinical Significance in Facial and Neck Surgery. *Cureus.* 2020 Feb 29;12(2):e7148. doi: 10.7759/cureus.7148.
11. Netter FH. Neck. In: Hansen JT, Benninger B, Brueckner JK, Carmichael SW, Granger NA, Tubbs RS, editors. *Atlas of Human Anatomy.* 5th ed. Philadelphia: Elsevier; 2011. p.26-34.
12. Norton MR. Bilateral accessory digastric muscles. *Br J Oral Maxillofac Surg.* 1991; 29: 167–68. doi: 10.1016/0266- 4356(91)90030-9.
13. Reyes G, Contreras C, Ramírez LM, Ballesteros LE. The digastric muscle's anterior accessory belly: case report. *Med Oral Patol Oral Cir Bucal.* 2007 Sep 1;12(5):E341-3. PMID: 17767094.
14. Sowman PF, Flavel SC, McShane CL et al. Asymmetric activation of motor cortex controlling human anterior digastric muscles during speech and target- directed jaw movements. *J Neurophysiol.* 2009;102:159-66. doi: 10.1152/jn.90894.2008.

15. Turan-Ozdemir S, Oygucu IH, Kafa IM. Bilateral abnormal anterior bellies of digastric muscles. *Anat Sci Int.* 2004 Jun;79(2):95-7. doi: 10.1111/j.1447-073x.2004.00065x.
16. Uzun A, Aluclu A, Kavakli A. Bilateral accessory anterior bellies of the digastric muscle and review of the literature. *Auris Nasus Larynx.* 2001 Apr;28(2):181-3. doi: 10.1016/s0385-8146(00)00100-0.

3. CONCLUSÃO

A partir dos resultados observado neste estudo, pode-se concluir que o ventre anterior do MD apresenta diferentes formas de variações anatômicas. Tais variações ocorrem com o aparecimento de ventre musculares acessórios, que podem apresentar comprimento e largura diferentes, assim como diferenças na direção das fibras musculares.

REFERÊNCIAS*

- Aszmann OC, Ebmer JM, Dellon AL. The anatomic basis for the innervated mylohyoid/digastric flap in facial reanimation. *Plast Reconstr Surg*. 1998;102(2):369-72.
- Atamaz Pinar Y, Govsa F, Bilge O. The anatomical features and surgical usage of the submental artery. *Surg Radiol Anat*. 2005; 27: 201–205. doi: 10.1007/s00276-005-0317-8.
- Atkekin M, Kurtoglu Z, Ozturk AH. A bilateral and symmetrical variation of the anterior belly of the digastric muscle. *Acta Med Okayama*. 2003; 57: 205 - 207.
- Celik HH, Aldur MM, Ozdemir B, Akşit MD. Abnormal digastric muscle with unilateral quadrification of the anterior belly. *Clin Anat*. 2002; 15: 32–34. doi: 10.1002/ca.1088.
- Coquerelle M, Prados-Frutos J, Rojo R, et al. The fetal origin of the human chin. *Evol Biol*. 2017; 44(3): 295–311. doi: 10.1007/s11692-017-9408-9. [L]
[SEP]
- Hollinshead WH. The Jaws, Palate and Tongue, In: *Anatomy for Surgeons*. vol 1, 3rd print, A Hoeber – Harper Book, New York. 1961:340-400.
- Mangalagiri AS, Razvi MR. Variations in the anterior belly of digastric. *Int J Health Sci (Qassim)*. 2009; 3: 257–262.
- Natsis K, Piagkou M, Lazaridis N, Anastasopoulos N. Asymmetry of the Accessory Anterior Digastric Muscle Bellies: The Clinical Significance in Facial and Neck Surgery. *Cureus*. 2020 Feb 29;12(2):e7148. doi: 10.7759/cureus.7148.
- Norton MR. Bilateral accessory digastric muscles. *Br J Oral Maxillofac Surg*. 1991; 29: 167–68. doi: 10.1016/0266-4356(91)90030-9.
- Ozgur Z, Govsa F, Ozgur T. The cause of the difference in the submental region: aberrant muscle bundles of the anterior belly of the digastric muscle. *J Craniofac Surg*. 2007 Jul;18(4):875-81. doi: 10.1097/scs.0b013e31806844da.
- Reyes G, Contreras C, Ramirez LM, Ballesteros LE. The digastric muscle's anterior accessory belly: Case report. *Med Oral Patol Oral Cir Bucal*. 2007;12: E341-343.
- Tan ST. Anterior belly of digastric muscle transfer: a useful technique in head and neck surgery. *Head Neck*. 2002; 24: 947–954. doi: 10.1002/hed.10150.

* De acordo com as normas da UNICAMP/FOP, baseadas na padronização do International Committee of Medical Journal Editors - Vancouver Group. Abreviatura dos periódicos em conformidade com o PubMed

Traini M. Bilateral accessory digastric muscles. *Surgical and Radiologic Anatomy*. 1983; 5: 199 - 200.

Turan-Ozdemir S, Oygucu IH, Kafa IM. Bilateral abnormal anterior bellies of digastric muscles. *Anat Sci Int*. 2004 Jun;79(2):95-7. doi: 10.1111/j.1447-073x.2004.00065.x

Uzun A, Aluclu A, Kavakli A. Bilateral accessory anterior bellies of the digastric muscle and review of the literature. *Auris Nasus Larynx*. 2001; 28: 181–183. doi: 10.1016/S0385-8146(00)00100-0.

ANEXOS

ANEXO 1: Certificação do Comitê de Ética em Pesquisa

	UNICAMP - FACULDADE DE ODONTOLOGIA DE PIRACICABA DA UNIVERSIDADE DE CAMPINAS - FOP/UNICAMP	
---	---	---

PARECER CONSUBSTANCIADO DO CEP

DADOS DO PROJETO DE PESQUISA

Título da Pesquisa: Estudo morfológico e morfométrico das variações anatômicas do ventre anterior do músculo digástrico

Pesquisador: DAYANE JAQUELINE GROSS

Área Temática:

Versão: 2

CAAE: 59065622.6.0000.5418

Instituição Proponente: Faculdade de Odontologia de Piracicaba - Unicamp

Patrocinador Principal: Financiamento Próprio

DADOS DO PARECER

Número do Parecer: 5.475.994

	UNICAMP - FACULDADE DE ODONTOLOGIA DE PIRACICABA DA UNIVERSIDADE DE CAMPINAS - FOP/UNICAMP	
---	---	--

Continuação do Parecer: 5.475.994

Tipo Documento	Arquivo	Postagem	Autor	Situação
Informações Básicas do Projeto	PB_INFORMAÇÕES_BÁSICAS_DO_PROJETO_1948449.pdf	19/06/2022 15:16:04		Aceito
Outros	Respostaparecer.pdf	19/06/2022 15:15:17	DAYANE JAQUELINE GROSS	Aceito
Projeto Detalhado / Brochura Investigador	Projeto.pdf	19/06/2022 15:13:54	DAYANE JAQUELINE GROSS	Aceito
Outros	Autarquivo.pdf	26/05/2022 21:31:20	DAYANE JAQUELINE GROSS	Aceito
TCLE / Termos de Assentimento / Justificativa de Ausência	TCLE.pdf	26/05/2022 21:30:13	DAYANE JAQUELINE GROSS	Aceito
Declaração de Pesquisadores	DeclaraPesquisadores.pdf	26/05/2022 21:29:59	DAYANE JAQUELINE GROSS	Aceito
Declaração de Instituição e Infraestrutura	DeclaraInstituicao.pdf	26/05/2022 21:29:46	DAYANE JAQUELINE GROSS	Aceito
Folha de Rosto	Folhaderosto.pdf	26/05/2022 21:29:21	DAYANE JAQUELINE GROSS	Aceito

Situação do Parecer:

Aprovado

Necessita Apreciação da CONEP:

Não

PIRACICABA, 19 de Junho de 2022

Assinado por:
jacks jorge junior
(Coordenador(a))

ANEXO 2: Comprovante de Aceite no Periódico

[FM] Preliminary editor decision #91331

Professor Janusz Moryś <fm.journals@viamedica.pl>
04:23

16 de agosto de 2022

Responder a: jmorys@pum.edu.pl

Para: PhD Alexandre Rodrigues Freire <alerfreire@gmail.com>

PhD Alexandre Rodrigues Freire:

We kindly inform that we have reached a preliminary decision regarding your submission to Folia Morphologica, entitled "Morphological study of the anatomical variations of anterior belly of digastric muscle in Brazilian cadavers".

The final decision will be made after the submission is verified in the anti-plagiarism system.

Our preliminary decision is: **Acceptance without modification**

Professor Janusz Moryś

Phone +91 466 1543

jmorys@pum.edu.pl

Folia Morphologica

Department of Normal Anatomy, Pomeranian Medical University

Al. Powstańców Wielkopolskich 72, 70–110 Szczecin, Poland

tel. (+48 91) 466 15 43, e-mail: jmorys@pum.edu.pl

ANEXO 3: Comprovante de originalidade e anti-plágio

ESTUDO MORFOLÓGICO E MORFOMÉTRICO DAS VARIAÇÕES ANATÔMICAS DO VENTRE ANTERIOR DO MÚSCULO DIGÁSTRICO

RELATÓRIO DE ORIGINALIDADE

13%	10%	10%	4%
ÍNDICE DE SEMELHANÇA	FONTES DA INTERNET	PUBLICAÇÕES	DOCUMENTOS DOS ALUNOS

FONTES PRIMÁRIAS

1	Submitted to Saint George's University Documento do Aluno	2%
2	José Aderval Aragão. "Variações Anatômicas - Volume 2", Editora Científica Digital, 2022 Publicação	2%
3	rbc.org.br Fonte da Internet	1%

ANEXO 4: Comprovante de autorização para publicação do artigo



Manuscript ID #91331 - authorisation by Editors of <Folia Morphologica>

Lukasz Stolarczyk <lukasz.stolarczyk@viamedica.pl>
Para: Alexandre R Freire <alerfreire@gmail.com>

Hello,

Yes, you can include this article in any repository, so also in the University database.

This is also explained at the Journal website at the panel dealing with Open Access Policy:

https://journals.viamedica.pl/fofia_morphologica/about/editorialPolicies#custom-0

Best regards,

Lukasz Stolarczyk MD, PhD

Scientific Medical Information, Via Medica

Ul. Swietokrzyska 83; 80-180 Gdansk, Poland

Tel.: +48 320 94 94 (reception desk)

Tel: + 48 320 94 97 (direct line)

Skype: lukestolarczyk

[Texto das mensagens anteriores oculto]

The screenshot displays the website of Folia Morphologica. The header includes the journal's logo and a navigation menu with links: VM Homepage, Journals, Folia Morphologica, Ahead Of Print, Current Issue, Archives, News, Subscription, and About. A search bar is located on the right. Below the navigation bar, there are shortcuts for 'Submit an article', 'Author Guidelines (new)', 'Ahead Of Print', and 'Reviewers 2021'. The main content area shows the breadcrumb 'Folia Morphologica > About the Journal > Editorial Policies' followed by three expandable sections: 'Focus and Scope', 'Peer Review Process', and 'Open Access Policy'. The 'Open Access Policy' section is expanded, revealing text about Creative Commons Attribution-Non Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) and the journal's self-archiving policy. At the bottom, there is a 'Regulations' section and a cookie notice stating 'Important: This website uses cookies. More >>'. A QR code and contact information for Via Medica sp. z o.o. are also present.

FOFIA MORPHOLOGICA

VM Homepage > Journals > Folia Morphologica Ahead Of Print Current Issue Archives News Subscription About

Search

Shortcuts Submit an article Author Guidelines (new) Ahead Of Print Reviewers 2021

Folia Morphologica > About the Journal > Editorial Policies

Focus and Scope

Peer Review Process

Open Access Policy

Articles published in open access are available under **Creative Commons Attribution-Non Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0)** allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

Via Medica (Publisher) self-archiving policy establishes that authors can archive only accepted manuscripts including final PDF version and/or 'ahead of print' in any repository. In details, the author may only post their version provided acknowledgment is given to the original source of publication and a link is inserted to the published article on Via Medica (Publisher) website. Apart from the link mentioned above, the acknowledgment should be provided with the full bibliographic record with DOI number.

Regulations

Important: This website uses cookies. More >>

By "Via Medica sp. z o.o." sp.k., Świętokrzyska 73, 80-180 Gdańsk, Poland
tel.: +48 58 320 94 94, faks: +48 58 320 94 60, e-mail: viamedica@viamedica.pl

17 0411 005 001