

02 FEB 1987

BEST APPROXIMANTS FROM CERTAIN SUBSETS
OF BOUNDED FUNCTIONS

João B. Prolla

Departamento de Matemática
Universidade Estadual de Campinas
Campinas, S.P., Brazil

ABSTRACT. Let A be a subalgebra of $C(T, \mathbb{R})$, where T is a compact Hausdorff space. It is well known that the uniform closure of A is proximal in $C(T, \mathbb{R})$ equipped with the sup-norm. In this paper we show that the uniform closure of $A^+ = \{f \in A; f \geq 0\}$, say V , is proximal too. Moreover, for any bounded non-empty subset $B \subset C(T, \mathbb{R})$, the set $\text{cent}(B; V)$ of relative Chebyshev centers of B (with respect to V) is non-empty. The proof relies on a generalization of Bernstein's Theorem on approximation of a positive continuous function f on $[0, 1]$ by its Bernstein polynomials $B_n(f)$.

Universidade Estadual de Campinas
Instituto de Matemática, Estatística e Ciência da Computação
IMECC - UNICAMP
Caixa Postal 6065
13.081 - Campinas, SP
BRASIL

O conteúdo do presente Relatório Técnico é de única responsabilidade do autor.

Janeiro - 1987

I. M. E. C. C.
BIBLIOTECA

Let T be a topological space and let $C_b(T; \mathbb{R})$ be the Banach space of all bounded continuous real-valued functions on T , equipped with the sup-norm, $\|f\| = \sup\{|f(t)|; t \in T\}$. When T is compact, $C_b(T; \mathbb{R}) = C(T; \mathbb{R})$, the space of all continuous real-valued functions on T . In this case it is well known that any closed subalgebra A of $C(T; \mathbb{R})$ is proximal and several proofs have been presented. Smith and Ward [4] extended this result by proving that any closed subalgebra A of $C(T; \mathbb{R})$, for compact T , has the relative Chebyshev center property in $C(T; \mathbb{R})$. Applying this result to the algebra $A = C(T; \mathbb{R})$ one gets that $C(T; \mathbb{R})$, for compact T , admits Chebyshev centers. This result was obtained for $T = [a, b]$ by Kadets and Zamyatin [6], and for any compact T by Garkavi [7]. It was extended by Mach [8]: indeed, it follows from Theorems 3 and 4 of Mach [8] that, for any topological space T , the algebra $C_b(T; \mathbb{R})$ has the relative Chebyshev center property in $\ell_\infty(T; \mathbb{R})$ and the map $B \rightarrow \text{cent}(B; C_b(T; \mathbb{R}))$ is lower semicontinuous. See also Mach [7]. The result that $C_b(T; \mathbb{R})$ admits Chebyshev centers, for any topological space T , was also noticed Franchetti and Cheney [4]. All these results were generalized and extended by Prolla, Chiacchio and Roversi [10], who showed that any closed subalgebra $A \subset C_b(T; \mathbb{R})$, for an arbitrary topological space T , has the relative Chebyshev center property in both $C_b(T; \mathbb{R})$ and $\ell_\infty(T; \mathbb{R})$ and the map $B \rightarrow \text{cent}(B; A)$ is Lipschitz continuous in the Hausdorff metric d_H with Lipschitz constant not greater than 2. This result was proved using among other things the Stone-Weierstrass Theorem. Since we extended recently this theorem to a description of the closure of A^+ , for compact T (see [11] or [12]), it is natural to attempt to extend this result of [10] to the uniform closure of A^+ . Our Theorem 3 below achieves this objective, even for $A \subset \ell_\infty(T; \mathbb{R})$.

Let us explain our notation and terminology. For any Banach space E , the open and closed balls of center a and radius r are denoted, respectively, by $B(a; r)$ and $\bar{B}(a; r)$. If V is any non-empty subset of E and $a \in E$, then

$$\text{dist}(a; V) := \inf\{\|a - v\|; v \in V\}.$$

We denote by $P_V(a)$ the set of all best approximants to a from V , i.e.,

J.M.E.C.C.
BIBLIOTECA

$$P_V(a) := \{v \in V; \|v - a\| = \text{dist}(a; V)\}.$$

If $P_V(a) \neq \emptyset$ for all $a \in E$, we say that V is proximal in E . If $B \subset E$ is any bounded non-empty subset, then

$$\text{rad}(B; V) := \inf \left\{ \sup_{f \in B} \|f - v\|; v \in V \right\}$$

is called the *relative Chebyshev radius of B with respect to V* . When $V = E$ we write simply $\text{rad}(B)$ and call it the *Chebyshev radius of B* . An element $v \in V$ such that

$$\sup_{f \in B} \|f - v\| = \text{rad}(B; V)$$

is called a *relative Chebyshev center of B with respect to V* , and we denote by $\text{cent}(B; V)$ the set of all such elements. When $V = E$, we write simply $\text{cent}(B)$ and its elements are called the *Chebyshev centers of B* .

When $\text{cent}(B; V) \neq \emptyset$ for any bounded $B \subset E$, we say that V has the *relative Chebyshev center property in E* . When $\text{cent}(B) \neq \emptyset$ for all such B , we say that E admits *Chebyshev centers*.

If T is any non-empty set, we denote by $\ell_\infty(T; \mathbb{R})$ the vector space of all bounded real-valued functions defined on T . When we equip $\ell_\infty(T; \mathbb{R})$ with the sup-norm

$$\|f\| = \sup \{|f(t)|; t \in T\}$$

it becomes a Banach space. When T is a topological space, then the vector subspace of all elements of $\ell_\infty(T; \mathbb{R})$, which are continuous on T , is denoted by $C_b(T; \mathbb{R})$. Since it is closed in $\ell_\infty(T; \mathbb{R})$, it is a Banach space too. When T is compact, then all continuous real-valued functions on T are bounded, i.e., $C(T; \mathbb{R}) = C_b(T; \mathbb{R})$ for compact T . For any topological space T , the space $C_b(T; \mathbb{R})$ is isometrically, algebraically and lattice isomorphic to $C(K; \mathbb{R})$ for some compact Hausdorff space K . When T is a completely regular Hausdorff space, then we may take K to be the Stone-Cech compactification of T . The set of all $f \in \ell_\infty(T; \mathbb{R})$ such that $f(t) \geq 0$, for any $t \in T$, is denoted by $\ell_\infty^+(T; \mathbb{R})$. For any subset $A \subset \ell_\infty(T; \mathbb{R})$, $A^+ := A \cap \ell_\infty^+(T; \mathbb{R})$.

The following definition was introduced in [10].

DEFINITION 1. Let V be a closed non-empty subset of a Banach space E , and let \mathcal{B} be a class of bounded non-empty subsets of E . We say that the pair (V, \mathcal{B}) has property (C) in E , if given $B \in \mathcal{B}$, $w \in V$, $r > 0$ and $\epsilon > 0$ such that $V \cap \bigcap_{f \in B} \bar{B}(f; r) \neq \emptyset$ and $\|f - w\| < r + \epsilon$, for all $f \in B$, there exists $v \in V$ such that $\|v - w\| \leq \epsilon$ and $\|f - v\| \leq r$ for all $f \in B$.

Let us say that V has property (C) in E , if the pair $(V, \mathcal{B}(E))$ has property (C) in E , where $\mathcal{B}(E)$ is the class of all bounded non-empty subsets of E . Clearly, if V has property (C) in E , and F is a closed vector subspace such that $V \subset F \subset E$, then V has property (C) in the Banach space F too.

The following result was proved in [10]. (See Proposition 2.2 and Theorem 2.4 of [10].)

THEOREM 1. Let V be a closed non-empty subset of a Banach space E . If V has property (C) in E , and F is any closed vector subspace of E containing V , then

(1) $\text{cent}(B; V) \neq \emptyset$, for every bounded and non-empty subset B of F .

(2) The map $B \rightarrow \text{cent}(B; V)$ is Lipschitz d_H -continuous, with Lipschitz constant not greater than 2, i.e.,

$$d_H(\text{cent}(K; V), \text{cent}(L; V)) \leq 2 d_H(K, L)$$

for any pair K, L of bounded and non-empty subsets of F .

(3) V is proximal in F .

(4) $d_H(P_V(f), P_V(g)) \leq 2 \|f - g\|$ for any pair f, g in F .

(5) The metric projection P_V admits a continuous selection.

REMARK. The Hausdorff metric d_H is defined as follows:

$$d_H(A, B) = \inf \{r > 0; A \subset B + rU, B \subset A + rU\}$$

where $U = \{v \in E; \|v\| \leq 1\}$, for any pair A, B of bounded and non-empty subsets of E .

THEOREM 2. Let K be a closed and non-empty subset of $\mathcal{L}_\infty(T; \mathbb{R})$ such that, for any pair $w, h \in K$ and $\varepsilon > 0$ the function $((w + \varepsilon) \wedge h)v(w - \varepsilon)$ belongs to K . Let $T_0 \subset T$, and let $K_0 = \{f \in K; f(t) = 0 \text{ for all } t \in T_0\}$. If $V = K$ or if $V = K_0$, if K_0 is non-empty, and $E = \mathcal{L}_\infty(T; \mathbb{R})$, then (1)-(5) of Theorem 1 are true.

PROOF. We have to prove that V has property (C) in $\mathcal{L}_\infty(T; \mathbb{R})$, and it suffices to show that $V = K$ has property (C) in $\mathcal{L}_\infty(T; \mathbb{R})$. Indeed, K_0 is closed too and if $w, h \in K_0$, then by hypothesis $g = ((w + \varepsilon) \wedge h)v(w - \varepsilon)$ belongs to K , since w and h belong to K . Now, if we take $t \in T_0$, then $w(t) = h(t) = 0$, and therefore $g(t) = 0$. Hence $g \in K_0$.

We claim that K has property (C) in $\mathcal{L}_\infty(T; \mathbb{R})$. Indeed, let $B \subset \mathcal{L}_\infty(T; \mathbb{R})$ be a bounded non-empty subset, let $w \in K$, $r > 0$ and $\varepsilon > 0$ be given with $K \cap \bigcap_{f \in B} \overline{B}(f; r) \neq \emptyset$ and $\|f - w\| < r + \varepsilon$ for all $f \in B$. Choose $h \in K$ such that $\|f - h\| \leq r$, for all $f \in B$. Let $v = ((w + \varepsilon) \wedge h)v(w - \varepsilon)$. Then $v \in K$ and $\|v - w\| \leq \varepsilon$. We claim that $\|f - v\| \leq r$ for all $f \in B$. Indeed, let $x \in T$ and $f \in B$ be given.

CASE 1. $|h(x) - w(x)| \leq \varepsilon$.

Then $v(x) = h(x)$ and $|f(x) - v(x)| = |f(x) - h(x)| \leq r$.

CASE 2. $h(x) - w(x) > \varepsilon$.

Then $v(x) = w(x) + \varepsilon$ and $-r \leq f(x) - h(x) < f(x) - w(x) - \varepsilon < r + \varepsilon - \varepsilon = r$.

CASE 3. $h(x) - w(x) < -\varepsilon$.

Then $v(x) = w(x) - \varepsilon$ and $-r = -(r + \varepsilon) + \varepsilon < f(x) - w(x) + \varepsilon < f(x) - h(x) \leq \leq r$. \square

REMARK. It is obvious from the proof of Theorem 2, that whenever a set $K \subset \mathcal{L}_\infty(T; \mathbb{R})$ is such that, for any pair $w, h \in K$ and $\varepsilon > 0$ the function $((w + \varepsilon) \wedge h)v(w - \varepsilon)$ belongs to K , then the set $K_0 = \{f \in K; f(t) = 0 \text{ for all } t \in T_0\}$ has the same property, for any subset $T_0 \subset T$. Hence, to each corollary listed below, there is a corresponding result for K_0 .

whenever K_0 is non-empty. Most of the time we will not state explicitly the corresponding corollary.

COROLLARY 1. Let $a, b \in \ell_\infty(T; \mathbb{R})$, with $a \leq b$, let $V = [a, b] := \{h \in \ell_\infty(T; \mathbb{R}) ; a(x) \leq h(x) \leq b(x), \text{ for all } x \in T\}$ and let $E = \ell_\infty(T; \mathbb{R})$. Then (1)-(5) of Theorem 1 are true.

PROOF. It is easy to see that for any pair w, h in $[a, b]$, the function $((w + \varepsilon) \wedge h) \vee (w - \varepsilon)$ belongs to $[a, b]$. \square

REMARK. Franchetti and Cheney proved the proximality of any order interval in any Banach lattice (see [4, Lemma 3.5]). Roversi proved that $[a, b] \subset \ell_\infty(T; \mathbb{R})$ has the relative Chebyshev center property in $\ell_\infty(T; \mathbb{R})$ (see [13, Proposition 2.6]). Notice that Corollary 1 applies to the set $V = \{h \in \ell_\infty(T; \mathbb{R}) ; h(T) \subset [a, b]\}$ when $[a, b] \subset \mathbb{R}$. Indeed, this case corresponds to take a and b in Corollary 1 to be constant functions.

COROLLARY 2. Let (T, \leq) be a preordered set and let V be the subset of all $f \in \ell_\infty(T; \mathbb{R})$ which are non-decreasing (resp. non-increasing) on T , and let $E = \ell_\infty(T; \mathbb{R})$. Then (1)-(5) of Theorem 1 are true.

PROOF. The set V is a closed sublattice of $\ell_\infty(T; \mathbb{R})$ and $w \pm \varepsilon$ belong to V , for each $w \in V$ and $\varepsilon > 0$. \square

REMARK. When T is a topological space and V is as in Corollary 1 or 2 then $V \cap C_b(T, \mathbb{R})$ has the same property in $\ell_\infty(T, \mathbb{R})$ as V , and analogous results can be formulated. Roversi had proved that the closed sublattice V of Corollary 2 has the relative Chebyshev center property in $\ell_\infty(T; \mathbb{R})$. (See Proposition 2.4 of [13].)

COROLLARY 3. Let K be a closed sublattice of $\ell_\infty(T; \mathbb{R})$ such that for any $w \in K$ and $\varepsilon > 0$, the functions $w + \varepsilon$ and $w - \varepsilon$ belong to K . Let $T_0 \subset T$, and $K_0 = \{f \in K ; f(t) = 0 \text{ for all } t \in T_0\}$. If $V = K$ or $V = K_0$ and $E = \ell_\infty(T; \mathbb{R})$, then (1)-(5) of Theorem 1 are true.

PROOF. Clearly, K satisfies the hypothesis of Theorem 2. \square

REMARK. The hypothesis of Corollary 3 are verified if K is a closed sublattice such that $K + K \subset K$ and K contains the constants; in particular, if K is a closed sublattice containing the constants which is also a convex cone. Hence Corollary 3 is a generalization of an Approximation Theorem of Blatter and Seever [2], [3]. Under the latter hypothesis they proved that K_0 is proximal in $\ell_\infty(T; \mathbb{R})$. Their proof uses their theory of interposition of functions. In [3] they establish a formula for $\text{dist}(f; K)$ in terms of the quasi-proximity defined by K on T . The approximation theorem of Blatter and Seever extends an approximation theorem of Nachbin [9, Appendix, §5, Theorem 6] which proves that any closed lattice cone $K \subset C(T; \mathbb{R})$, containing the constants, is proximal in $C(T; \mathbb{R})$, for T a compact Hausdorff space. (When $T_0 = \emptyset$, then Blatter and Seever's result follows from Nachbin's). Nachbin also proved a formula for $\text{dist}(f; K)$.

In [10] it is considered the case in which V is a closed vector subspace of $\ell_\infty(T; \mathbb{R})$. Then Theorem 2 takes the following simplified form.

THEOREM 2'. Let V be a closed vector subspace of $\ell_\infty(T; \mathbb{R})$ such that, for any $h \in V$ and $\epsilon > 0$, the function $(v \wedge h)v(-\epsilon)$ belongs to V . Then V has property (C) in $\ell_\infty(T; \mathbb{R})$.

Using Theorem 2' the following result was proved in [10].

THEOREM 3. Let V be a closed subalgebra of $\ell_\infty(T; \mathbb{R})$, and let $E = \ell_\infty(T; \mathbb{R})$. Then (1)-(5) of Theorem 1 are true.

The proof of Theorem 3 is reduced to the case of a closed subalgebra V of $C(K; \mathbb{R})$, where K is a compact Hausdorff space, and in this case the proof that V satisfies the hypothesis of Theorem 2' uses the Stone-Weierstrass Theorem. (See [10].) Since any closed subalgebra of $C_b(T; \mathbb{R})$, is closed in $\ell_\infty(T; \mathbb{R})$, Theorem 3 implies our next result.

COROLLARY 4. Let T be a topological space. Let V be a closed subalgebra of $C_b(T; \mathbb{R})$ and let $E = \ell_\infty(T; \mathbb{R})$ or $E = C_b(T; \mathbb{R})$. Then (1)-(5) of Theorem 1 are true. If T is locally compact, and V is a closed subalgebra of $C_0(T; \mathbb{R})$ and $E = \ell_\infty(T; \mathbb{R})$, then (1)-(5) of Theorem 1 are true.

Let us now extend Theorem 3 and Corollary 4 to the uniform closure of the set of positive elements of a given subalgebra A . Firstly, we show that our version of the Stone-Weierstrass theorem ([11] or [12]), describing the uniform closure of A^+ , for $A \subset C(K; \mathbb{R})$, K compact, can be used to prove that such a closed convex cone satisfies the hypothesis of Theorem 2.

LEMMA 1. Let A be a subalgebra of $C(K; \mathbb{R})$, where K is a compact Hausdorff space. For any w and h in the uniform closure of A^+ and any $\varepsilon > 0$, the function $((w + \varepsilon) \wedge h) \vee (w - \varepsilon)$ belongs to the uniform closure of A^+ .

PROOF. Let $g = ((w + \varepsilon) \wedge h) \vee (w - \varepsilon)$. By Theorem 3, Prolla [11] there exists a point $x \in K$ such that

$$\text{dist}(g; A^+) = \text{dist}(g_{[x]}; A_{[x]}^+)$$

where $[x]$ is the equivalence class of x mod. A^+ , i.e., $[x] = \{t \in K; a(t) = a(x) \text{ for all } a \in A^+\}$. Since both w and h belong to the uniform closure of A^+ , they are constant on $[x]$. Let w_0 and h_0 be constant value of w and h , respectively, on the set $[x]$. Notice that $w_0 \geq 0$ and $h_0 \geq 0$. Hence $((w_0 + \varepsilon) \wedge h_0) \vee (w_0 - \varepsilon) \geq 0$.

CASE 1. For any $a \in A^+$, $a(x) = 0$.

In this case $w_0 = h_0 = 0$, and $g(t) = 0$ for all $t \in [x]$. Hence $\|g - 0\|_{[x]} = 0$ and, since $0 \in A^+$, $\text{dist}(g_{[x]}; A_{[x]}^+) = 0$.

CASE 2. For some $a \in A^+$, $a(x) > 0$.

Let $f_0 = (((w_0 + \varepsilon) \wedge h_0) \vee (w_0 - \varepsilon)) / a(x)$ and $f = f_0 a$. Then $f \in A^+$ and $f(t) = g(t)$ for all $t \in [x]$. Hence $\|g - f\|_{[x]} = 0$ and $\text{dist}(g_{[x]}; A_{[x]}^+) = 0$.

In both cases, $\text{dist}(g; A^+) = 0$ and therefore g belongs to the uniform closure of A^+ in $C(K; \mathbb{R})$. \square

THEOREM 4. Let A be a subalgebra of $\ell_\infty(T; \mathbb{R})$, let V be the uniform

closure of A^+ in $\ell_\infty(T; \mathbb{R})$ and let $E = \ell_\infty(T; \mathbb{R})$. Then (1)-(5) of Theorem 1 are true.

PROOF. Let K be the Stone-Cech compactification of T equipped with the discrete topology. Then $\ell_\infty(T; \mathbb{R})$ is isometrically, algebraically and lattice isomorphic to $C(K; \mathbb{R})$. The result now follows from Lemma 1 and Theorem 2. \square

COROLLARY 5. Let T be a topological space and let A be a subalgebra of $C_b(T; \mathbb{R})$. If V denotes the uniform closure of A^+ in $C_b(T; \mathbb{R})$ and $E = \ell_\infty(T; \mathbb{R})$, then (1)-(5) of Theorem 1 are true.

PROOF. The algebra A is a subalgebra of $\ell_\infty(T; \mathbb{R})$ and the uniform closure of A^+ in $C_b(T; \mathbb{R})$ is the same as the uniform closure of A^+ in $\ell_\infty(T; \mathbb{R})$, since $C_b(T; \mathbb{R})$ is closed in $\ell_\infty(T; \mathbb{R})$. \square

COROLLARY 6. Let T be a locally compact space and let A be a subalgebra of $C_0(T; \mathbb{R})$. If V denotes the uniform closure of A^+ in $C_0(T; \mathbb{R})$ and $E = \ell_\infty(T; \mathbb{R})$, then (1)-(5) of Theorem 1 are true.

PROOF. The algebra A is a subalgebra of both $C_b(T; \mathbb{R})$ and $\ell_\infty(T; \mathbb{R})$ and the uniform closure of A^+ in $C_0(T; \mathbb{R})$ is the same as the uniform closure of A^+ in $C_b(T; \mathbb{R})$ and in $\ell_\infty(T; \mathbb{R})$, since $C_0(T; \mathbb{R})$ is closed in both $C_b(T; \mathbb{R})$ and $\ell_\infty(T; \mathbb{R})$. \square

COROLLARY 7. Let T be a topological space (resp. a locally compact space), let $V = C_b^+(T; \mathbb{R})$ (resp. $V = C_0^+(T; \mathbb{R})$), and let $E = \ell_\infty(T; \mathbb{R})$. Then (1)-(5) of Theorem 1 are true.

COROLLARY 8. Let $V = \ell_\infty^+$ (resp. C_0^+) and let $E = \ell_\infty$. Then (1)-(5) of Theorem 1 are true.

PROOF. In Corollary 7 take $T = \mathbb{N}$ with the discrete topology. \square

COROLLARY 9. Let $\varphi \geq 0$ be defined on $T \times T$, and let $V = \{f \in \ell_\infty(T; \mathbb{R}); |f(t) - f(u)| \leq \varphi(t, u) \text{ for all } (t, u) \in T \times T\}$. Then (1)-(5) of Theorem 1, are true.

PROOF. It is easily seen that $((w + \varepsilon) \wedge h) \vee (w - \varepsilon)$ belongs to V , whenever w and h belong to V . Indeed, V is a lattice containing $w \pm \varepsilon$ for any $w \in V$ and $\varepsilon > 0$. \square

As an example of application of Corollary 9, assume that (T, d) is a metric space (or even a pseudo-metric space) and let $\varphi(t, u) = Md(t, u)^\alpha$ for some fixed $M > 0$ and $\alpha \in \mathbb{R}$. Then V consists of all $f \in \ell_\infty(T; \mathbb{R})$ such that $|f(t) - f(u)| \leq Md(t, u)^\alpha$ for all $(t, u) \in T \times T$, i.e., all $f \in \text{Lip}_\alpha$ with Lipschitz constant not greater than M .

In order to state our last corollary let us recall the definition of $C_{\mathbb{R}}(T; \mathbb{R})$ when T is a locally compact Hausdorff space. For any $f \in C_b(T; \mathbb{R})$ and $v \in \mathbb{R}$, we say that $\lim_{t \rightarrow \infty} f(t) = v$ if, given $\varepsilon > 0$

there exists a compact subset $K \subset T$ such that $|f(t) - v| < \varepsilon$ for all $t \in T$, $t \notin K$. Following Amir and Deutsch [1], $C_{\mathbb{R}}(T; \mathbb{R})$ denotes the closed subalgebra of $C_b(T; \mathbb{R})$ of all functions that have "limit at infinity". When $T = \mathbb{N}$ with the discrete topology, we write $c = C_{\mathbb{R}}(\mathbb{N}; \mathbb{R})$.

COROLLARY 10. Let T be a locally compact Hausdorff space (resp. $T = \mathbb{N}$ with the discrete topology), let $V = C_{\mathbb{R}}^+(T; \mathbb{R})$ (resp. $V = c^+$), and let $E = \ell_\infty(T; \mathbb{R})$ (resp. $E = \ell_\infty$). Then (1)-(5) of Theorem 1 are true.

REFERENCES

- [1] AMIR, D. and F. DEUTSCH , Approximation by certain subspaces in the Banach space of continuous vector-valued functions, *J. Approximation Theory* 27 (1979), 254-270 .
- [2] BLATTER, J. and G. L. SEEVER , Inclusions, interposition and approximation. In: *Approximation Theory* (G. G. Lorentz, editor), Academic Press, New York, 1973, pp. 257-261.
- [3] BLATTER, J. and G. L. SEEVER , Interposition and lattice cones of functions, *Trans. Amer. Math. Soc.* 222 (1976), 65-96.
- [4] FRANCHETTI, C. and E. W. CHENEY , Simultaneous approximation and restricted Chebyshev centers in function spaces. In: *Approximation Theory and Applications* (Z. Ziegler, editor), Academic Press, New York, 1981, pp. 65-88 .
- [5] GARKAVI, A. L. , The best possible net and the best possible cross-section of a set in normed space, *Amer. Math. Soc. Transl.* 39 (1964), 111-132.
- [6] KADETS, I. M. and V. ZAMYATIN , Chebyshev centers in the space $C[a, b]$, *Teor. Funkciĭ. Funkcional. Anal. i Prilozhen* 7 (1968), 20-26. (Russian).
- [7] MACH, J. , On the existence of best simultaneous approximation , *J. Approximation Theory* 25 (1979), 258-265.
- [8] MACH, J. , Best simultaneous approximation of bounded functions with values in certain Banach spaces, *Math. Ann.* 240 (1979), 157-164.
- [9] NACHBIN, L. , *Topology and Order*, Van Nostrand Math. Studies 4, Van Nostrand, Princeton, 1965.
- [10] PROLLA, J. B. , CHIACCHIO, A. O. and M. S. M. ROVERSI , Relative Chebyshev centers in subalgebras of function spaces, to appear.

- [11] PROLLA, J. B. , Approximation by positive elements of subalgebras of real-valued functions, *Proc. International Workshop on Multivariate Approximation*, Santiago, Chile, December 15-19 , 1986 .
- [12] PROLLA, J. B. , A generalized Bernstein approximation theorem , to appear.
- [13] ROVERSI, M. S. M. , Best approximation of bounded functions by continuous functions, *J. Approximation Theory* 41 (1984), 135-148.
- [14] SMITH, P. W. and J. D. WARD, Restricted centers in subalgebras of $C(X)$, *J. Approximation Theory* 15 (1975), 54-59.