LIMITING PROPERTIES OF THE EMPIRICAL PROBABILITY GENERATING FUNCTION OF STATIONARY RANDOM SEQUENCES AND PROCESSES

by

Mauro S. Marques
Universidade Estadual de Campinas
IMECC, Caixa Postal 1170
13081 Campinas, SP - Brasil

and

Víctor Pérez-Abreu*
Centro de Investigación en Matemáticas, A. C.
A. P. 402, Guanajuato, Gto., 36000 - México

ABSTRACT

Let $X_1, X_2, \ldots$ be a sequence of strictly stationary discrete random variables with probability generating function $\phi(t)$ and let $\hat{\phi}_n(t) = n^{-1}\sum_{i=1}^{n} t^{X_i}$ be the empirical probability generating function of the first n random variables. A strong law of large numbers and the weak convergence of $n^{1/2}(\hat{\phi}_n - \phi)$ to a Gaussian process are discussed. These results are extended to a sequence of stationary multivariate discrete random variables and to integer valued stationary stochastic processes. Applications to the statistical analysis of discrete data are discussed.

AMS 1980 Subject Classification: Primary 60F05, 60F15.
Secondary 62E20, 62G99

Key words and phrases: Empirical probability generating function, discrete distribution, Gaussian process, Banach space valued random variable, Central limit theorem.

---

* Research of this author was partially supported by Office of Naval Research Grant No. N00014-88-J-1069.

Novembro – 1988

1. **INTRODUCTION.** Let $\{P_k\}_{k\geq 0}$ be a discrete distribution with probability generating function

$$(1.1) \qquad \phi(t) = \sum_{k=0}^{\infty} t^k P_k \qquad |t| \leq 1$$

and let $X_1, X_2, \ldots$ be a sequence of nonnegative integer valued random variables with the same distribution $\{P_k\}_{k\geq 0}$. For each $n \geq 1$ let

$$(1.2) \qquad \hat{\phi}_n(t) = n^{-1} \sum_{j=1}^{n} t^{X_j} \qquad |t| \leq 1$$

be the empirical probability generating function (epgf) of the first n observations.

Under the assumption of $\{X_i\}_{i\geq 1}$ independent, the use of some sample generating functions in statistical inference is by now well known. Thus, for example, Press (1972), Heathcote (1972) and Feuerverger and Mureika (1977) have suggested the use of the empirical characteristic function $\hat{c}_n(t) = n^{-1}\sum_{j=1}^{n} exp(itX_j)$ to construct statistical procedures for stable laws, while Epps et al (1982) have proposed the empirical moment generating function $\hat{m}_n(t) = n^{-1}\sum_{j=1}^{n} exp(tX_j)$ to test hypotheses of goodeness of fit in the presence of nuisance parameters. The limiting behavior of $\hat{c}_n$ has been studied, among others, by Feuerverger and Mureika (1977) and Csörgő (1981a, 1981b), while Maĭboroda (1986) has proved a law of large numbers and a central limit theorem for $\hat{m}_n$. Recently, Kocherlakota and Kocherlakota (1986) and O'Reilly et al (1988) have used the empirical probability generating function $\hat{\phi}_n$ to construct goodness of fit tests for discrete random variables, leading to easier procedures than with the use of $\hat{c}_n$ or $\hat{m}_n$.

The purpose of this paper is to study the limiting behavior of the empirical probability generating function $\hat{\phi}_n$ of a stationary sequence of integer valued random variables $X_1, X_2, \ldots$. In Section 2 we prove a law of large numbers (Theorem 1) for $\hat{\phi}_n$ and a central limit theorem (Theorem 2) for the empirical probability generating process $Y_n(t) = n^{1/2}(\hat{\phi}_n(t)-\phi(t))$ in the Banach space of continuous functions on $[-1,1]$. In Section 3 we discuss extensions of Theorems 1 and 2 to a stationary sequence of multivariate discrete random variables and to integer valued stationary stochastic processes. Applications to the statistical analysis of discrete data are discussed in Section 4.

2. THE EPGF OF A STATIONARY SEQUENCE. Let $(\Omega, \mathcal{F}, P)$ be a probability space where there is defined a stationary sequence $\{X_n\}_{n\geq 1}$ of nonnegative integer valued random variables with the same distribution $\{P_k\}_{k\geq 0}$ and probability generating function $\phi(t)$. Let $\hat{\phi}_n(t)$ be the epgf given by (1.2) and

$$Y_n(t) = n^{1/2}(\hat{\phi}_n(t)-\phi(t)) \qquad |t| \leq 1 \tag{2.1}$$

be the epgf process. We shall denote by $C[-1,1]$ the Banach space of real valued continuous functions on the interval $[-1,1]$ with norm $\|f\| = \sup_{-1\leq t\leq 1} |f(t)|$. Clearly ϕ_n, ϕ and Y_n belong to $C[-1,1]$.

The first result of this paper gives the $\|\cdot\|$-consistency of $\hat{\phi}_n$ as an estimator of ϕ.

THEOREM 1. Let $\{X_n\}_{n\geq 1}$ be a strictly stationary ergodic integer valued random sequence. Then

$$\|\hat{\phi}_n - \phi\| \to 0 \quad \text{a.s.} \tag{2.2}$$

Proof. Define for $n \geq 1$

$$P_{k,n} = n^{-1}\sum_{j=1}^{n} 1_{\{X_j=k\}} \qquad k \geq 0 . \tag{2.3}$$

Then the epgf can be expressed as

$$\hat{\phi}_n(t) = \sum_{k=0}^{\infty} t^k P_{k,n} \qquad t \in [-1,1] \tag{2.4}$$

and therefore for $t \in [-1,1]$ and $n \geq 1$

$$|\hat{\phi}_n(t)-\phi(t)| \leq \sum_{k=0}^{\infty} |P_{k,n} - P_k| \leq 2 . \tag{2.5}$$

Since the stationary sequence $\{X_n\}_{n\geq 1}$ is ergodic, we have that for any $k \geq 0$

$$P_{k,n} = n^{-1} \sum_{i=1}^{n} 1_{\{k\}}(X_i) \xrightarrow[n\to\infty]{} P(X_1=k) = P_k \quad \text{a.s.} \tag{2.6}$$

Then by the dominated convergence theorem, from (2.5) and (2.6) we obtain that

$$\sup_{-1\leq t\leq 1} |\hat{\phi}_n(t)-\phi(t)| \xrightarrow[n\to\infty]{} 0 \quad \text{a.s.} \tag{2.7}$$

and the proof is complete. ■

The next step is to study the central limit theorem or weak convergence of the processes $\{Y_n\}_{n\geq 1}$ in the Banach space $C[-1,1]$, under some mixing conditions on the sequence of random variables $\{X_n\}_{n\geq 1}$. Following Bradley (1986) we introduce some notation to be used in the remaining of the work: For two σ-algebras $\mathcal{F}_1$ and $\mathcal{F}_2$ of the basic probability space $(\Omega,\mathcal{F},P)$ define the dependence coefficients

$$\psi(\mathcal{F}_1,\mathcal{F}_2) := \sup_{A\in\mathcal{F}_1, B\in\mathcal{F}_2} |P(A\cap B)-P(A)P(B)|/(P(A)P(B)) \tag{2.8}$$

$$\varphi(\mathcal{F}_1,\mathcal{F}_2) := \sup_{A\in\mathcal{F}_1, B\in\mathcal{F}_2, P(A)>0} |P(B|A)-P(B)| \tag{2.9}$$

and let $\mathcal{F}_j^k = \sigma(X_i : j \leq i \leq k)$. For each $n = 1,2,3,\ldots$ define

(2.10) $$\psi(n) := \sup_{j\geq 1} \psi(\mathcal{F}_1^j, \mathcal{F}_j^\infty)$$

(2.11) $$\varphi(n) := \sup_{j\geq 1} \varphi(\mathcal{F}_1^j, \mathcal{F}_j^\infty)$$

We recall that the sequence of random variables $\{X_n\}_{n\geq 1}$ is said to be φ-mixing if $\lim_{n\to\infty} \varphi(n) = 0$ and ψ-mixing if $\lim_{n\to\infty} \psi(n) = 0$. Also the sequence $\{X_n\}_{n\geq 1}$ is φ-mixing (see Peligrad (1986)) if and only if

(2.12) $$|EXY-EXEY| \leq 2\phi^{1/2}(n)\|X\|_1 \|Y\|_\infty$$

for every $X \in L_1(\mathcal{F}_1^m)$, $Y \in L_\infty(\mathcal{F}_{n+m}^\infty)$ and for every m; and is ψ-mixing (see Peligrad (1986)) if and only if

(2.13) $$|EXY-EXEY| \leq \psi(n)E|X|E|Y|$$

for every $X \in L_1(\mathcal{F}_1^m)$, $Y \in L_1(\mathcal{F}_{n+m}^\infty)$ and for every m.

Our second result of this work provides a central limit theorem for the processes $\{Y_n\}_{n\geq 1}$ under two mixing conditions on the sequence of random variables $\{X_n\}_{n\geq 1}$. As usual we denote weak convergence by $\Rightarrow$.

THEOREM 2. Let $\{X_n\}_{n\geq 1}$ be a strictly stationary random sequence with probability generating function $\phi(t)$ and $EX_1^2 < \infty$.

a) Assume $\{X_n\}_{n\geq 1}$ is a φ-mixing and $\sum_{i=1}^\infty \varphi^{1/2}(i) < \infty$. Then $Y_n \Rightarrow Y$ in $C[-1,1]$ where Y is a Gaussian process with mean zero and covariance

(2.14) $$K(s,t) = \phi(st)-\phi(s)\phi(t) + \sum_{k=1}^\infty r_k(s,t) + \sum_{k=1}^\infty r_k(t,s)$$

where $r_k(s,t) = COV(s^{X_1}, t^{X_{1+k}})$ $k \geq 0$.

b) Assume $\{X_n\}$ is ψ-mixing and $\sum_{i=1}^\infty \psi(i) < \infty$. Then $Y_n \Rightarrow Y$ in $C[-1,1]$ where Y is a Gaussian process, with mean zero and covariance (2.14).

Proof. We first observe that if we define the centered sequence of Banach space valued random variables $Z_i(t) = t^{X_i} - \phi(t)$ $|t| \le 1$ and $\mathcal{F}_k^j = \sigma(Z_i : k \le i \le j)$, then $\mathcal{F}_k^j \subset \mathcal{F}_k^j$ and if $\{X_n\}$ is ψ-mixing (repectively φ-mixing) then $\{Z_n\}$ is ψ-mixing (respectively φ-mixing) with the dependence coefficients for the sequence $\{Z_n\}$ being no greater that the corresponding ones for $\{X_n\}$.

Let $S_n(t) \doteq \sum_{i=1}^{n} Z_i(t)$ and $\mathcal{S}_k(t) = E(Z_1(t)Z_{k+1}(t))$ $k = 0,1,2,\ldots$.

Since the sequence $\{Z_i(t)_{n\ge1}\}$ is stationary

$$\sigma_n^2(t) = VAR(S_n(t)) = n\mathcal{S}_0(t) + 2\sum_{k=1}^{n-1}(n-k)\mathcal{S}_k(t) . \tag{2.15}$$

On the other hand using (2.12) we have that

$$\sum_{k=1}^{\infty}\mathcal{S}_k(t) \le 2\mathcal{S}_0(t)\sum_{k=1}^{\infty}\varphi^{1/2}(k) \qquad |t| \le 1 \tag{2.16}$$

and using (2.13) we obtain

$$\sum_{k=1}^{\infty}\mathcal{S}_k(t) \le \mathcal{S}_0(t)\sum_{k=1}^{\infty}\psi(k) \qquad |t| \le 1 . \tag{2.17}$$

Then

$$n^{-1}\sigma_n^2(t) \to \sigma^2(t) \quad \text{as} \quad n \to \infty \tag{2.18}$$

where

$$\sigma^2(t) = \mathcal{S}_0(t) + 2\sum_{k=1}^{\infty}\mathcal{S}_k(t) \qquad |t| \le 1 . \tag{2.19}$$

The weak convergence of the finite dimensional distributions of $Y_n(t) = n^{-1/2}S_n(t)$ to those of a Gaussian process follows using the Cramer-Wold device and, under assumption (a), Theorem 20.1 in Billingsley (1968), and under assumption (b), the main theorem in Nakhapetiyan (1981). To prove tightness of $\{Y_n(t)\}_{n\ge1}$ in $C[-1,1]$ we shall apply Theorem 12.3 in Billingsley (1968). Using (2.12) we have

$$E(Y_n(t)-Y_n(s))^2 \le 2E(Z_1(t)-Z_1(s))^2\sum_{k=1}^{\infty}\varphi^{1/2}(k) \quad s,t \in [-1,1] \tag{2.20}$$

and using (2.13)

$$E(Y_n(t)-Y_n(s))^2 \le E(Z_1(t)-Z_1(s))^2\sum_{k=1}^{\infty}\psi(k) \quad s,t \in [-1,1] . \tag{2.21}$$

Next it is easily shown that

$$E(Z_1(t)-Z_1(s))^2 \le \theta(t-s)^2 \tag{2.22}$$

where $\theta = \sum_{k=1}^{\infty} k^2 P_k < \infty$. Then by Theorem 12.3 in Billingsley (1968) $\{Y_n\}_{n\ge1}$ is tight in $C[-1,1]$ under assumptions a) or b). Finally by Theorem 8.1 of Billingsley $Y_n \Rightarrow Y$ in $C[-1,1]$. To find the covariance of Y we observe that for $n \ge 1$ and $s,t \in [-1,1]$

$$E(Y_n(t)Y_n(s)) = r_0(s,t) + n^{-1}\sum_{k=1}^{n-1}(n-k)r_k(s,t) + n^{-1}\sum_{k=1}^{n-1}(n-k)r_k(t,s) \tag{2.23}$$

As in proving (2.16) and (2.17) we can show that

$$\sum_{k=1}^{\infty} r_k(s,t) \le 2r_0(s,t)\sum_{k=1}^{\infty}\varphi^{1/2}(k) \qquad s,t \in [-1,1] \tag{2.24}$$

$$\sum_{k=1}^{\infty} r_k(s,t) \le r_0(s,t)\sum_{k=1}^{\infty}\psi(k) \qquad s,t \in [-1,1] \tag{2.25}$$

and therefore $E(Y_n(t)Y_n(s)) \xrightarrow[n\to\infty]{} K(s,t)$ where K is given by (2.14) ∎

For the sake of completness we write the conclusions of Theorems 1 and 2 for a sequence of independent random variables.

THEOREM 3. Let $\{X_n\}_{n\ge1}$ be a sequence of independent integer valued random variables with p.g.f. $\phi(t)$ and $EX_1^2 < \infty$. Then

a) $\|\hat{\phi}_n - \phi\| \xrightarrow[n\to\infty]{} 0$ a.s.

b) $Y_n \Rightarrow Y$ in $C[-1,1]$ where $Y(t)$ is a zero mean Gaussian process with covariance

$$K(s,t) = \phi(st)-\phi(s)\phi(t) \qquad s,t \in [-1,1]\ . \tag{2.26}$$

3. EXTENSIONS TO MULTIVARIATE DISTRIBUTIONS AND STATIONARY STOCHASTIC PROCESSES.

We start this section by presenting extensios of Theorems 1 and 2 to multivariate discrete distributions. Let $\underline{X} = (X_1,\ldots,X_r)$ be an r-dimensional discrete random vector with probability generating function

$$\phi(\underline{t}) = E(t_1^{X_1}\ldots t_r^{X_r}) \quad \underline{t} = (t_1,\ldots,t_r) \tag{3.1}$$

and let $\underline{X}^n = (X_1^n,\ldots,X_r^n)$ $n \geq 1$ be a multivariate stationary independent sequence of random variables with the same p.g.f. ϕ. Define the multivariate empirical probability generating function of the first n observations as

$$\hat{\phi}_n(t) = \frac{1}{n}\sum_{i=1}^{n} t_1^{X_1^i}\ldots t_r^{X_r^i} \quad \underline{t} = (t_1,\ldots,t_r) \quad |t| \leq 1 . \tag{3.2}$$

Let $C([-1,1]^r)$ be the metric space of continuous functions from $[-1,1]^r$ to $\mathbb{R}$ with the norm

$$|||f||| = \sup_{\underline{t}\in[-1,1]^r} |f(\underline{t})| . \tag{3.3}$$

We introduce the following notation: for $\underline{t},\underline{s} \in [-1,1]^r$ $\underline{t}\cdot\underline{s} = (t_1s_1,\ldots,t_rs_r)$.

THEOREM 4. If the sequence $\underline{X}^n$ $n \geq 1$ is ergodic $|||\hat{\phi}_n-\phi||| \to 0$ a.s. as $n \to \infty$.

Proof. Let $P_{k_1\ldots k_r} = P(X_1=k_1,\ldots,X_r=k_r)$ and

$$\hat{P}^n_{k_1\ldots k_r} = \frac{1}{n}\sum_{i=1}^{n} 1\left\{X_1^i=k_1,\ldots,X_r^i=k_r\right\}. \tag{3.4}$$

Then

$$\phi(\underline{t}) = \sum_{k_1\ldots k_r=0}^{\infty} t_1^{k_1}\ldots t_r^{k_r} P_{k_1\ldots k_r} \qquad |\underline{t}| \leq 1 \tag{3.5}$$

and

$$\hat{\phi}_n(\underline{t}) = \sum_{k_1\ldots k_r=0}^{\infty} t_1^{k_1}\ldots t_r^{k_r} P_{k_1\ldots k_r} \qquad |\underline{t}| \leq 1 . \tag{3.6}$$

Since the sequence $\underline{X}^n$ is stationary and ergodic

(3.7) $$\hat{P}^n_{k_1\ldots k_r} \xrightarrow[n\to\infty]{} P_{k_1\ldots k_r} \quad \text{a.s.}$$

Then the proof of the theorem ends quite analogous to the proof Theorem 1.

■

THEOREM 5. Let $Y_n(\underline{t}) = n^{1/2}(\hat{\phi}_n(\underline{t})-\phi(\underline{t}))$ and $EX_i^2 < \infty$ for $i=1,\ldots,r$. Let $\varphi(n)$ and $\psi(n)$ be the dependence coefficients for $\{\underline{X}^n\}_{n\geq 1}$ given by (2.10) and (2.11).

a) If $\{\underline{X}^n\}_{n\geq 1}$ is φ-mixing and $\sum_{n=1}^{\infty}\varphi^{1/2}(n) < \infty$ then $Y_n \Rightarrow Y$ in $C([-1,1]^r)$ where Y is a Gaussian random field with zero mean and covariance.

(3.8) $$K(\underline{s},\underline{t}) = \phi(\underline{s},\underline{t})-\phi(\underline{s})\phi(\underline{t}) + \sum_{k=1}^{\infty} r_k(\underline{s},\underline{t}) + \sum_{k=1}^{\infty} r_k(\underline{t},\underline{s}) \qquad \underline{s},\underline{t} \in [-1,1]^r$$

where $r_k(\underline{s},\underline{t}) = COV(s_1^{X_1^1}\ldots s_r^{X_r^1}, t_1^{X_1^{1+k}}\ldots t_r^{X_r^{1+k}})$

b) If $\{\underline{X}^n\}_{n\geq 1}$ is ψ-mixing and $\sum_{n=1}^{\infty}\psi(n) < \infty$ then $Y_n \Rightarrow Y$ in $C([-1,1]^r)$ where Y is a Gaussian random field with zero mean and covariance (3.8).

The proof of the theorem is quite analogous to the proof of Theorem 2 using the fact that

(3.9) $$E|Y_1(\underline{t})-Y_1(\underline{s})|^2 \leq \theta^2 e^2(\underline{s},\underline{t})$$

where $e(\underline{s},\underline{t}) = \max_{1\leq i\leq r} |t_i-s_i|$ and

(3.10) $$\theta^2 = \sum_{k=0}^{\infty} k^2(P_k(1)+\ldots+P_k(r)) < \infty .$$

Let us now consider the case of a stationary stochastic process. Let $\{X_\Delta\}_{\Delta\geq 0}$ be a measurable strictly stationary nonnegative integer

valued stochastic process and let $\phi(t) = E(t^{X_\Delta})$ $|t| \le 1$ be the probability generating function of X_Δ. For $S \ge 0$ define the empirical probability generating function

$$\hat{\phi}_S(t) = S^{-1} \int_0^S t^{X_\Delta} d\Delta \quad |t| \le 1 \tag{3.11}$$

and let

$$Y_S(t) = S^{1/2}(\hat{\phi}_S(t) - \phi(t)) \tag{3.12}$$

THEOREM 6.

a) If $\{X_\Delta\}_{\Delta \ge 0}$ is an ergodic process

$$\|\hat{\phi}_S - \phi\| \to 0 \quad \text{a.s.} \quad \text{as } S \quad \infty .$$

b) If the process $\{X_\Delta\}_{\Delta \ge 0}$ is φ-mixing and $\int_0^\infty \varphi^{1/2}(\Delta)d\Delta < \infty$ then $Y_n \Rightarrow Y$ in $C[-1,1]$ where Y is a Gaussian process with zero mean and covariance

$$K(t_1,t_2) = \phi(t_1 t_2) - \phi(t_1)\phi(t_2) + \int_0^\infty r_\Delta(t_1,t_2)d\Delta + \int_0^\infty r_\Delta(t_2,t_1)d\Delta \tag{3.13}$$

where $r_\Delta(t_1,t_2) = COV(t_1^{X_0} t_2^{X_\Delta})$.

c) If the process $\{X_\Delta\}_{\Delta \ge 0}$ is ψ-mixing and $\int_0^\infty \psi(\Delta)d\Delta < \infty$ then $Y_n \Rightarrow Y$ in $C[-1,1]$ where the process Y is as in (b).

The proof of the theorem is done using the device discussed in Kolmogorov (1931): First deduce the result for the stationary random sequence

$$\tilde{Y}_n(t) = \int_n^{n+1} Y_\Delta(t)d\Delta$$

and use the measurability condition.

4. APPLICATIONS. We now mention some applications to statistics. Let $X_1, X_2, \ldots$ be a strictly stationary sequence having a discrete distribution F_0 with probability generating function $\phi(t)$ $|t| \le 1$. The statistics

$$d = n \int_0^1 (\hat{\phi}_n(t) - \phi(t))^2 dt$$

can be used to test a goodness of fit for the specified distribution F_0. For example, if F_0 is the Poisson distribution with parameter $\lambda > 0$ then

$$d = n^{-1} \sum_{i=1}^{n} \sum_{j=1}^{n} 1/(X_i + X_j + 1) + (2\lambda)^{-1} n[1 - exp(-2\lambda)]$$
$$- 2 \sum_{i=1}^{n} [(-1)^{X_i+1} X_i!/(\lambda^{X_i+1}) exp(-\lambda)$$
$$+ \sum_{j=0}^{X_i} (-1)^j X_i (X_i - 1) \ldots (X_i - j + 1)/(\lambda^{X_i+1})]$$

In this case, under the assumption of independence and using Theorem 3, the distribution of d has been computed in O'Reilly et al (1988).

REFERENCES

[1] Billingsley, P. (1968). *Convergence of Probability Measures.* Wiley, New York.

[2] Bradley, R. (1986). Basic properties of strong mixing conditions, in E. Eberlein and M. S. Taqqu eds: *Dependence in Probability and Statistics: A Survey of Recent Results.* Birkhäuser, Boston, 166-192.

[3] Csörgo, S. (1981a). Limit behaviour of the empirical characteristic function. *Ann. Probab.* 9, 130-144.

[4] Csörgo, S. (1981b). The empirical characteristic process when parameters are estimated, in J. Gani and V. K. Rohatgi eds: *Contributions to Probability Collection of Papers Dedicated to Eugene Lukacs.* Academic Press, New York, 215-230.

[5] Epps, T. W., K. J. Singleton and L. B. Pulley (1982). A test of separated families of distributions based on the empirical moment generating function. *Biometrika* 69, 391,399.

[6] Feuerverger, A. and R. A. Mureika (1977). The empirical characteristic function and its applications. *Ann Statist.* 5, 88-97.

[7] Heathcote, C. R. (1972). A test of goodness of fit for symmetric random variables. *Austral. J. Statist.* 14, 172-181.

[8] Kocherlakota, S. and K. Kocherlakota (1986). Goodness of fit tests for discrete distributions. *Commun. Statist. Theor. Meth.* 15, 815-829.

[9] Kolmogorov, A. N. (1931). Über die analytischen Methoden in der Wahrscheinlichkeitsrechnung. *Mat. Ann.* 104, 415-458.

[10] Maĭboroda, R. E. (1986). An estimator of the moment generating function of a random variable from the results of observations. *Theor. Probab. and Math. Statist.* 32, 141-151.

[11] Nakhapetiyan, B. S. (1981). On a criterion of weak convergence. *Theory. Probab. Appl.* 21, 370-377.

[12] O'Reilly, F., V. Pérez-Abreu and R. Rueda (1988). The empirical probability generating function with estimated parameters: A Poisson goodness of fit test. *Preprint.*

[13] Peligrad, M. (1986). Recent advances in the central limit theorem and its weak invariance principle for mixing sequences of random variables, in E. Eberlein and M. S. Taqqu eds: *Dependence in Probability and Statistics: A Survey of Recent Results.* Birkhäuser, Boston, 193-223.

[14] Press, S. J. (1972). Estimation in univariate and multivariate stable distributions. *J. Amer. Statist. Assoc.* 67, 842-846.

RELATÓRIOS TÉCNICOS — 1988

01/88 – A Linear Continuous Transportation Problem – *Enrique D. Andjel, Tarcísio L. Lopes* and *José Mario Martínez.*

02/88 – A Splitting Theorem for Complete Manifolds With Non–Negative Curvature Operator – *Maria Helena Noronha.*

03/88 – Mathematical Physics of the Generalized Monopole without String – *W. A. Rodrigues Jr., M. A. Faria–Rosa, A. Maia Jr.* and *E. Recami.*

04/88 – A Family of Quasi–Newton Methods with Direct Secant Updates of Matrix Factorizations – *José Mário Martínez.*

05/88 – Rotation Numbers of Differential Equations. A Framework in the Linear Case – *Luis San Martin.*

06/88 – A Geometrical Theory of non Topological Magnetic Monopoles – *Marcio A. Faria–Rosa* and *Waldyr A. Rodrigues Jr.*

07/88 – Cosmic Walls and Axially Symmetric Sigma Models – *Patricio S. Letelier* and *Enric Verdaguer.*

08/88 – Verificação do Nível de Enlace do Protocolo X–25 – *Célio C. Guimarães* e *Edmundo R. M. Madeira.*

09/88 – A Numerically Stable Reduced–Gradient Type Algorithm for Solving Large–Scale Linearly Constrained Minimization Problemas – *Herminio Simões Gomes* and *José Mário Martínez.*

10/88 - On Integral Bases of Some Ring of Integers – *Nelo D. Allan.*

11/88 – Generating Inexact–Newton Methods Using Least Change Secant Update Procedures – *José Mario Martínez.*

12/88 – Polarized Partition Relations of Higher Dimension – *Walter Alexandre Carnielli* and *Carlos Augusto Di Prisco.*

13/88 – Teoria e Prática no Planejamento de Experimentos – *Armando M. Infante.*

14/88 – On Closed Twisted Curves – *Sueli I. R. Costa.*

15/88 – Green's Function and Isotropic Harmonic Oscillator – *E. Capelas de Oliveira.*

16/88 – A Hopf Bifurcation Theorem for Evolution Equations of Hyperbolic Type – *Aloisio Freiria Neves* and *Hermano de Souza Ribeiro.*

17/88 – Nonnegatively Curved Submanifolds in Codimension Two – *Maria Helena Noronha.*

18/88 – A Comment on the Twin Paradox and the Hafele–Keating Experiment – *W. A. Rodrigues Jr.* and *E. C. Oliveira.*