

R 2792

RT-IMECC
IM/4078

**A NEW CONSTRUCTION OF THE
CASIMIR OPERATORS FOR THE
FANTAPPIÉ-de SITTER GROUP**


E. A. Notte Cuello
and
E. Capelas de Oliveira

Dezembro

RP 62/92

Relatório de Pesquisa

**Instituto de Matemática
Estatística e Ciência da Computação**



**UNIVERSIDADE ESTADUAL DE CAMPINAS
Campinas - São Paulo - Brasil**

R.P.
IM/62/92



2000870883

IMECC

Periódico

ABSTRACT - We obtain the Casimir operators associated to the Fantappiè-de Sitter Group (isomorphic to the 5-dimensional pseudo-rotation group) which is the group of motions admitted by the de Sitter cosmological spacetime, using the generalized derivative operators.

IMECC - UNICAMP
Universidade Estadual de Campinas
CP 6065
13081 Campinas SP
Brasil

O conteúdo do presente Relatório de Pesquisa é de única responsabilidade dos autores.

Dezembro - 1992

I. M. E. C. C.
BIBLIOTECA

A NEW CONSTRUCTION OF THE CASIMIR OPERATORS FOR THE FANTAPPIÉ-de SITTER GROUP

E. A. Notte Cuello and E. Capelas de Oliveira
Departamento de Matemática Aplicada - IMECC
Universidade Estadual de Campinas
13081-970 Campinas São Paulo Brasil

Abstract

We obtain the Casimir operators associated to the Fantappié-de Sitter Group (isomorphic to the 5-dimensional pseudo-rotation group) which is the group of motions admitted by the de Sitter cosmological spacetime, using the generalized derivative operators.

1. INTRODUCTION

The de Sitter space is the curved spacetime which has been most studied by quantum field theorists because together with the anti-de Sitter space are the unique maximally symmetric curved spacetimes⁽¹⁾.

The symmetry group of de Sitter space is the ten parameters group $SO(4, 1)$ of homogeneous Lorentz transformations in the 5-dimensional embedding space known as the de Sitter group⁽²⁾.

There are many possible coordinatizations for the study the de Sitter space time, for example the steady-state universe (parametrization) of Bondi & Gold⁽³⁾ and Hoyle⁽⁴⁾ which covers the half of the de Sitter manifold and the static system which also only covers the half of the de Sitter manifold⁽⁵⁾. Tagirov⁽⁶⁾ studies the Einstein universe using a conformal time where the coordinates cover the whole of the de Sitter manifold.

The theory of hyperspherical models of universe (tied to the integer numbers) proposed by Fantappié⁽⁷⁾ and perfected by Arcidiacono^(8,9) based on group theory is a original way to study the cosmological problem. In this theory it is necessary to distinguish the absolute spacetime (with constant curvature) effective seat of the physics events from the infinite relative space time (tangents) where each observer localize and see the phenomena. Then we use a flat representation of the de Sitter universe on their tangent spaces. Among the infinite representations we use the Beltrami⁽¹⁰⁾ geodesic representation where the geodesics of the hyperspherical spacetime corresponds to the straight lines of the flat tangent space time of the observer's location.

It follows that the group of motions in itself of the de Sitter universe is represented by the so called Fantappiè-de Sitter Group (Isomorphic to the 5-dimensional pseudo-rotation group.) i.e. by the projectivities of the tangent space, which change in itself the Cayley-Klein absolute of equation

$$R^2 A^2 = (x_1)^2 + (x_2)^2 + (x_3)^2 + (x_0)^2 + R^2 = 0$$

where $x_0 = ict$. (For the definition of A see eq (2.2))

This paper, which is the first of three, is organized as follow: in section two we present how to pass from the de Sitter formulation to the orthogonal coordinates using the Beltrami geodesic representation and we obtain the formulas with relates the derivatives; in section three we discuss the Fantappiè-de Sitter group and we obtain the explicit formulas for the invariant associated operators (Casimir operators); in section four, using spherical coordinates we present the comutation relations and construct explicitly the Casimir invariant operators and finally we present ours comments.

In the following papers we solve the equations obtained from the second and fourth orders Casimir invariant operators.

2. ORTHOGONAL COORDINATES AND DERIVATIVES

In this section we consider how to pass from the Beltrami representation $x_\mu (\mu = 0, 1, 2, 3)$ of the de Sitter Universe to the homogeneous coordinates $\xi_A (A = 0, 1, 2, 3, 4)$ of the embedded space $R_{4,1}$. They are related by (see Apendice).

$$x_\mu = R \frac{\xi_\mu}{\xi_4} \quad (2.1)$$

satisfying the relation of normalization $\xi_A \xi_A = R^2$ where R is the radius of the de Sitter universe.

Introducing the following notation

$$A^2 = 1 + \alpha^2 - \gamma^2 = 1 + \alpha_\mu \alpha_\mu \quad (2.2)$$

where

$$\alpha_\mu = \frac{1}{R} x_\mu \quad \text{and} \quad \gamma = \frac{1}{R} ct = \frac{t}{t_0}$$

we can remove the ξ_4 coordinate, then we have the following relations

$$\xi_A = \frac{R}{A} \quad \text{and} \quad \xi_\mu = \frac{x_\mu}{A}.$$

To obtain the relation for the partial derivatives we consider a function $\varphi(\xi_A)$ being an homogeneous function of degree N in all five variables ξ_A , and using Euler's theorem for homogeneous functions, we have

$$\xi_A \partial_A \varphi(\xi_A) = N \varphi(\xi_A) \quad (2.3)$$

where we have put $\partial_A = \partial/\partial \xi_A$. Using the definition of homogeneous function we can write

$$\varphi\left(R \frac{\xi_4}{\xi_4}, R \frac{\xi_1}{\xi_4}, \dots\right) = \left(\frac{R}{\xi_4}\right)^N \varphi(\xi_A) \quad (2.4)$$

and finally we get the following relation

$$R^N \varphi(\xi_A) = (\xi_4)^N \varphi(R, x_\mu) \quad (2.5)$$

where the function in the right hand side is a function obtained from $\varphi(\xi_A)$ with the substitutions, $\xi_4 \rightarrow R$ and $\xi_\mu \rightarrow x_\mu$.

Deriving eq. (2.7) firstly in relation to ξ_4 and secondly in relation ξ_μ we obtain, respectively

$$R \frac{\partial}{\partial \xi_4} \varphi(\xi_A) = A^{1-N} (N - x_\mu \partial_\mu) \varphi(R, x_\mu) \quad (2.8a)$$

and

$$\frac{\partial}{\partial \xi_\mu} \varphi(\xi_A) = A^{1-N} \partial_\mu \varphi(R, x_\mu) \quad (2.8b)$$

where A is given by eq. (2.2) and we have put $\partial_\mu \equiv \partial/\partial x_\mu$.

Introducing a function $\psi(x_\mu)$ defined by

$$\psi(x_\mu) = A^{-N} \varphi(R, x_\mu) \quad (2.9)$$

in the above equations we can finally write the derivatives, respectively, as follows

$$R \frac{\partial}{\partial \xi_4} \varphi(\xi_A) = \left(\frac{N}{A} - A x_\mu \partial_\mu\right) \psi(x_\mu) \quad (2.10a)$$

and

$$\frac{\partial}{\partial \xi_\mu} \varphi(\xi_A) = \left(A \partial_\mu + \frac{N}{A R^2} x_\mu\right) \psi(x_\mu) \quad (2.10b)$$

Then, we have solved the problem to pass of the 5-dimensional formulation, ξ_A , to spacetime formulation, x_μ , i.e. in orthogonal cartesian coordinates. The relations eq. (2.10a) and eq. (2.10b) are the link between the two formulations.

3. THE FANTAPPIÉ-de SITTER GROUP

In this section we present the Fantappié-de Sitter Group and write its invariant operators.

The Fantappié-de Sitter Group - isomorphic to the 5-dimensional pseudo rotation group - is the group of motions admitted by a cosmological space with line element given by

$$-ds^2 = A^2 dx_\mu dx_\mu = A^2 [(dx_1)^2 + (dx_2)^2 + (dx_3)^2 + (dx_0)^2]$$

where $x_0 = ict$, and $R^2 A^2 = R^2 + \rho^2 - x_0^2$ and $\rho^2 = (x_1)^2 + (x_2)^2 + (x_3)^2$.

This space can be embedded in a flat 5-dimensional space time, being the x_μ , the Beltrami projection from the "sfere" with equation

$$\sum_{A=0}^4 \xi_A \xi_A = (\xi_1)^2 + (\xi_2)^2 + (\xi_3)^2 + (\xi_4)^2 - (\xi_0)^2 = R^2 \quad (3.1)$$

The coordinates are related by the following expressions

$$x_\mu = R \frac{\xi_\mu}{\xi_4} \quad \xi_\mu = \frac{1}{A} x_\mu \quad \xi_4 = \frac{1}{A} R$$

where $\mu = 0, 1, 2, 3$ and for the differential operator we have

$$\frac{\partial}{\partial \xi_\mu} = A \partial_\mu + \frac{N}{AR^2} x_\mu \quad (3.3a)$$

and

$$\frac{\partial}{\partial \xi_4} = \frac{1}{R} \left(\frac{N}{A} - A x_\mu \partial_\mu \right) \quad (3.3b)$$

where N is a parameter, R is the radius of the de Sitter universe and A is given by eq. (2.2).

The generators of the 5-dimensional pseudo rotation group satisfy⁽²⁾,

$$\begin{aligned} -i[J_{k\lambda}, J_{\mu\nu}] &= \delta_{k\nu} J_{\lambda\mu} - \delta_{k\mu} J_{\lambda\nu} + \delta_{\lambda\mu} J_{k\nu} - \delta_{\lambda\nu} J_{k\mu} \\ -i[\pi_\lambda, J_{\mu\nu}] &= \delta_{\lambda\mu} \pi_\nu - \delta_{\lambda\nu} \pi_\mu \\ -i[\pi_\mu, \pi_\nu] &= -\frac{1}{R^2} J_{\mu\nu} \end{aligned}$$

where $\pi_\mu = \frac{1}{R} J_{0\mu}$. We note that to $R \rightarrow \infty$ we have

$$\pi_\mu \rightarrow p_\mu$$

which is the four dimensional operator associate with the translations of the Minkowski space time, when $R \rightarrow \infty$ we obtain the Lie Algebra of the non homogeneous Lorentz

group.

Introducing

$$p_\mu \rightarrow -i\partial_\mu$$

we have a representation for the Fantappiè-de Sitter group which is given by the 5-dimensional angular momentum operators

$$J_{AB} = -i\hbar \left(\xi_A \frac{\partial}{\partial \xi_B} - \xi_B \frac{\partial}{\partial \xi_A} \right) \equiv L_{AB}$$

where $A, B = 0, 1, 2, 3, 4$ which in terms of the Beltrami coordinates are given by

$$L_{\mu\nu} = x_\mu p_\nu - x_\nu p_\mu \quad (3.4a)$$

and

$$\pi_\lambda \equiv \frac{1}{R} L_{0\lambda} = A^2 p_\lambda + \frac{1}{R^2} x_\mu L_{\lambda\mu} \quad (3.4b)$$

where $\mu, \nu, \lambda = 0, 1, 2, 3$.

We note that in the above equations (where π_μ are the analogous of the momentum operators in the Minkowski space) that the linear momentum and the angular momentum mix in a unique tensor. This mixing is due to the fact that transformation of displacements are the analogous of the translations and therefore the energy-momentum operators are not conserved in relation to the Fantappiè-de Sitter group.

Now, we consider the explicit form to the ten operators. Introducing the T_0 -operator, representing the temporal translations, defined by

$$L_{40} \equiv \frac{i}{c} R T_0 = -i\hbar \left(\xi_4 \frac{\partial}{\partial \xi_0} - \xi_0 \frac{\partial}{\partial \xi_4} \right)$$

we have

$$T_0 = -\hbar c \left(\partial_0 + \frac{1}{R^2} x_0 x_\mu \partial_\mu \right) \quad (3.5)$$

where $\partial_\mu = \partial/\partial x_\mu$ and $\mu = 0, 1, 2, 3$.

The T_μ -operators, representing the "spatial translations", are defined by

$$L_{\mu 4} \equiv R T_\mu = -i\hbar \left(\xi_\mu \frac{\partial}{\partial \xi_4} - \xi_4 \frac{\partial}{\partial \xi_\mu} \right)$$

and we obtain

$$T_\mu = \frac{i\hbar}{R^2} (R^2 \partial_\mu + x^\nu x_\nu \partial_\mu) \quad (3.6)$$

where $\mu = 1, 2, 3$ and $\nu = 0, 1, 2, 3$.

Introducing the V_μ -operators which are related to the center of mass inertia momentum, given by

$$L_{0\mu} \equiv icV_\mu = -i\hbar \left(\xi_0 \frac{\partial}{\partial \xi_\mu} - \xi_\mu \frac{\partial}{\partial \xi_0} \right)$$

we have

$$V_\mu = -\frac{\hbar}{c} (x_0 \partial_\mu - x_\mu \partial_0) \quad (3.7)$$

where $\mu = 1, 2, 3$.

Finally, we introduce L_λ -operators, representing the spatial rotations, defined by

$$L_{\mu\nu} \equiv L_\lambda = -i\hbar \left(\xi_\mu \frac{\partial}{\partial \xi_\nu} - \xi_\nu \frac{\partial}{\partial \xi_\mu} \right)$$

and we obtain

$$L_\lambda = -i\hbar (x_\mu \partial_\nu - x_\nu \partial_\mu) \quad (3.8)$$

where $\mu, \nu, \lambda = 1, 2, 3$ and in the above expressions \hbar and c have the usual meanings.

Now, we can write the two invariant operators of the Fantappiè-de Sitter group (Casimir operators) using T_0 , T_μ , V_μ and L_μ as follow

$$I_2 = (T^2 - \frac{1}{c^2} T_0^2) + \frac{1}{R^2} (L^2 - c^2 V^2) = M^2 \quad (3.9a)$$

and

$$I_4 = (\vec{L} \cdot \vec{T})^2 - \frac{1}{c^2} (T_0 \vec{L} + c^2 \vec{T} \times \vec{V})^2 - \frac{c^2}{R^2} (\vec{L} \cdot \vec{V})^2 = N^2 \quad (3.9b)$$

where M^2 and N^2 are constants.

We note that, in the limit $R \rightarrow \infty$ we obtain

$$I_2 \rightarrow m^2 \quad \text{and} \quad I_4 \rightarrow m^2 s(s+1)$$

where m and s are, respectively, the rest mass and the spin which characterize the representations of the Poincaré Group^[2]. Then, the representations of the Fantappiè-de Sitter group are labeled by eigenvalues of I_2 and I_4 which generalizes the usual mass and spin. Yet, a particle in a Fantappiè-de Sitter universe has not a well defined mass and a spin but eigenvalues of the I_2 and I_4 invariant operators.

4. COMUTATION RELATIONS AND CASIMIR OPERATORS

In this section we introduce a spherical coordinate sistem (r, θ, ϕ) and we obtain the explicit comutation relations and the explicit form of Casimir operators, in these coordinates. This result is important for the following papers^(14,15).

The relativistic spherical coordinates are given by $x_0 = t$, $x_3 = r \cos \theta$, $x_2 = r \sin \theta \sin \phi$ and $x_1 = r \sin \theta \cos \phi$, and we obtain ten differential operators in the explicit forms given by

$$T_0 = -\hbar c \left[\left(1 + \frac{t^2}{R^2}\right) \frac{\partial}{\partial t} + \frac{tr}{R^2} \frac{\partial}{\partial r} \right]$$

$$T_1 = -\frac{i\hbar}{R^2} \left[(r^2 + R^2) \sin \theta \cos \phi \frac{\partial}{\partial r} + \frac{R^2}{r} \left(\cos \theta \cos \phi \frac{\partial}{\partial \theta} - \frac{\sin \phi}{\sin \theta} \frac{\partial}{\partial \phi} \right) + rt \sin \theta \cos \phi \frac{\partial}{\partial t} \right]$$

$$T_2 = -\frac{i\hbar}{R^2} \left[(r^2 + R^2) \sin \theta \sin \phi \frac{\partial}{\partial r} + \frac{R^2}{r} \left(\cos \theta \sin \phi \frac{\partial}{\partial \theta} + \frac{\cos \phi}{\sin \theta} \frac{\partial}{\partial \phi} \right) + rt \sin \theta \sin \phi \frac{\partial}{\partial t} \right]$$

$$T_3 = -\frac{i\hbar}{R^2} \left[(r^2 + R^2) \cos \theta \frac{\partial}{\partial r} - \frac{R^2}{r} \sin \theta \frac{\partial}{\partial \theta} + rt \cos \theta \frac{\partial}{\partial t} \right]$$

$$V_1 = \frac{\hbar}{c} \left[t \left(\sin \theta \cos \phi \frac{\partial}{\partial r} + \frac{1}{r} \cos \phi \cos \theta \frac{\partial}{\partial \theta} - \frac{1}{r} \frac{\sin \phi}{\sin \theta} \frac{\partial}{\partial \phi} \right) - r \sin \theta \cos \phi \frac{\partial}{\partial t} \right]$$

$$V_2 = \frac{\hbar}{c} \left[t \left(\sin \theta \sin \phi \frac{\partial}{\partial r} + \frac{1}{r} \cos \theta \sin \phi \frac{\partial}{\partial \theta} + \frac{1}{r} \frac{\cos \phi}{\sin \theta} \frac{\partial}{\partial \phi} \right) - r \sin \theta \sin \phi \frac{\partial}{\partial t} \right]$$

$$V_3 = \frac{\hbar}{c} \left[t \left(\cos \theta \frac{\partial}{\partial r} - \frac{1}{r} \sin \theta \frac{\partial}{\partial \theta} \right) - r \cos \theta \frac{\partial}{\partial t} \right]$$

$$L_1 = i\hbar \left(-\sin \phi \frac{\partial}{\partial \theta} - \cot \theta \cos \phi \frac{\partial}{\partial \phi} \right)$$

$$L_2 = i\hbar \left(-\cos \phi \frac{\partial}{\partial \theta} + \cot \theta \sin \phi \frac{\partial}{\partial \phi} \right)$$

$$L_3 = i\hbar \frac{\partial}{\partial \phi}$$

Now, considering a cyclic permutation of the index μ , ν and λ we obtain the following comutation relations for the differential operators,

$$[T_0, T_\mu] = -i\hbar \frac{c^2}{R^2} V_\mu \quad [T_0, V_\mu] = -i\hbar T_\mu \quad [T_0, R_\mu] = 0$$

$$[T_\mu, T_\nu] = -\frac{i\hbar}{R^2} L_\lambda \quad [T_\mu, V_\nu] = -\frac{i\hbar}{c^2} \delta_{\mu\nu} T_0 \quad [T_\mu, L_\nu] = i\hbar T_\lambda$$

$$[V_\mu, V_\nu] = \frac{i\hbar}{c^2} L_\lambda \quad [L_\mu, V_\nu] = -i\hbar V_\lambda \quad [L_\mu, L_\nu] = -i\hbar L_\lambda$$

where $\mu, \nu, \lambda = 1, 2, 3$.

Finally, we obtain the explicit forms for the Casimir operators, introducing the differential operators given above in eq. (3.9a) and (3.9b).

The Casimir operator of second order is given by

$$I_2 = -\hbar^2 A^2 \left\{ \left(1 + \frac{r^2}{R^2}\right) \frac{\partial^2}{\partial r^2} + 2rt \frac{\partial^2}{\partial r \partial t} - \left(1 - \frac{t^2 c^2}{R^2}\right) \frac{\partial^2}{\partial (ct)^2} + \right. \\ \left. + \frac{2}{r} \left(1 + \frac{r^2}{R^2}\right) \frac{\partial}{\partial r} + \frac{2t}{R^2} \frac{\partial}{\partial t} + \frac{1}{r^2} \mathcal{L}^2 \right\}$$

where we have put $t \rightarrow ict$ and the \mathcal{L}^2 operator is

$$\mathcal{L}^2 = \frac{\partial^2}{\partial \theta^2} + \cot \theta \frac{\partial}{\partial \theta} + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \phi^2}.$$

We note that when $R \rightarrow \infty$ the above equation reduces to, the D'Alembert wave operator, i.e.,

$$\lim_{R \rightarrow \infty} I_2 \equiv \square = \hbar^2 \left(\Delta - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right)$$

where Δ is the Laplacian operators written in spherical coordinates.

For the fourth order Casimir differential operator we have

$$I_4 = 4 \frac{c^2 \hbar^2}{R^4} \mathcal{L}^2 \mathcal{H}^2$$

where

$$\mathcal{H}^2 = t^2 r^2 \frac{\partial^2}{\partial r^2} + (R^2 - t^2) \frac{\partial^2}{\partial t^2} - 2rt(R^2 - t^2) \frac{\partial^2}{\partial r \partial t} + r(2t^2 - R^2) \frac{\partial}{\partial r} - 2t(R^2 - t^2) \frac{\partial}{\partial t}$$

and \mathcal{L}^2 is given above.

5. COMMENTS

In this paper we have discussed an alternative way to obtain the Casimir invariant operators of the Fantappi -de Sitter Group which is isomorphic to 5-dimensional pseudo rotation group.

It is clear the dependence in the two Casimir invariant operators in both spatial and temporal parts, given by the T_0 , T_μ , V_μ and L_μ operators. In consequence, a particle in a Fantappi -de Sitter universe has not a well defined mass and a spin but has constant eigenvalues of the I_2 and I_4 Casimir invariant operators.

The next point is solve the generalized Klein-Gordon wave equation for the scalar field^(11, 12, 13) obtained from the second order Casimir invariant operator and the

equation obtained from the fourth order Casimir invariant operator, which must generalize the concept of mass and spin^(14,15). These topics are presented in another paper.

Acknowledgements

One of us (ECO) is grateful to Professor G. Arcidiacono for giving inspiration for the study of the present theme and to Professor W. A. Rodrigues Jr for many discussions about it and for carefully reading the manuscript. I am also grateful to CNPq for a research grant.


APENDICE

The de Sitter space can be represented as the surface of a four-dimensional pseudosphere (of a hyperbolic character in one direction) embedded in a five-dimensional space. It is described by five coordinates $\xi_1, \xi_2, \xi_3, \xi_4, \xi_0$ connected by the relation of normalization condition

$$\xi_1^2 + \xi_2^2 + \xi_3^2 + \xi_4^2 - \xi_0^2 = R^2$$

R being the radius of the "sphere".

To see how to pass from the five dimensional formulation to the four-dimensional orthogonal coordinates $x_\mu (\mu = 0, 1, 2, 3)$ we consider the Beltrami representation (geodesic representations) as in the figure:



where

$$x_\mu = R \frac{\xi_\mu}{\xi_4}$$

Introducing $\rho^2 = x^\mu x_\mu = -(x_0)^2 + (x_1)^2 + (x_2)^2 + (x_3)^2$ and using normalization condition we can write

$$\xi_4 = \frac{R}{(1 + \rho^2/R^2)^{1/2}}.$$

Then, the relations to pass from the pentadimensional formulation to the four-dimensional formulation are given by

$$\xi_4 = \frac{R}{A} \quad \text{and} \quad \xi_\mu = \frac{x_\mu}{A}$$

where $A^2 = 1 + \rho^2/R^2$

REFERENCES

- (1) S. Weinberg (1972) *Gravitation and Cosmology: Principles and Applications of the General theory of Relativity* (New York: Wiley)
- (2) F. Gürsey (1963) *Introduction to Group Theory in Relativity, Groups and Cosmology* edited by Dewitt and Dewitt (Gordon Breach)
- (3) H. Bondi and T. Gold, *Mon. Not. R. Astron. Soc.* **108**, 252 (1948)
- (4) F. Hoyle, *Mon. Not. R. Astron. Soc.* **108**, 372 (1948)
- (5) G. W. Gibbons and S. W. Hawking, *Phys. Rev. D* **15**, 2738 (1977)
- (6) E. A. Tagirov, *Ann. Phys* **76**, 561 (1973)
- (7) L. Fantappiè, *Opere Scelte*, (1973) Ed. Unione Matematica Italiana (Bologna, Italia)
- (8) G. Arcidiacono, *Gen. Rel. Grav.* **7**, 885(1976), **8**, 865(1977) and **9**, 949(1978)
- (9) G. Arcidiacono (1986) *Projective, Relativity, Cosmology and Gravitation*, Hadronic Press Inc., Massachusetts, USA and (1987) *Relatività e Cosmologia*, Veschi, Roma
- (10) E. Beltrami, *Ann. di Matem.* **8**, 135 (1865)
- (11) G. Arcidiacono and E. Capelas de Oliveira, *Hadronic Journal* **14**, 347 (1991)
- (12) G. Arcidiacono and E. Capelas de Oliveira, *Hadronic Journal* **14**, 353 (1991)
- (13) E. Capelas de Oliveira, "On the solution of homogeneous generalized wave equation" to appear in *J. Math. Phys* (1992)
- (14) E. A. Notte Cuello and E. Capelas de Oliveira "D'Alembert equation in the de-Sitter Universe" to be published (1993)
- (15) E. A. Notte Cuello and E. Capelas de Oliveira "Dirac equation in the de Sitter Universe" to be published (1993).

RELATÓRIOS DE PESQUISA — 1992

- 01/92 Uniform Approximation the: Non-locally Convex Case — *João B. Prolla.*
- 02/92 Compactificação de $L^r_{\text{loc}}(Q)$ com τ Finito — *A. M. Sette and J. C. Cifuentes.*
- 03/92 Um Modelo para Aquisição da Especificação — *Cecilia Inés Sosa Arias and Ariadne Carvalho.*
- 04/92 Convergence Estimates for the Wavelet Galerkin Method — *Sônia M. Gomes and Elsa Cortina.*
- 05/92 Optimal Chemotherapy: A Case Study with Drug Resistance, Saturation Effect and Toxicity — *M. I. S. Costa, J. L. Boldrini and R. C. Bassanezi.*
- 06/92 On the Paper "Cauchy Completeness of Elementary Logic" of D. Mundici and A. M. Sette — *J. C. Cifuentes.*
- 07/92 What is the EM Algorithm for Maximum Likelihood Estimation in PET and How to Accelerate it — *Alvaro R. De Pierro.*
- 08/92 Bifurcation from infinity and multiple solutions for an elliptic system — *Raffaele Chiappinelli and Djairo G. de Figueiredo.*
- 09/92 Approximation Processes for Vector-Valued Continuous Functions — *João B. Prolla.*
- 10/92 Aplicação do Método de Fraïssé à Compactificação de Lógicas com Quantificadores Co-filtro — *A. M. Sette and J. C. Cifuentes.*
- 11/92 Absolutely Summing Holomorphic Mappings — *Mário C. Matos.*
- 12/92 The Feynman-Dyson Proof of Maxwell Equations and Magnetic Monopoles — *Adolfo A. Jr. and Waldyr A. R. Jr.*
- 13/92 A Generalized Dirac's Quantization Condition for Phenomenological Non-abelian Magnetic Monopoles — *Adolfo M. Jr. and Waldyr A. R. Jr.*
- 14/92 Multiplicity Results for the 1-Dimensional Generalized p -Laplacian — *Pedro Ubilla.*
- 15/92 Nowhere Vanishing Torsion Closed Curves Always Hide Twice — *Sueli R. Costa. and Maria Del Carmen R. Fuster.*
- 16/92 Uniform Approximation of Continuous Convex-Cone-Valued Functions — *João B. Prolla.*
- 17/92 Monotonically Dominated Operators on Convex Cones — *A. O. Chiacchio, J. B. Prolla, M. L. B. Queiroz and M. S. M. Roversi.*
- 18/92 Testing the Concept of a Photon as an Extended Object in a Variation of Francon's Experiment — *V. Buonamano, A. J. R. Madureira and L. C. B. Ryff.*
- 19/92 A New Trust Region Algorithm for Boun Constrained Minimization — *Ana Friedlander, José Mario Martínez and Sandra A. Santos.*

- 20/92 A Priori Estimates for Positive Solutions of Semilinear Elliptic Systems Via Hardy-Sobolev Inequalities — Ph. Clément, D.G. de Figueiredo and E. Mitidieri.
- 21/92 On the Resolution of Linearly Constrained Convex Minimization Problems — Ana Friedlander, José Mario Martínez and Sandra A. Santos.
- 22/92 Convergência no Espectro Real de um Anel — J. C. Cifuentes.
- 23/92 Parallel Methods for Large Least Norm Problems — Alvaro R. De Pierro, and Jose M. Lopes.
- 24/92 A Generalization of Dirac Non-Linear Electrodynamics, and Spinning Charged Particles — Waldyr A. Rodrigues Jr., Jayme Vaz Jr. and Erasmo Recami.
- 25/92 Bifurcation of Certain Singularities of Sliding Vector Fields — Marco Antonio Teixeira.
- 26/92 Density of Infimum-Stable Convex Cone — João B. Prolla.
- 27/92 An Extended Decomposition Through Formalization in Product Spaces — Alvaro R. De Pierro.
- 28/92 Accelerating Iterative Algorithms for Symmetric Linear Complementarity Problems — Alvaro R. De Pierro and José Marcos Lopes.
- 29/92 Free Maxwell Equations, Dirac Equation, and Non-Dispersive de Broglie Wavepackets — Waldyr A. Rodrigues Jr., Jayme Vaz Jr. and Erasmo Recami.
- 30/92 New Results on the Equivalence of Regularization and Truncated Iteration for Ill-posed Problems — Reginaldo J. Santos.
- 31/92 Topological Defects: A Distribution Theory Approach — Patricio S. Letelier and Anzhong Wang.
- 32/92 Some Applications of the Generalized Khintchine's Inequality — Mário C. Matos.
- 33/92 A Technique for Finding The Center of A Polytope — Antonio Carlos Moretti.
- 34/92 On The Convergence of the Column-Updating Method — José Mario Martínez.
- 35/92 Post-Linear Formalism For Gravitating Strings: Crossed Straight Strings Collision — D. V. Gal'tsov, Yu. V. Gruts and P. S. Letelier.
- 36/92 A Simple Model of Life Expectancy With Subjective Parameters — R. C. Bassanezi and L. C. Barros.
- 37/92 On the Growth of Caiman Crocodilus Yacare: A Strategy for Optimal Farming — Alejandro B. Engel and R. C. Bassanezi.
- 38/92 The Dirac Operator and the Structure of Riemann-Cartan-Weyl Spaces — Quintino A. G. de Souza and Waldyr A. Rodrigues, Jr.
- 39/92 On the Nature of the Gravitational Field — Waldyr A. Rodrigues, Jr. and Quintino A. G. de Souza.
- 40/92 Spinning Strings and Cosmic Dislocations — D. V. Gal'tsov and P. S. Letelier.
- 41/92 Natural Ultrabornological, Non-Complete, Normed Function Spaces — Antonio Gilioli (edited by Klaus Floret and Chaim S. Höning).
- 42/92 Parallel and Semi-Parallel Immersions Into Space Forms — Francesco Mercuri.

- 43/92 Periodic Motions of Sucker Rod Pumping Systems — *Aloisio Freiria Neves and José Luiz Boldrini.*
- 44/92 Algorithms for Solving Nonlinear Systems of Equations — *José Mario Martínez.*
- 45/92 Spin and Electron Structure — *Matěj Pavšič, Erasmo Recami, Waldyr A. Rodrigues Jr. and G. Daniele Maccarrone, Fabio Raciti, Giovanni Salesi.*
- 46/92 The Initial Value Problem For the Boussinesq Equations in a Time-Dependent Domain — *Carlos Conca Rozende and Marko Rojas Medar.*
- 47/92 A New Expansion for Jacobi Function — *E. Capelas de Oliveira.*
- 48/92 On the Equivalence of Convergences of Fuzzy Sets — *Marko Rojas Medar and Heriberto Román Flores.*
- 49/92 Monotonicity and Symmetry of Solutions of Elliptic Systems in General Domains — *Djairo G. de Figueiredo.*
- 50/92 Tendências em Programação Não Linear de Grande Porte — *José Mario Martínez.*
- 51/92 Teoría y Práctica en Optimización Numérica — *José Mario Martínez.*
- 52/92 The Clifford Bundle and the Nature of the Gravitational Field — *Waldyr A. Rodrigues Jr. and Quintino A. G. de Souza.*
- 53/92 Completamento e Compactificação de Espaços Zero-Dimensionais — *J. C. Cifuentes.*
- 54/92 Some Results on Polarized Partition Relations of Higher Dimension — *Walter Alexandre Carnielli and Carlos Augusto Di Prisco.*
- 55/92 The Generalized Wave Equation: Revisited — *G. C. Ducati and E. Capelas de Oliveira.*
- 56/92 The Four-Vertex Theorem for Constant Curvature Manifolds — *Sueli I. R. Costa and Marcelo Firer.*
- 57/92 The Riemann-Cartan-Weyl Geometry Generated by a Dirac-Hestenes Spinor Field — *Diego Lucio Rapoport-Campodónico, Waldyr A. Rodrigues, Jr., Quintino A. G. de Souza and Jayme Vaz Jr.*
- 58/92 On the Equivalence of Dirac and Maxwell Equations, and Quantum Mechanics — *Jayme Vaz Jr. and Waldyr A. Rodrigues.*
- 59/92 Convergência Fraca de Medidas em Espaços Métricos Separáveis e Completos — *Mario Antonio Gneri.*
- 60/92 Homoclinic Orbits for a Quasilinear Hamiltonian System — *Pedro Ubilla.*
- 61/92 Quasi-Newton Methods and the Solution of Some Fluid Mechanics Problems — *Márcia A. Gomes - Ruggiero, Daniel N. Kozakevich and José Mario Martínez.*