

UNIVERSIDADE ESTADUAL DE CAMPINAS
SISTEMA DE BIBLIOTECAS DA UNICAMP
REPOSITÓRIO DA PRODUÇÃO CIENTÍFICA E INTELLECTUAL DA UNICAMP

Versão do arquivo anexado / Version of attached file:

Versão do Editor / Published Version

Mais informações no site da editora / Further information on publisher's website:

<https://academic.oup.com/botlinnean/article/194/4/439/5890103>

DOI: 10.1093/botlinnean/boaa059

Direitos autorais / Publisher's copyright statement:

©2020 by Wiley. All rights reserved.

DIRETORIA DE TRATAMENTO DA INFORMAÇÃO

Cidade Universitária Zeferino Vaz Barão Geraldo

CEP 13083-970 – Campinas SP

Fone: (19) 3521-6493

<http://www.repositorio.unicamp.br>

Molecular phylogenetic insights into the evolution of *Eriosema* (Fabaceae): a recent tropical savanna-adapted genus

ELISA SILVA CÂNDIDO^{1,*}, MOHAMMAD VATANPARAST^{2,3,*},
WANDERLEIA DE VARGAS⁴, LUÍSA M. P. A. BEZERRA⁴, GWILYM P. LEWIS⁵,
VIDAL F. MANSANO^{1,6}, ANDRÉ O. SIMÕES¹, MARCOS J. SILVA⁷, CHARLES STIRTON⁸,
ANA MARIA G. A. TOZZI¹ and ANA PAULA FORTUNA-PEREZ^{1,4,*}

¹Departamento de Biologia Vegetal, Programa de Pós-Graduação em Biologia Vegetal, Instituto de Biologia, Universidade Estadual de Campinas, 13083-862, Campinas, São Paulo, Brasil

²Department of Geosciences and Natural Resource Management, Rolighedsvej 23, 1958 Frederiksberg C., University of Copenhagen, Denmark

³US National Herbarium (US), Department of Botany, Smithsonian Institution-NMNH, MRC 166, 10th and Constitution Avenue, Washington, D.C., 20560, USA

⁴Departamento de Botânica, Instituto de Biociências, Programa de Pós-Graduação em Ciências Biológicas (Botânica), Instituto de Biociências, Universidade Estadual Paulista, 18618-000, Botucatu, São Paulo, Brasil.

⁵Comparative Plant and Fungal Biology Department, Royal Botanic Gardens, Kew, Richmond, Surrey, TW9 3AB, UK

⁶Instituto de Pesquisas do Jardim Botânico do Rio de Janeiro, DIPEQ, Rua Pacheco Leão, 915, Jardim Botânico, Rio de Janeiro, 22460-030, RJ, Brasil.

⁷Departamento de Botânica, Instituto de Ciências Biológicas, Universidade Federal de Goiás, CP 131, 74001-970, Goiânia, GO, Brasil.

⁸Bolus Herbarium, Department of Biological Sciences, University of Cape Town, 105 Rondebosch, South Africa.

Received 4 February 2020; revised 30 April 2020; accepted for publication 17 June 2020

Eriosema comprises c. 150 species and has a pantropical distribution and two centres of diversity, Africa and America. The species occur in tropical savannas and grasslands, including the cerrado in Brazil. They have adapted to these environments by developing specialized underground organs, and an abundance of trichomes. Here we present the first comprehensive molecular phylogenetic analysis of *Eriosema*, including species from its entire distribution range and generating 391 new DNA sequences. We sampled 140 species from nine genera of Cajaninae, of which 94 (60% of the genus) were *Eriosema*. Our analyses were based on the nuclear ITS and plastid *rpl32* and *trnQ* regions, and used maximum likelihood and Bayesian phylogenetic analyses of individual and combined data sets. In all analyses, *Eriosema* was resolved as monophyletic, but its interspecific relationships are not well resolved. *Rhynchosia* is not monophyletic, and some African *Rhynchosia* spp. emerged together as sister to *Eriosema*. Our study supports the monophyly of *Adenodolichos*, *Dunbaria*, *Flemingia* and *Cajanus*, but *Chrysoscias* and *Bolusafr* formed a clade that is sister to a group of *Rhynchosia* spp. *Paracalyx* was resolved as paraphyletic and nested among African *Rhynchosia* spp. Divergence time analysis suggested that the *Eriosema* lineages diverged 6.5–10.7 Mya. Two major lineages have diversified in *Eriosema*, one including most of the African species (4.41–6.68 Mya), the other mainly composed of

*Corresponding authors. E-mail: ana.fortuna@unesp.br;
elisasilvacandido@gmail.com; mov@ign.ku.dk

[†]These authors contributed equally to this work.

the South American cerrado species (3.56–5.78 Mya). These results revealed that *Eriosema* is a recent and tropical savanna-adapted group, and its diversification occurred in the late Miocene in parallel with the expansion of C₄ grasslands.

ADDITIONAL KEYWORDS: Cajaninae – cerrado – divergence time – grassland – Leguminosae – recent diversification – *Rhynchosia* – systematics.

INTRODUCTION

Eriosema (DC.) Desv. comprises c. 150 species and is the second-largest genus of subtribe Cajaninae (tribe Phaseoleae), Fabaceae (Greear, 1970; LPWG, 2017). It has two centres of diversification, one in Africa (including Madagascar) with c. 110 species and the other in the New World (from Mexico to northern Argentina, except for Chile) with c. 40 species. In addition, one species, *E. chinense* Vogel, occurs in Southeast Asia and northern Australia (Greear, 1970; van der Maesen, 2003; Schrire, 2005). The genus is used throughout its distribution range for traditional medicine and food. For instance, in South Africa, 'Zulu' people use the roots as a medicine for the treatment of erectile dysfunction and/or impotence (Ojewole, 2007). In the Neotropics, Brazilian traditional communities use some species as laxatives or as an anti-inflammatory medicine (Hirschmann & De Arias 1990; Rodrigues & Carvalho, 2001; Santos et al., 2016) and in the Northern Territory of Australia, root tubers of *E. chinense* have been used as human food (Smith, 1991).

Previous molecular phylogenetic studies, although including a limited number of sampled taxa, have demonstrated that subtribe Cajaninae and *Eriosema* are monophyletic (Bruneau, Doyle & Doyle, 1995; Kajita et al., 2001; Egan, Vatanparast & Cagle, 2016; LPWG, 2017). Cajaninae, as traditionally recognized, comprise ten genera: *Adenodolichos* Harms (20 species), *Bolusafr* Kuntze Harms (one species); *Cajanus* DC. (34 species), *Carrissoa* Baker f. (one species), *Chrysoscias* E.Mey. (three or four species), *Dunbaria* Wight & Arn. (20 species), *Eriosema* (150 species), *Flemingia* Roxb. ex W.T.Aiton (30–35 species), *Paracalyx* S.I.Ali (six species) and *Rhynchosia* Lour. (230 species) (Greear, 1970, 1978; Schrire, 2005). *Eriosema* is morphologically similar and phylogenetically closely related to *Rhynchosia* (Greear, 1970; Doyle & Doyle, 1993; Bruneau et al., 1995; Kajita et al., 2001; Egan et al., 2016; LPWG, 2017). These two genera are the largest and the only pantropical members of Cajaninae, all other genera of the subtribe occurring exclusively in the Palaeotropics, including Oceania (Greear, 1970, 1978). To date, there is no comprehensive molecular phylogenetic study focused on *Eriosema* or on any other genus of Cajaninae. There are, however,

some studies focused on population genetics of *Cajanus* (Nadimpalli et al., 1993; Kassa et al., 2012; Saxena et al., 2014; Kaila et al., 2016).

Eriosema spp. are mainly distributed in tropical savannas, grasslands and mountainous areas of Africa and in Brazilian cerrado (Fig. 1; Greear, 1970; Cândido et al., 2019). The main life form and morphological characteristics of *Eriosema* spp. that have adapted to these environments are their herbaceous, sub-shrubby or shrubby habit (Fig. 2A, C, G), frequently with a specialized underground root system such as a woody rootstock (Fig. 2A–C), the apex of which has a bud-generating portion often called a xylopodium, and an abundance of trichomes throughout the plant (Fig. 2D–H). In addition, secretory structures that are peculiar to Cajaninae are usually present, such as bulbous-based trichomes, vesicular glands and secretory-based trichomes (Greear, 1970; Lackey, 1978; Cândido et al., 2016; Vargas et al., 2018; Cândido et al., 2019).

Savannas evolved in close association with frequent fires and native megafaunal herbivory (Buisson et al., 2019; Dwyer et al., 2000). As a result, these environments have become resilient and often dependent on these disturbances, and the plants that occur in these adverse environments have evolved peculiar adaptive morphologies (Coutinho, 1990; Bond & Keeley, 2005; Beerling & Osborne, 2006; Simon et al., 2009; Ratnam et al., 2011; Parr et al., 2014; Veldman et al., 2015; Bond & Zaloumis 2016; Buisson et al., 2019). This type of vegetation began to evolve c. 30 Mya, with the emergence of C₄ photosynthetic grasslands (Scholes & Archer, 1997; Edwards et al., 2010; Van der Werf et al., 2010; Parr et al., 2014). Fossil records of pollen and phytoliths show an expansion of C₄ grasses between 6 and 8 Mya (Cerling et al., 1997; Parr et al., 2014).

Tropical savanna and grassland environments cover c. 20% of the global land surface (Parr et al., 2014; Buisson et al., 2019). In Brazil, cerrado is a tropical moist savanna and is the most diverse tropical savanna in the world; it is considered a biodiversity hotspot with a high number of endemic species, the existence of which is extremely threatened by habitat loss (Myers et al., 2000; Hoekstra et al., 2005; Simon et al., 2009; Sano et al., 2010) (Fig. 1).

In the savannas, there is a diverse array of habitats, including woodland, wooded-grassland,




Figure 1. Tropical savannas and grassland environments in Africa and South America, the natural habitat of *Eriosema*. A, Bergville (KwaZulu-Natal), South Africa. B, D, Rio Preto State Park (Minas Gerais), Brazil. C, Ingomankulu (KwaZulu-Natal), South Africa. E, G, Umtamvuna Nature Reserve (KwaZulu-Natal), South Africa. F, Chapada dos Veadeiros National Park (Goiás), Brazil. H, Cerrado of São João da Aliança (Goiás), Brazil. Photographs: A, C, E, David Styles; B, D, Luísa M.P.A. Bezerra; F, H, Elisa Cândido; G, Braam Van Wyk.



Figure 2. *Eriosema* morphology adapted to the savanna. A–C, *E. crinitum* showing the morphology of the underground root system. D, *E. defoliatum* showing trichomes on the inflorescence and flowers. E, *E. floribundum* showing trichomes throughout the plant. F, *E. irwinii* showing trichomes on mature fruits. G, The small subshrub habit of *E. macrostipulatum*. H, *E. rufum* var. *macrostachyum* showing trichomes on fruits. Photographs: A–C, E, Ambar Soldevila; D, F, Elisa Cândido; G, Marcelo Simon; H, Isa L. Morais.

bushy-grassland and grassland (Lock, 2006), in which *Eriosema* spp. grow. These habitats form a gradient of woody vegetation ranging from a canopy cover of up to 40%, to open grassland with a limited occurrence of shrubs (White, 1983; Pennington, Prado & Pendry, 2000; Lock, 2006; Cândido *et al.*, 2019). The characteristics of these habitats include a single dry season lasting more than four months, tussock-forming and xeromorphic narrow-leaved grasses, a single shrub or grass layer, often the occurrence of regular fires, natural grasslands in sites with seasonal waterlogging, shallow soils and a high metallic ion concentration (Lock, 2006). At higher elevations in the savanna biome, a vegetational mosaic of mountain-top 'islands' dominated by seasonal drought and rocky or sandy soils of low fertility occurs. In the Neotropics, these rocky 'islands' are referred to as 'campos rupestres' (literally rocky fields), and the Espinhaço Range in eastern Brazil is one of the main areas for this habitat. Many *Eriosema* spp. occur in campo rupestre vegetation (Gear, 1970; Fortuna-Perez *et al.*, 2013b, 2017, 2018; Cândido *et al.*, 2014a, b, 2019) (Fig. 1D). In tropical Africa, similar vegetation to the Espinhaço Range is found in the Chimanimani mountains on the Zimbabwe–Mozambique border (Lock, 2006) where one of the most abundant shrubs is *Eriosema montanum* Baker f. (Phipps & Goodier, 1962).

Given the lack of a comprehensive phylogenetic evaluation of *Eriosema*, its intrinsic ecological relationship with savanna and grassland environments and its pantropical distribution, we decided to construct a molecular phylogenetic tree for the genus, examining its infra- and intergeneric relationships. Also, we estimated the timing of diversification of the genus in tropical savannas. We addressed the following questions. (1) Can we confirm that *Eriosema* is a monophyletic group? (2) What are the relationships of *Eriosema* with other genera in *Cajaninae*? (3) How can the transatlantic disjunct distribution of *Eriosema* be explained? (4) Where and when did major diversification events occur in the tropical savanna-adapted genus *Eriosema*?

MATERIAL AND METHODS

TAXON SAMPLING

Ninety-three species (98 accessions) of *Eriosema* were included in this study, of which 58 species are from Africa, 34 are from America and *E. chinense* is from Asia. The sampling covers the entire distribution range of the genus and represents 62% of the species (Appendix S1). To assess relationships among genera in *Cajaninae*, we also sampled one *Chrysoscias* sp., the monospecific genus *Bolusafrax*, two species each

of *Adenodolichos* and *Paracalyx*, three species each of *Cajanus*, *Dunbaria* and *Flemingia*, and 32 species of *Rhynchosia* (Appendix S1). *Galactia striata* (Jacq.) Urb. and *Camptosema grandiflorum* Benth. from subtribe Diocleinae were included as outgroups (Queiroz *et al.*, 2015). The monospecific genus *Carrisosa* (*C. angolense* Baker f.), was not included in this study as we were unable to access material for DNA analysis. This species was known only from a type specimen housed in the herbarium of the University of Coimbra (COI), Portugal. Recently, this species has been collected in Angola by David Goyder (Royal Botanic Gardens, Kew) and we have received this material. It will be included in our next manuscript on the phylogenetics of *Cajaninae*.

DNA EXTRACTION, AMPLIFICATION AND SEQUENCING

Most samples for DNA extraction were obtained from herbarium collections (Appendix S1). Samples of some species that occur in Brazil and a few from Africa were collected in the field and stored in silica gel. Total genomic DNA was isolated using the NucleoSpin Plant II kit (Macherey-Nagel, Düren, Germany), based mainly on the manufacturer's protocols, but with a modification at the beginning of the procedure, i.e. adding 4 µL of RNase A stock solution, and for herbarium samples we left the heater on for c. 3 h. About 1 cm² of dried leaf material of each sample was ground with a pestle and mortar using liquid nitrogen or in a Geno Grinder 2010 machine (SPEX Sample Prep LLC, Metuchen, NJ, USA) for 2 min at 2 rpm. Before the final selection of plastid DNA markers for phylogenetic analyses, we evaluated various markers to examine their phylogenetic informativeness among selected closely related *Eriosema* spp. (data not shown). After an initial screening step, we selected *rpl32-trnL* and *trnQ-5'rps16* intergenic spacers for subsequent steps. These were amplified using *trnL* (UAG)/*rpl32-F* and *trnQ* (UUG)/*rps16x1* primers, respectively (Shaw *et al.*, 2007). To amplify the ITS region (including ITS1 and ITS2 spacer regions and the 5.8S ribosomal subunit) of nuclear ribosomal DNA (nrDNA), primer pairs 17SE and 26SE were used (Sun *et al.*, 1994) (Appendix S2).

The PCR reactions were prepared in 25 µL solution containing 2 µL 10× PCR buffer, 1.5 µL 50 mM MgCl₂, 1 µL 2.5 mM dNTPs, 1 µL KB Extender (Invitrogen), 1 µL 5%DMSO, 0.5 µL bovine serum albumin, 0.5 µL betaine, 2 µL each primer (5 pM), 0.2 µL Platinum Taq DNA polymerase (Invitrogen) and 1.5 µL DNA template, adjusted with deionized water. The amplification program for the plastid regions consisted of 35 cycles of 95 °C for 1 min, 55 °C for 1 min and 72 °C for 1 min and a final extension 72 °C for 7 min and 10 °C

constant at the end of the reaction. The ITS region was amplified with 11 cycles using the touchdown strategy: 94 °C for 3 min, 94 °C for 1 min; –1 °C per cycle from 65 °C; 72 °C for 1 min; 15 cycles back to step 2, 94 °C for 1 min; 50 °C for 1 min; 72 °C for 4 min; 21 cycles back to step 2; 72 °C for 1 min and 15 °C constant at the end of the reaction (Appendix S2). After the PCR, 2 µL of the PCR solution was run on 1% agarose gel to detect successful amplification. The amplified PCR products were purified using the NucleoSpin Gel and PCR clean-up kit (Macherey-Nagel, Düren, Germany) and sent for sequencing with the same primers used for the amplification. DNA sequencing reactions were performed in an automatic sequencer ABI 3730xl (Macrogen Inc., Seoul, South Korea).

ALIGNMENT AND PHYLOGENETIC ANALYSIS

Forward and reverse sequences were assembled and aligned into contigs with Geneious v.8.1.9 (Kearse *et al.*, 2012). All sequences generated in this study were deposited in GenBank (Appendix S1). A BLAST search (Altschul *et al.*, 1990) was conducted for all sequences to check for possible contamination. DNA regions were aligned using MAFFT v.7.369b (Katoh & Standley, 2013) using the L-INS-i method with ‘--localpair’ and ‘--maxiterate 1000’ and ‘--adjustdirection’ options. Alignments were corrected and edited by eye in AliView v.1.23 (Larsson, 2014).

Our phylogenetic analyses are based on maximum likelihood (ML) and Bayesian inference (BI) approaches. We inferred the maximum-likelihood tree using IQ-TREE v.1.6.12 (Nguyen *et al.*, 2015). ModelFinder (Kalyaanamoorthy *et al.*, 2017) tested 286 DNA models for accurate phylogenetic estimates. We obtained branch supports with the ultrafast bootstrap feature, ‘-bb 10000’ (Hoang *et al.*, 2018), and performed the SH-like approximate likelihood ratio test (SH-aLRT) (Guindon *et al.*, 2010) by using the ‘-alrt 1000’ option to assess branch supports. We ran IQ-TREE on each marker independently, concatenated plastid regions and concatenated plastid regions plus ITS (four data matrices in total). BI was performed using a Markov chain Monte Carlo (MCMC) method, as implemented in MrBayes, v.3.2.6 (Ronquist, Huelsenbeck & Teslenko, 2012) using parallel version (mb-mpi). We used MrModeltest v.2.3 (Nylander, 2004) to find the best substitution model for each region separately. For *rpl32-trnL* and ITS SYM+G and for the *trnQ-5'rps16* HKY+G were selected based on the Akaike information criterion (AIC) (Akaike, 1974). Two independent runs were performed on four data matrices as mentioned above. For each BI run, we set the number of generations to 20 million, sampling every 2000 generations, discarding 20% of the trees as burn-in. All runs were inspected to check that the

average standard deviation of split frequencies was < 0.1 (Ronquist *et al.*, 2011). Convergence of MCMC inference was evaluated using Tracer v.1.7.1 (Rambaut *et al.*, 2018). The phylogenetic analyses were completed on the Smithsonian Institution High-Performance Cluster (<https://doi.org/10.25572/SHPC>).

DIVERGENCE TIME ANALYSIS

Estimates of divergence time of lineages in *Cajaninae* were obtained using BEAST v.1.8.2 (Drummond *et al.*, 2012) using the ITS matrix. A literature search revealed no fossil records of *Cajaninae* for use as a calibration point, and calibration of the root of the tree was implemented based on the estimation from Lavin, Herendeen & Wojciechowski (2005). A normal distribution was applied with the mean of 27.8 Mya and a standard deviation of 1.6 according to Lavin *et al.* (2005). An uncorrelated relaxed molecular clock with a lognormal distribution of rates and a Yule speciation model was used. Two independent runs, each with 50 million generations were performed sampling every 5000 generations. A maximum clade-credibility (MCC) tree was retrieved using TreeAnnotator v.1.8.2 (Drummond *et al.*, 2012) after discarding the first 20% of trees as burn-in. We used Tracer v.1.7.1 (Rambaut *et al.*, 2018) to assess Effective Sample Size (ESS) and convergence.

RESULTS

DATA MATRICES AND PHYLOGENETIC ANALYSIS

Our dataset includes 140 species from nine genera of *Cajaninae* and *Galactia striata* and *Camptosema grandiflorum* from *Diocleinae* as outgroups. In this study, we produced 391 new DNA sequences (123, 141 and 127 for ITS, *rpl32-trnL* and *trnQ-5'rps16*, respectively; Appendix S1). The alignment lengths of ITS, *rpl32-trnL* and *trnQ-5'rps16* were 818, 832 and 789 bp, respectively, of which 460, 154 and 187 bp were potentially parsimony informative (Appendix S3).

The ML and BI analyses of the combined dataset (ITS + plastid regions) revealed that *Cajaninae* and *Adenodolichos*, *Dunbaria*, *Flemingia* and *Cajanus* are monophyletic (BP = 100, PP = 1.00, Fig. 3A, Appendix S4). *Rhynchosia* was recovered as polyphyletic and emerged as three separate lineages. *Rhynchosia volubilis* Lour. (clade I), the type species of the genus, alone is sister to the *Cajanus* clade, although the bootstrap support for their ancestral node is only moderate (BP = 95, Fig. 3A). The second lineage of *Rhynchosia*, clade II (*Rhynchosia* Old World + (*Rhynchosia* Old World + *Paracalyx*) + *Rhynchosia* AM), in which most of the sampled species are placed,




Figure 3. A scheme of the phylogenetic tree for *Eriosema* from maximum likelihood analysis of the ITS/5.8S and plast DNA dataset (*rpl32-trnL* and *trnQ-5'rps16*). A, Detail of the other genera of Cajaninae. B, Detail of *Eriosema*. Numbers along branches are bootstrap values and posterior probabilities.

also includes *Paracalyx* with moderate support (BP = 81; Fig. 3A), and a clade of American species (*Rhynchosia* AM) (BP = 90; Fig. 3A).

Chrysoscias and *Bolusafr* form a clade (BP = 99, PP = 1.00, Fig. 3A; Appendix S4) that is sister to the third lineage of *Rhynchosia* [*Rhynchosia* AF (Africa)],



Figure 3. Continued.

clade III (Fig. 3A, BP=99, PP=1.00). This clade includes the African species *Rhynchosia adenodes* Eckl. & Zeyh., *R. fleckii* Schinck, *R. aureovillosa* Hauman var. *humbertii* Hauman, *R. sublobata* (Schumach.) Meikle and *R. nyasica* Baker (Appendix S4). Together, clades III and *Chrysoscias* + *Bolusafr* are sister to the *Eriosema* clade (Fig. 3A). In all analyses, *Eriosema* was resolved as monophyletic with maximum support (BS = 100, PP = 1.00, Fig. 3A, B, Appendix S4). However, interspecific relationships in *Eriosema* so far have not been fully resolved. Inside the *Eriosema* clade, one group comprising only American species (subclade IV) has moderate bootstrap support (BP = 78; Fig. 3B) and there is no good resolution of species relationships inside this American group. Likewise, there is no intra-cladal support within the groupings of African species. The only Asian species, *Eriosema chinense* [*Eriosema* AS (Asian)], formed part of clade V with African *E. psoraleoides* (Lam.) G. Don (BP = 99, PP = 1.00, Fig. 3B), and this clade is the sister group of the rest of *Eriosema*.

DIVERGENCE TIME ANALYSIS

Age estimates are presented for relevant crown nodes (mean and 95% credibility intervals). Cajaninae were estimated to have diversified 24.29–30.64 Mya (Fig. 4A, Appendix S5). The *Adenodolichos* clade diverged first (3.25–6.78 Mya) in Cajaninae (Fig. 4A, Appendix S5). *Flemingia* and *Dunbaria* diverged c. 2.09–4.45 and 4.09–7.21 Mya, respectively. The diversification time of *Cajanus* + *Rhynchosia volubilis* is estimated at 11.53–17.70 Mya (Fig. 4A, Appendix S5). *Cajanus* itself diversified 3.61–6.61 Mya (Fig. 4A, Appendix S5). The clade that includes most *Rhynchosia* spp. + *Paracalyx* is estimated to date from 11.18–16.91 Mya (Fig. 4A, Appendix S5). The African clade of *Rhynchosia fleckii*, *R. adenodes*, *R. nyasica*, *R. aureovillosa* var. *humbertii* and *R. sublobata* plus *Bolusafr* *bituminosa* (L.) Kuntze and *Chrysoscias calycina* E. Mey. diversified 10.59–16.44 Mya (Fig. 4A, Appendix S5). Stem and crown group ages of *Eriosema* were estimated as 12.25–18.11 and 6.50–10.79 Myr, respectively (Fig. 4B, Appendix S5). Group 1 of *Eriosema* comprises species that occur in Africa and has an estimated crown group age of 4.41–6.68 Myr (Fig. 4B, Appendix S5). The group that includes American *Eriosema* spp. (Group 2) diverged from African species c. 3.56–5.78 Mya (Fig. 4B, Appendix S5). Species in this group occur mainly in the cerrado and campos rupestres vegetation types in Brazil, especially in the states of Goiás and Minas Gerais. The age of the crown clade formed by the only Asian species, *E. chinense*, and the African species *E. psoraleoides* (Group 3) was estimated at 1.73–4.64 Myr (Fig. 4B, Appendix S5).

DISCUSSION

MONOPHYLY OF *ERIOSEMA*

Our phylogenetic study confirms the monophyly of *Eriosema* (Fig. 3B, Appendix S4). Previous phylogenetic studies focusing on Phaseoleae or the entire Fabaceae (e.g. Doyle & Doyle, 1993; Bruneau *et al.*, 1995; Kajita *et al.*, 2001; Egan *et al.*, 2016; LPWG, 2017), albeit including a limited number of *Eriosema* spp., also supported the monophyly of the genus. Among these studies, the most recent and well sampled so far, based on the plastid *matK* gene (LPWG, 2017), included only 12 *Eriosema* spp. (8% of the genus). In our initial screening of loci for this project, the *matK* gene was sequenced for 13 accessions of *Eriosema* spp., but we found the genetic diversity and parsimony informativeness of *matK* to be lower than the two plastid markers that we used in this study.

Interspecific relationships in *Eriosema* have not yet been resolved (Fig. 3B, Appendix S4), even using some selective plastid DNA markers in combination with nrITS. In all our analyses, there is a tendency for species to cluster in geographical groups, although not always with strong support, with the American species (BP 78%, Fig. 3B) and the African species forming two large groups (Fig. 3B, Appendix S4). In some analyses, the African species are the first diverging group in the *Eriosema* clade, possibly suggesting an African origin of the genus, but our results are not able to fully confirm this hypothesis. One possible explanation for this may be related to the recent and rapid diversification of *Eriosema* (10.79–6.50 Mya) (Fig. 4B, Appendix S5). Future studies that focus on the evolution of *Eriosema* should incorporate phylogenomic techniques that would obtain high resolution phylogenetic trees, which should better explain the morphological and biogeographical relationships in the genus.

Morphologically, *Eriosema* shares with other genera of Cajaninae the papilionaceous flowers (yellow or red), unifoliate or trifoliate leaves, fruits elastically dehiscent, peculiar secretory structures, absence of canavanine (except in *Adenodolichos*) and absence of bracteoles (except in *Adenodolichos*) (Grear, 1970; Lackey, 1978; Lewis *et al.*, 2005; Cândido *et al.*, 2016; Vargas *et al.*, 2018). Clade III, which is the sister group of *Eriosema*, comprises African species of *Rhynchosia*, *Bolusafr* and *Chrysoscias* (Fig. 3A) and morphologically these three genera are similar. This similarity was also observed by Lackey (1981), who described *Bolusafr* as a 'viscid *Rhynchosia*-like vine'. The representatives of clade III are usually trifoliate, have a climbing habit, flowers yellow and red and calyx lobes exceeding the corolla (Vargas, 2019).

Of the *Eriosema* spp. occurring in the Americas, those belonging to the *Eriosema campestre* Benth. complex need nomenclatural attention. Taxa of




Figure 4. Analysis performed on the maximum clade credibility tree from divergence dating analysis (BEAST) using lognormal uncorrelated relaxed clock models for the combined data (ITS/5.8S, *rpl32-trnL* and *trnQ-5rps16*). A, Detail of the genera of Cajaninae (except *Eriosema*). B, Detail of *Eriosema*. Blue bars represent 95% confidence intervals (CI) for the estimated mean dates. Posterior probabilities are indicated at nodes.

this complex [*E. campestre* Benth. var. *campestre*, *E. campestre* var. *macrophyllum* (Grear) Fortunato, *E. campestre* var. *pubescens* (Chodat & Hassl.) Grear and *E. campestre* var. *delicatula* Fortunato]

do not form a group in our analyses (Fig. 3B) and the varieties would be better recognized as distinct species, as also supported by the anatomical studies of Seixas, Fortuna-Perez & Rodrigues (2019). Another




Figure 4. Continued.

complex that requires attention is *E. simplicifolium* (DC.) G.Don with two varieties: var. *simplicifolium* and var. *micranthum* Grear, because these appeared in different lineages in our analyses (Fig. 3B). *Eriosema simplicifolium* var. *micranthum* has an

erect habit (prostrate in var. *simplicifolium*), smaller flowers and longer inflorescences than the typical variety (Cândido *et al.*, 2019). Fortunato (1993) synonymized *E. simplicifolium* var. *micranthum* under the typical variety, but our study clearly

shows that *E. simplicifolium* var. *micranthum* should be elevated to species rank. Also, *E. pycnanthum* Benth. var. *pycnanthum* and *E. pycnanthum* var. *veadeirense* Grear appeared in different lineages in the phylogenetic tree (Fig. 3B). Cândido *et al.* (2019) stated that the varieties of *E. pycnanthum* are distinguished morphologically mainly by the length of the inflorescence (3.5–6.5-cm long in *E. pycnanthum* var. *pycnanthum* vs. 1.5–3.0-cm long in *E. pycnanthum* var. *veadeirense*) and leaflet texture (membranaceous to chartaceous in *E. pycnanthum* var. *pycnanthum* vs. thin-coriaceous to coriaceous in *E. pycnanthum* var. *veadeirense*) (Cândido *et al.*, 2019). Our analyses suggest that *E. pycnanthum* var. *veadeirense* would be better recognized at the rank of species.

HISTORICAL BIOGEOGRAPHY OF *ERIOSEMA*

Eriosema has a pantropical, intercontinental disjunct distribution, a common pattern in plants in general (Thorne, 1972; Givnish & Renner, 2004; Bartish *et al.*, 2011) and in Fabaceae (e.g. Fortuna-Perez *et al.*, 2013a; Vatanparast *et al.*, 2013; Moura *et al.*, 2016). According to molecular dating analyses, *Eriosema* originated in the Miocene, with an estimated crown-group age of 10.79–6.50 Myr and a stem group age of 18.11–12.25 Myr (Fig. 4B, Appendix S5). There are two major groups of species in *Eriosema*. Group 1 includes species that occur in Africa and has an estimated crown group age of 6.68–4.41 Myr (Fig. 4B, Appendix S5). This lineage comprises the greatest species diversity in the genus and is morphologically characterized by having racemose inflorescences, flowers occurring along the inflorescence rachis and, generally, inflorescences longer than the leaves (Fig. 5). A xylopodium, a specialized underground root system, is a synapomorphic character for species of this group. Group 2, which encompasses the majority of the New World species, has a divergence time of c. 5.78–3.56 Mya in the late Miocene (Fig. 4B, Appendix S5). It comprises species that occur mainly in the cerrado and campos rupestres vegetation in Brazil (Fig. 1B, D, F, H), including *E. brevipes* Grear (Fig. 6B), *E. defoliatum* Benth. (Fig. 6F), *E. elegans* Fort.-Perez & M.J.Silva (Fig. 6G), *E. floribundum* Benth. (Fig. 6H), *E. hatschbachii* Fort.-Perez & G.P.Lewis (Fig. 6K) and *E. irwinii* Grear (Fig. 6L), and a few species occurring in open areas of Mexico, Central and South America. The species in this clade have a racemose inflorescence and flowers crowded at the apex of the inflorescence, generally not surpassing the leaves in length (Fig. 6). Characters shared between species of the ‘Old World’ and ‘New World’ include the uni- to trifoliate leaves, the abundant, long trichomes and glands throughout the plant and the presence of a specialized underground root system (Fig. 2A–C).

According to a previous study (Li *et al.*, 2013), as observed in this study for Cajaninae, during the late Oligocene some lineages of phaseoloid members of Fabaceae (mainly woody plants) underwent a rapid adaptive radiation, probably in response to the warming and increased aridity during this geological period; taxa rapidly evolved to exploit new niche opportunities in Africa, Asia and Australia. According to our results, *Eriosema* probably first appeared in Africa or Asia, during the late Miocene, (Fig. 3B, Appendix S4) and the great diversification in Africa and America was subsequently driven by rapid adaptation to these new ecological opportunities and the range expansion of C₄ grasses.

Our results strongly support the Old World as being the centre of origin of *Eriosema*. The chronogram resulting from the dating analysis indicates that the unique shifting event of *Eriosema* from the Old World to America occurred c. 6.89–4.64 Mya (Fig. 4B, Appendix S5). Considering the age of this migratory event and the estimates of the young age of the African-Neotropical disjunctions in *Eriosema*, continental drift or boreotropical migration can be discarded as explanations for this ampho-Atlantic migration. Some studies have reported local ants or beetles as the dispersal agents of *Eriosema* spp. (van der Pijl, 1982; Kergoat *et al.*, 2011). Transoceanic long-distance dispersal from Africa/Asia to the Neotropics is the most likely explanation. Several molecular dating studies report that the current distribution of taxa has occurred after these transoceanic long-distance dispersal events; these include studies in Fabaceae (e.g. Herendeen, Crepet & Dilcher, 1992; Lavin *et al.*, 2004; Lavin, Herendeen & Wojciechowski, 2005; Schrire *et al.*, 2005; Pennington, Richardson & Lavin, 2006; Simon *et al.*, 2011; Fortuna-Perez *et al.*, 2013a; Snak *et al.*, 2016; Cardoso *et al.*, 2017; Tosso *et al.*, 2018; Vasconcelos *et al.*, 2020). Future studies of *Eriosema* should explore in more depth the biogeographical hypotheses outlined here, and discuss in more detail the ancestral areas and geoclimatic events coexisting at the time of *Eriosema* speciation.

DIVERSIFICATION AND SAVANNA-ADAPTED MORPHOLOGY OF *ERIOSEMA*

Tropical savanna and grassland environments evolved c. 30 Myr and expanded worldwide in the late Miocene. Their evolution was simultaneous in Africa and America, and coincided with the substantial increase in C₄ grasses, which currently account for 25% of all terrestrial photosynthesis (Scholes & Archer, 1997; Cerling *et al.*, 1997; Keeley & Rundel, 2005; Beerling & Osborne, 2006; Edwards *et al.*, 2010; Van der Werf *et al.*, 2010; Parr *et al.*, 2014; Maurin *et al.*, 2014; Pennington & Hughes, 2014). In accordance with




Figure 5. African *Eriosema* species. A, *E. dregei*. B, *E. ellipticifolium*. C, *E. guenzii*. D, *E. luteopetalum*. E, *E. populifolium*. F, *E. preptum*. G, *E. rossii*. H, *E. salignum*. I, *E. umtamvunense*. Photographs: David Styles.




Figure 6. American *Eriosema* species. A, *E. benthamianum*. B, *E. brevipes*. C, *E. campestre* var. *campestre*. D, *E. congestum*. E, *E. crinitum* var. *crinitum*. F, *E. defoliatum*. G, *E. elegans*. H, *E. floribundum*. I, *E. glabrum*. J, *E. heterophyllum*. K, *E. hatschbachii*. L, *E. irwinii*. M, *E. laxiflorum*. Photographs: A–C, F, G, L, Elisa Cândido; D, E, I, Ana Paula Fortuna-Perez; H, Ambar Soldevila; J, Gustavo H. Shimizu; M, Marcos Silva.

the age estimate of *Eriosema* (10.7–6.5 Myr; Fig. 4B, Appendix S5) the diversification of the genus occurred in parallel with the savanna biome expansion. Time-calibrated phylogenetic trees of other plant taxa also suggest that savanna lineages started to diversify < 10 Mya and that the rapid diversification of these lineages coincided with the increase and dominance of C₄ grasses in tropical savannas around the world (e.g. Simon *et al.*, 2009, 2011; Simon & Pennington, 2012; Souza *et al.*, 2013; Trovó *et al.*, 2013; Maurin *et al.*, 2014; Souza-Neto, Cianciaruso & Collevatti, 2015; Rando *et al.*, 2016).

Eriosema is a recent, tropical savanna-adapted subshrub/shrub group and the current distribution of its representatives occurs, essentially, in the tropical savannas and grasslands of the African continent and the Brazilian cerrado (Grear, 1970; Cândido *et al.*, 2019) (Fig. 1). *Eriosema*, like other genera that evolved and occur in these environments, displays specialized morphological, physiological and ecological survival strategies. Morphologically, we highlight the underground root system and the presence of trichomes throughout the plant (Fig. 2), including secretory structures (bulbous-based trichomes, vesicular glands and secretory-based trichomes) (Grear, 1970; Lackey, 1978; Cândido *et al.*, 2016; Vargas *et al.*, 2018; Cândido *et al.*, 2019). Grear (1970) named the underground specialized root system a xylopodium, and this terminology has been followed by subsequent studies (e.g. Stirton, 1975, 1981, 1986; Stirton & Gordon-Gray, 1978; Grear, 1978; Miotto, 1988; Fortunato, 1993, 1999; Fortuna-Perez *et al.*, 2013, 2017, 2018; Cândido *et al.*, 2014a, b, 2016; Oliveira, Fortuna-Perez & Silva, 2018). However, only now are detailed anatomical studies being conducted to investigate the variation in the thickened underground root system found in *Eriosema* spp. A preliminary survey has found that in some species this root system can take the form of a xylopodium, a tuberous or non-tuberous root (often daucate and with or without smaller side appendages) or a rhizome (Piva *et al.*, 2020; Stirton, C, unpubl. data). Future investigations should focus on the ecological functions of this underground system. Root systems in African savannas are often different from those in coastal and upland grassland systems (Stirton, C, unpubl. data).

Regarding the trichome types encountered in *Eriosema*, Vargas *et al.* (2018) carried out the most comprehensive anatomical and micro-morphological study of representatives of Cajaninae, confirming that only vesicular glands are a unifying character of the subtribe. Another recent study analysed the ultrastructure and histochemistry of glandular trichomes in the leaflets of species of Cajaninae, including *E. rufum* (Kunth) G. Don and

E. simplicifolium, two species widely distributed in the American continent. The authors suggested that glandular trichomes and their exudates (lipids and phenolic compounds) act as a defence against herbivory and also help to prevent damage by ultraviolet radiation. This is especially important for the survival of *Eriosema* spp. that have diversified in environments with high light intensity and high temperatures (Vargas *et al.*, 2019).

Although there exists no published reproductive phenological study of *Eriosema* spp., there are some examples reporting the intrinsic ecological relationship between *Eriosema* spp. and savanna environments. *Eriosema laxiflorum* Harms (Fig. 6M), a small subshrub (10–26-cm tall) with an erect habit and a tufted (caespitose) appearance, is endemic to the cerrado of central Brazil and only blooms directly after fire. All flowering individuals of this species collected by us were encountered in recently burnt vegetation (Cândido *et al.*, 2019). *Eriosema kraussianum* Meisn. is a small hysteroanthous pyrophyte from the grasslands of Kwa-Zulu Natal in South Africa that only flowers after fire. This post-fire flowering phenomenon has also been reported for other angiosperm species in similar environments (Coutinho, 1976; Conceição & Orr, 2012; Conceição *et al.*, 2013; Fidelis & Blanco, 2014). Two species which are endemic to the Brazilian cerrado, *Eriosema defoliatum* Benth. (Fig. 6F) and *Eriosema congestum* Benth. (Fig. 6D), only flower and fruit in the dry season when they are leafless, so that leaves and flowers are rarely collected together (Cândido *et al.*, 2019). This reproductive trait needs to be better investigated.

MORPHOLOGICAL AND PHYLOGENETIC RELATIONSHIPS BETWEEN *ERIOSEMA* AND *RHYNCHOSIA*: THE TWO MOST SPECIES-RICH GENERA OF CAJANINAE

Rhynchosia, the largest genus of Cajaninae, comprises c. 230 species (Schrire, 2005). The genus is not, however, monophyletic, and one clade of African species is the sister group of *Eriosema*. Egan *et al.* (2016) also demonstrated *Rhynchosia* to be paraphyletic, although they only sampled two species in their study of *Pueraria* DC. (subtribe Glycininae, tribe Phaseoleae). The close morphological and phylogenetic relationship between *Eriosema* and *Rhynchosia* has also been demonstrated in other previous studies, albeit with lower sampling of the two genera (Doyle & Doyle, 1993; Bruneau *et al.*, 1995; Kajita *et al.*, 2001; LPWG, 2017). Our phylogenetic analyses support the two genera as distinct evolutionary lineages (Fig. 3, Appendix S4).

Eriosema and *Rhynchosia* together comprise c. 380 species, representing the large majority of species in Cajaninae; they are the only genera in the subtribe

that have species in the New World. The two genera share several morphological characteristics, including typically papilionaceous and mostly yellow flowers; two-seeded dehiscent fruits; uni- or trifoliate leaves and stipules usually lanceolate to triangular (Oliveira *et al.*, 2018; Bezerra *et al.*, 2019a, b, c; Cândido *et al.*, 2019). Traditionally, the two genera have been separated in morphological studies mainly by the apparent differing insertion point of the funicle in relation to the hilum of the seed (Baker, 1895; Grear, 1970; 1978; Miotto, 1988; Cândido *et al.*, 2014a). The distinction between these genera based only on this character is not, however, consistent or reliable. A suite of characters is needed to reliably separate *Rhynchosia* from *Eriosema*, including habit, stipules (free or connate), presence or absence of stipels, inflorescence type, seed colour, hilum shape, extension of the strophiole and fruit constriction between the seeds (Cândido *et al.*, 2019).

In southern Africa, the two genera are easily identified by the shape of the fruits (box-like rectangular and shaggy-haired in *Eriosema* versus attenuate at both ends and less prominently hairy in *Rhynchosia*), the seedlings (epicotyls shaggy-haired in *Eriosema* and finely pubescent in *Rhynchosia*); the presence of dense glands on the keel petals and back of the standard petal (ubiquitous in *Eriosema*, rare in *Rhynchosia*) and the patterns formed on leaflets by species of Agromyzidae (leaf-miners) and species of Bruchidae (seed weevils) that attack the fruits (Stirton, C, unpubl. data). These traits might provide useful diagnostic characters if surveyed across the geographical distribution of the two genera. In addition, although the two genera have an essentially pantropical distribution, some *Rhynchosia* spp. extend to temperate regions and forest environments, whereas *Eriosema* spp. are restricted to savanna, savanna-like environments or grasslands (Fig. 1) (Grear, 1978; Bezerra *et al.*, 2019a, b; Cândido *et al.*, 2019).

INTERGENERIC RELATIONSHIPS IN CAJANINAE

Cajaninae, with c. 500 species, are the largest subtribe of Phaseoleae (Fig. 3, Appendix S4). They are included in core Phaseoleae *sensu* Schrire (2005) with Psoraleeae, Glycininae, Erythrinae and the genus *Butea* Roxb. ex Willd. (Egan *et al.*, 2016). In our study we sampled nine out of the ten genera traditionally placed in Cajaninae (Appendix S1). Material of the monospecific *Carrissoa* Baker f. (*C. angolense* Baker f.) will be included in future studies of the subtribe. *Adenodolichos* is strongly supported as monophyletic (Fig. 3A, Appendix S4). Bruneau *et al.* (1995) resolved *Adenodolichos* as sister to all other representatives of Cajaninae, but in our study its position is not fully resolved. *Adenodolichos* is the only genus in the

subtribe that possesses bracteoles on its flowers and the chemical compound canavanine, and Lackey (1977, 1981) considered it as anomalous in Cajaninae. Egan *et al.* (2016) found that canavanine production is the ancestral condition in Phaseoleae although its presence has been lost and regained more than once in lineages of the tribe. *Adenodolichos* shares vesicular glands with other members of Cajaninae, and these glands are considered as a unifying character of the subtribe (Vargas *et al.*, 2018). The other genera of Cajaninae that have been recovered as monophyletic are *Cajanus*, *Dunbaria* and *Flemingia* (Fig. 3A, Appendix S4). *Rhynchosia* is separated into two distinct lineages; here we included *R. volubilis* (the type species, native to Asia and sister to the *Cajanus* clade) and *C. cinereus* (F.Muell. ex Benth.) F.Muell., *C. cajanifolius* (Haines) Maesen and *C. scarabaeoides* (L.) Thouars, all native to Asia and northern Australia (Fig. 3A, Appendix S4). *Paracalyx* emerged nested in a group of *Rhynchosia* spp. Egan *et al.* (2016) identified *Paracalyx* as sister to *Rhynchosia*. The species belonging to *Paracalyx* were originally included in *Cylista* Aiton and the type species of that genus is included in *Rhynchosia* (Egan *et al.*, 2016). Our study included the largest, to date, sampling of *Rhynchosia* spp. across its geographical range. Further work including a broader sample of *Cajanus*, *Paracalyx* and *Rhynchosia* and multiple accessions of *R. volubilis* is needed to completely unravel the relationships between these taxa.

CONCLUSIONS

Our study presents the first comprehensive molecular phylogenetic analysis of *Eriosema* and allied genera of Cajaninae. Our results support the monophyly of the subtribe and of *Adenodolichos*, *Eriosema*, *Dunbaria*, *Flemingia* and *Cajanus*. *Eriosema* originated in the Miocene (10.7–6.5 Mya), and its great diversification in Africa and the New World was driven by the opportunities generated by the appearance of new ecological niches and the range expansion of C₄ grasses, in parallel with the expansion of the savanna biome around the world. Future studies should focus on phylogenomic approaches to further elucidate evolutionary relationships in *Eriosema*. *Rhynchosia* is polyphyletic and should be reclassified based on its distinct evolutionary lineages. The close relationship between the evolution and diversification of *Eriosema* with the emergence of the savanna biome is clear from our analyses, and the conservation of *Eriosema* spp. is intimately connected to the conservation of this under-protected biome. Additional taxonomic, morphological, anatomical and biogeographical studies are necessary to better understand the evolution of and relationships between lineages of *Eriosema*.

ACKNOWLEDGEMENTS

This work is one of the results from the project 'Estudos taxonômicos, filogenéticos e biogeográficos no gênero *Eriosema* (DC.) Desv. (Leguminosae, Papilionoideae, Phaseoleae)', which is funded by FAPESP/Brazil - Fundação de Amparo à Pesquisa do Estado de São Paulo (process 2015/13386-0; 09/53002-5); CNPq/Brazil - Conselho Nacional de Desenvolvimento Científico e Tecnológico (process 400567/2016-4), both granted to A.P. Fortuna-Perez. The authors also thank Fundação de Amparo à Pesquisa do Estado de Minas Gerais/Brazil (process APQ-02323-12), and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) Print/UNESP (process 88887.373155/2019-00), for financial support granted to A.P. Fortuna-Perez; Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro/Brazil (process 203.007/2017) and CNPq (process 312766/2009-2) for financial support granted to V.F. Mansano; CNPq - REFLORA Project (process 563550/2010-4) for financial support granted to A. Tozzi; CAPES/Brazil and CNPq/Brazil (process 141628/2016-2) for a PhD scholarship to E.S. Cândido; the Rupert Barneby Award of The New York Botanical Garden awarded to E.S. Cândido. We thank the Biologia Vegetal post-graduate program at UNICAMP, the UNESP-Botucatu, the Instituto de Pesquisas JBRJ, the UFOP for the support to E.S. Cândido; the UFG for all the support in the field in the state of Goiás. We thank the curators and staff of all herbaria visited for providing access to specimens and photographs in their collections, and all the people who helped us in the field in Brazil. The majority of the analyses were conducted at the Smithsonian Institution (USA) High-Performance Cluster (<https://doi.org/10.25572/SIHP>).

REFERENCES

- Akaike H. 1974.** A new look at the statistical model identification. *IEEE Transactions on Automatic Control* **19**: 716–723.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990.** Basic local alignment search tool. *Journal of Molecular Biology* **215**: 403–410.
- Baker EG. 1895.** Revision of the African species of *Eriosema*. *Journal of Botany* **33**: 396–397.
- Bartish IV, Antonelli A, Richardson JE, Swenson U. 2011.** Vicariance or long-distance dispersal: historical biogeography of the pantropical subfamily Chrysophylloideae (Sapotaceae). *Journal of Biogeography* **38**: 177–190.
- Beerling DJ, Osborne CP. 2006.** The origin of the savanna biome. *Global Change Biology* **12**: 2023–2031.
- Bezerra LMPA, Cândido ES, Vargas W, Servilha JH, Monteiro TC, Fortuna-Perez AP. 2019a.** O gênero *Rhynchosia* Lour. (Leguminosae, Papilionoideae, Phaseoleae) no Brasil. *Rodriguésia* **70**: e04332017. 201.
- Bezerra LMPA, Oliveira ACS, Santos JS, Vargas W, Cândido ES, Monteiro TC, Vatanparast M, Fortuna-Perez AP. 2019b.** A new species of *Rhynchosia* (Leguminosae, Papilionoideae) from Bahia State, Brazil. *Phytotaxa* **406**: 84–90.
- Bezerra LMPA, Vargas W, Cândido ES, Monteiro TC, Vatanparast M, Fortuna-Perez AP. 2019c.** *Rhynchosia mineira* (Leguminosae: Papilionoideae), a new and critically endangered species from Minas Gerais, Brazil. *Kew Bulletin* **74**: 62.
- Bond W, Zaloumis NP. 2016.** The deforestation story: testing for anthropogenic origins of Africa's flammable grassy biomes. *Philosophical Transactions of the Royal Society B, Biological Sciences* **371**: 20150170.
- Bond WJ, Keeley JE. 2005.** Fire as a global 'herbivore': the ecology and evolution of flammable ecosystems. *Trends in Ecology & Evolution* **20**: 387–394.
- Bruneau A, Doyle JL, Doyle JJ. 1995.** Phylogenetic relationships in Phaseoleae: evidence from chloroplast restriction site characters. In: Crisp MD, Doyle JJ, eds. *Advances in legume systematics. Part 7. Phylogeny*. Kew: Royal Botanic Gardens, 309–330.
- Buisson E, Le Stradic S, Silveira FAO, Durigan G, Overbeck GE, Fidelis A, Fernandes GW, Bond WJ, Hermann JM, Mahy G, Alvarado ST, Zaloumis NP, Veldman JW. 2019.** Resilience and restoration of tropical and subtropical grasslands, savannas, and grassy woodlands. *Biological Reviews* **94**: 590–609.
- Cândido ES, Fortuna-Perez AP, Aranha Filho JLM, Bezerra LMPA. 2014a.** *Eriosema* (Leguminosae-Papilionoideae) no Sudeste do Brasil. *Rodriguésia* **65**: 885–916.
- Cândido ES, Fortuna-Perez AP, Bezerra LMPA, Aranha Filho JLM. 2014b.** A new species of *Eriosema* (Leguminosae, Papilionoideae, Phaseoleae) from Minas Gerais, Brazil. *Phytotaxa* **178**: 229–232.
- Cândido ES, Vargas W, Bezerra LMPA, Mansano VF, Vatanparast M, Lewis GP, Tozzi AMGA, Fortuna-Perez AP. 2019.** Taxonomic synopsis of *Eriosema* (Leguminosae: Papilionoideae, Phaseoleae) in Brazil. *Phytotaxa* **416**: 91–137.
- Cândido ES, Vargas W, Vatanparast M, Mansano VF, Machado SR, Fortuna-Perez AP. 2016.** A new species of *Eriosema* (Leguminosae, Papilionoideae, Phaseoleae) from Mato Grosso do Sul, Brazil, with a secretory structure novel to the genus. *Phytotaxa* **163**: 122–130.
- Cardoso D, Harris DJ, Wieringa JJ, São-Mateus WMB, Batalha-Filho H, Torke BM, Prenner G, Queiroz LP. 2017.** A molecular-dated phylogeny and biogeography of the monotypic legume genus *Haplormosia*, a missing African branch of the otherwise American-Australian Brongniartieae clade. *Molecular Phylogenetics and Evolution* **107**: 431–442.
- Cerling TE, Harris JM, MacFadden BJ, Leakey MG, Quade J, Eisenmann V, Ehleringer JR. 1997.** Global vegetation change through the Miocene/Pliocene boundary. *Nature* **389**: 153–158.
- Conceição AA, Alencar TG, Souza JM, Moura ADC, Silva GA. 2013.** Massive post-fire flowering events in a

- tropical mountain region of Brazil: high episodic supply of floral resources. *Acta Botanica Brasilica* **27**: 847–850.
- Conceição AA, Orr BJ. 2012.** Post-fire flowering and fruiting in *Vellozia sincorana*, a caulescent rosette plant endemic to Northeast Brazil. *Acta Botanica Brasilica* **26**: 94–100.
- Coutinho LM. 1976.** *Contribuição ao conhecimento do papel ecológico das queimadas na floração de espécies do cerrado*. Unpublished D. Phil. Thesis, Universidade de São Paulo, São Paulo, Brazil.
- Coutinho LM. 1990.** Fire in the ecology of the Brazilian cerrado. In: Goldammer JG, ed. *Fire in the tropical biota – ecosystem process and global challenges*. Berlin: Springer-Verlag, 82–105.
- Doyle JJ, Doyle JL. 1993.** Chloroplast DNA phylogeny of the papilionoid legume tribe Phaseoleae. *Systematic Botany* **18**: 309–327.
- Drummond AJ, Suchard MA, Xie D, Rambaut A. 2012.** Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Molecular Biology and Evolution* **29**: 1969–1973.
- Dwyer E, Pincock S, Gregoire JM, Pereira JMC. 2000.** Global spatial and temporal distribution of vegetation fire as determined from satellite observations. *International Journal of Remote Sensing* **21**: 1289–1302.
- Edwards EJ, Osborne CP, Stromberg CAE, Smith SA, C4 Grasses Consortium. 2010.** The origins of C₄ grasslands: integrating evolutionary and ecosystem science. *Science* **328**: 587–591.
- Egan AN, Vatanparast M, Cagle W. 2016.** Parsing polyphyletic *Pueraria*: delimiting distinct evolutionary lineages through phylogeny. *Molecular Phylogenetics and Evolution* **104**: 44–59.
- Fidelis A, Blanco C. 2014.** Does fire induce flowering in Brazilian subtropical grasslands? *Applied Vegetation Science* **17**: 690–699.
- Fortuna-Perez AP, Cândido ES, Silva MJ, Vargas W, Bezerra LMPA, Vatanparast M. 2018.** A noteworthy new species of *Eriosema* (Leguminosae, Papilionoideae, Phaseoleae) from Goiás State, Brazil, including an identification key. *Systematic Botany* **43**: 198–205.
- Fortuna-Perez AP, Lewis GP, Cândido ES, Bezerra LMPA, Tozzi AMA. 2013b.** *Eriosema hatschbachii* (Leguminosae, Papilionoideae), a new species from Minas Gerais, Brazil. *Kew Bulletin* **68**: 641–645.
- Fortuna-Perez AP, Silva MJ, Cândido ES, Vargas W, Monteiro TC, Vatanparast M. 2017.** *Eriosema elegans* (Leguminosae, Papilionoideae): a new species from the Highlands of Goiás State, Brazil. *Phytotaxa* **296**: 81–87.
- Fortuna-Perez AP, Silva MJ, Queiroz LP, Lewis GP, Simões AO, Tozzi AMGA, Sarkinen TE, Souza AP. 2013a.** Phylogeny and biogeography of the genus *Zornia* (Leguminosae: Papilionoideae: Dalbergieae). *Taxon* **62**: 723–733.
- Fortunato RH. 1993.** Cambios nomenclaturales en *Eriosema* (Fabaceae: Papilionoideae, Cajaninae). *Novon* **3**: 24–27.
- Fortunato RH. 1999.** Cambios nomenclaturales en *Eriosema* (Fabaceae: Papilionoideae, Cajaninae) II. *Kurtziana* **27**: 371–382.
- Givnish T, Renner S. 2004.** Tropical intercontinental disjunctions: Gondwana breakup, immigration from the boreotropics, and transoceanic dispersal. *International Journal of Plant Sciences* **165**: S1–S6.
- Greear JW. 1970.** A revision of the American species of *Eriosema* (Leguminosae-Lotoideae). *Memoirs of the New York Botanical Garden* **20**: 1–98.
- Greear JW. 1978.** A revision of the New World species of *Rhynchosia* (Leguminosae, Faboideae). *Memoirs of the New York Botanical Garden* **31**: 1–168.
- Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010.** New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Systematic Biology* **59**: 307–321.
- Herendeen PS, Crepet WL, Dilcher DL. 1992.** The fossil history of the Leguminosae: phylogenetic and biogeographic implications. In: Herendeen PS, Dilcher DL, eds. *Advances in legume systematics. Part 4. The fossil record*. Kew: Royal Botanic Gardens, 303–316.
- Hirschmann GS, De Arias AR. 1990.** A survey of medicinal plants of Minas Gerais, Brazil. *Journal of Ethnopharmacology* **29**: 159–172.
- Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018.** UFBoot2: Improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* **35**: 518–522.
- Hoekstra JM, Boucher TM, Ricketts TH, Roberts C. 2005.** Confronting a biome crisis: global disparities of habitat loss and protection. *Ecology Letters* **8**: 23–29.
- Kaila T, Chaduvla PK, Saxena S, Bahadur K, Gahukar SJ, Chaudhury A, Sharma TR, Singh NK, Gaikwad K. 2016.** Chloroplast genome sequence of pigeonpea (*Cajanus cajan* (L.) Millspaugh) and *Cajanus scarabaeoides* (L.) Thouars: genome organization and comparison with other legumes. *Frontier in Plant Sciences* **7**: 1847.
- Kajita K, Ohashi H, Tateishi Y, Bailey CD, Doyle JJ. 2001.** *rbcl* and legume phylogeny, with particular reference to Phaseoleae, Millettieae, and allies. *Systematic Botany* **26**: 515–536.
- Kalyanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermin LS. 2017.** ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods* **14**: 587–589.
- Kassa MT, Penmetse RV, Carrasquilla-Garcia N, Sarma BK, Datta S, Upadhyaya HD, Varshney RK, von Wettberg EJB, Cook DR. 2012.** Genetic patterns of domestication in pigeonpea (*Cajanus cajan* (L.) Millsp.) and wild *Cajanus* relatives. *PLoS One* **7**: e39563.
- Katoh K, Standley DM. 2013.** MAFFT: iterative refinement and additional methods. In: Russell DJ, ed. *Methods in molecular biology. Multiple sequence alignment methods*. New York: Springer, 131–146.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Mentjies P, Drummond A. 2012.** Geneious Basic: an integrated and extendable desktop

- software platform for the organization and analysis of sequence data. *Bioinformatics* **28**: 1647–1649.
- Keeley JE, Rundel PW. 2005.** Fire and the Miocene expansion of *C₄* grasslands. *Ecology Letters* **8**: 683–90.
- Kergoat GJ, Le Ru BP, Genson G, Cruaud C, Couloux A, Delobel A. 2011.** Phylogenetics, species boundaries and timing of resource tracking in a highly specialized group of seed beetles (Coleoptera: Chrysomelidae: Bruchinae). *Molecular Phylogenetics and Evolution* **59**: 746–760.
- Lackey JA. 1977.** A revised classification of the tribe Phaseoleae (Leguminosae: Papilionoideae) and its relation to canavanine distribution. *Botanical Journal of the Linnean Society* **74**: 163–178.
- Lackey JA. 1978.** Leaflet anatomy of Phaseoleae (Fabaceae, Papilionoideae) and its relation to taxonomy. *Botanical Gazette* **139**: 346–446.
- Lackey JA. 1981.** Phaseoleae. In: Polhill RM, Raven PH, eds. *Advances in legume systematics. Part 1*. Kew: Royal Botanic Gardens, 301–321.
- Larsson A. 2014.** AliView: a fast and lightweight alignment viewer and editor for large datasets. *Bioinformatics* **30**: 3276–3278.
- Lavin M, Herendeen PS, Wojciechowski MF. 2005.** Evolutionary rates analysis of Leguminosae implicates a rapid diversification of lineages during the Tertiary. *Systematic Biology* **54**: 575–594.
- Lavin M, Schrire BP, Lewis GP, Pennington RT, Delgado-Salinas A, Thulin M, Hughes CE, Matos AB, Wojciechowski MF. 2004.** Metacommunity process rather than continental tectonic history better explains geographically structured phylogenies in legumes. *Philosophical Transactions of the Royal Society of London, B, Biological Sciences* **359**: 1509–1522.
- Lewis G, Schrire BP, MacKinder B, Lock M. 2005.** *Legumes of the world*. Kew: Royal Botanic Gardens.
- Li H, Wang W, Lin L, Zhu X, Li J, Zhu X, Chen Z. 2013.** Diversification of the phaseoloid legumes: effects of climate change, range expansion and habit shift. *Frontiers in Plant Science* **4**: 386.
- Lock JM. 2006.** The seasonally dry vegetation of Africa: parallels and comparisons with the Neotropics. In: Pennington RT, Lewis GP, Ratter JA, eds. *Neotropical savannas and dry forests: plant diversity, biogeography and conservation*. Boca Raton: CRC Press, 449–467.
- LPWG (The Legume Phylogeny Working Group). 2017.** A new subfamily classification of the Leguminosae based on a taxonomically comprehensive phylogeny. *Taxon* **66**: 44–77.
- Maurin O, Davies TJ, Burrows JE, Daru BH, Yessoufou K, Muasya AM, van der Bank M, Bond WJ. 2014.** Savanna fire and the origins of the ‘underground forests’ of Africa. *New Phytologist* **204**: 201–214.
- Miotto STS. 1988.** Leguminosae, Faboideae, tribo Phaseoleae, subtribo Cajaninae. *Flora Ilustrada do Rio Grande do Sul, fascículo XIX. Boletim do Instituto de Biociências* **43**: 1–88.
- Moura TM, Vatanparast M, Tozzi AMGA, Forest F, Wilmot-Dear CM, Simon MF, Mansano VF, Kajita T, Lewis GP. 2016.** A molecular phylogeny and new infrageneric classification of *Mucuna* Adans. (Leguminosae-Papilionoideae) including insights from morphology and hypotheses about biogeography. *International Journal of Plant Sciences* **177**: 76–89.
- Myers N, Mittermeier RA, Mittermeier CG, Fonseca GAB, Kent J. 2000.** Biodiversity hotspots for conservation priorities. *Nature* **403**: 853–858.
- Nadimpalli RG, Jarret RL, Phatak SC, Kochert G. 1993.** Phylogenetic relationships of the pigeonpea (*Cajanus cajan*) based on nuclear restriction fragment length polymorphisms. *Genome* **36**: 216–23.
- Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015.** IQ-TREE: a fast and effective stochastic algorithm for estimating maximum likelihood phylogenies. *Molecular Biology and Evolution* **32**: 268–274.
- Nylander JAA. 2004.** *MrModeltest 2.3. Program distributed by the author*. Uppsala: Evolutionary Biology Centre, Uppsala University.
- Ojewole JAO. 2007.** African traditional medicines for erectile dysfunction: elusive dream or imminent reality? *Cardiovascular Journal of Africa* **18**: 213–215.
- Oliveira ACS, Fortuna-Perez AP, Silva JS. 2018.** Os gêneros *Eriosema* e *Rhynchosia* (Leguminosae - Papilionoideae - Phaseoleae) no nordeste brasileiro. *Rodriguésia* **69**: 1825–1850.
- Parr CL, Lehmann CER, Bond WJ, Hoffmann WA, Andersen AN. 2014.** Tropical grassy biomes: misunderstood, neglected, and under threat. *Trends in Ecology & Evolution* **29**: 205–213.
- Pennington RT, Hughes CE. 2014.** The remarkable congruence of New and Old World savannah origins. *New Phytologist* **204**: 4–6.
- Pennington RT, Prado DE, Pendry CA. 2000.** Neotropical seasonally dry forests and Quaternary vegetation changes. *Journal of Biogeography* **27**: 261–273.
- Pennington RT, Richardson JE, Lavin M. 2006.** Insights into the historical construction of species-rich biomes from dated plant phylogenies, neutral ecological theory and phylogenetic community structure. *New Phytologist* **172**: 605–616.
- Phipps JB, Goodier R. 1962.** A preliminary account of the plant ecology of the Chimanimani mountains. *Journal of Ecology* **50**: 291–319.
- Piva T, Fortuna-Perez AP, Vargas W, Machado SR. 2020.** Variations in the architecture and histochemistry of the gelatinous fibers in *Eriosema* (DC.) Desv. (Leguminosae) species from the Brazilian Cerrado. *Flora* **268**: 151624.
- Queiroz LP, Pastore JFB, Cardoso D, Snak C, Lima ALC, Gagnon E, Vatanparast M, Holland AE, Egan AN. 2015.** A multilocus phylogenetic analysis reveals the monophyly of a circumscribed papilionoid legume tribe Diocleae with well-supported generic relationships. *Molecular Phylogenetics and Evolution* **90**: 1–19.
- Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. 2018.** Posterior summarisation in Bayesian phylogenetics using Tracer 1.7. *Systematic Biology* **67**: 901–904.
- Rando JG, Zuntini AR, Conceição AS, Berg C, Pirani JR, Queiroz LP. 2016.** Phylogeny of *Chamaecrista* ser. *Coriaceae* (Leguminosae) unveils a lineage recently diversified in

- Brazilian campo rupestre vegetation. *International Journal of Plant Science* **177**: 3–17.
- Ratnam J, Bond WJ, Fensham RJ, Hoffmann WA, Archibald S, Lehmann CER, Anderson MT, Higgins SI, Sankaran M. 2011.** When is a 'forest' a savanna, and why does it matter? *Global Ecology Biogeography* **20**: 653–660.
- Rodrigues VEG, Carvalho DA. 2001.** Levantamento etnobotânico de plantas medicinais no domínio do cerrado na região do Alto Rio Grande - Minas Gerais. *Revista Ciência e Agrotecnologia* **25**: 102–123.
- Ronquist F, Huelsenbeck J, Teslenko M. 2011.** *Draft MrBayes version 3.2 manual: tutorials and model summaries*. Available at: <https://nbisweden.github.io/MrBayes/manual.html>.
- Ronquist F, Teslenko M, Van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP. 2012.** MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* **61**: 539–542.
- Sano EE, Rosa R, Brito JLS, Ferreira Jr LG. 2010.** Land cover mapping of the tropical savanna region in Brazil. *Environmental Monitoring and Assessment* **166**: 113–124.
- Santos MG, Almeida VG, Avelar-Freitas BA, Graef CFF, Gregório LE, Pereira WF, Brito-Melo GEA. 2016.** Phytochemical screening of the dichloromethane ethanolic extract of *Eriosema campestre* var. *macrophyllum* roots and its antiproliferative effect on human peripheral blood lymphocytes. *Revista Brasileira Farmacognosia* **26**: 464–470.
- Saxena RK, von Wettberg E, Upadhyaya HD, Sanchez V, Songok S, Saxena K, Kimurto P, Varshney RK. 2014.** Genetic diversity and demographic history of *Cajanus* spp. illustrated from genome-wide SNPs. *PLoS One* **9**: e88568.
- Scholes RJ, Archer SR. 1997.** Tree–grass interactions in savannas. *Annual Review of Ecology and Systematics* **28**: 517–544.
- Schrire BD. 2005.** Phaseoleae. In: Lewis G, Schrire B, Mackinder B, Lock M, eds. *Legumes of the world*. Kew: Royal Botanic Gardens, 393–431.
- Seixas DP, Fortuna-Perez AP, Rodrigues TM. 2019.** Leaf anatomical features of the *Eriosema campestre* Benth. (Leguminosae, Papilionoideae, Phaseoleae) complex and potential taxonomic implications. *Flora* **253**: 107–115.
- Shaw J, Lickey EB, Schilling EE, Small RL. 2007.** Comparison of whole chloroplast genome sequences to choose noncoding regions for phylogenetic studies in angiosperms: the tortoise and the hare III. *American Journal of Botany* **94**: 275–288.
- Simon MF, Grether R, Queiroz LP, Sarkinen TE, Dutra VF, Hughes CE. 2011.** The evolutionary history of *Mimosa* (Leguminosae): towards a phylogeny of the sensitive plants. *American Journal of Botany* **98**: 1201–1221.
- Simon MF, Grether R, Queiroz LP, Skema C, Pennington RT, Hughes CE. 2009.** Recent assembly of the cerrado, a Neotropical plant diversity hotspot, by *in situ* evolution of adaptations to fire. *Proceedings of the National Academy of Sciences of the United States of America* **106**: 20359–20364.
- Simon MF, Pennington RT. 2012.** Evidence for adaptation to fire regimes in the tropical savannas of the Brazilian cerrado. *International Journal of Plant Science* **173**: 711–723.
- Smith NM. 1991.** Ethnobotanical field notes from the Northern Territory, Australia. *Journal of the Adelaide Botanic Garden* **14**: 1–65.
- Snak C, Vatanparast M, Silva C, Lewis GP, Lavin M, Kajita T, Queiroz LP. 2016.** A dated phylogeny of the papilionoid legume genus *Canavalia* reveals recent diversification by a pantropical liana lineage. *Molecular Phylogenetics and Evolution* **98**: 133–146.
- Souza ER, Lewis GP, Forest F, Schnadelbach AS, van den Berg C, Queiroz LP. 2013.** Phylogeny of *Calliandra* (Leguminosae: Mimosoideae) based on nuclear and plastid molecular markers. *Taxon* **62**: 1200–1219.
- Souza-Neto AC, Cianciaruso MV, Collevatti RG. 2015.** Habitat shifts shaping the diversity of a biodiversity hotspot through time: insights from the phylogenetic structure of Caesalpinioideae in the Brazilian cerrado. *Journal of Biogeography* **43**: 340–350.
- Stirton CH. 1975.** A contribution to knowledge of the genus *Eriosema* (Leguminosae, Lotoideae) in Southern Africa (excluding Mozambique and Rhodesia). M.Sc. thesis, University of Natal.
- Stirton CH. 1981.** The *Eriosema cordatum* complex. II. The *Eriosema cordatum* and *E. nutans* groups. *Bothalia* **13**: 281–306.
- Stirton CH. 1986.** The *Eriosema squarrosus* complex (Papilionoideae, Fabaceae) in southern Africa. *Bothalia* **16**: 11–22.
- Stirton CH, Gordon-Gray KD. 1978.** The *Eriosema cordatum* complex. 1. The *Eriosema populifolium* group. *Bothalia* **12**: 395–404.
- Sun Y, Skinner DZ, Liang GH, Hulbert SH. 1994.** Phylogenetic analysis of *Sorghum* and related taxa using internal transcribed spacers of nuclear ribosomal DNA. *Theoretical and Applied Genetics* **89**: 26–32.
- Thorne RF. 1972.** Major disjunctions in the geographic ranges of seed plants. *The Quarterly Review of Biology* **47**: 365–411.
- Tosso F, Hardy OJ, Doucet JL, Daïnou K, Kaymak E, Migliore J. 2018.** Evolution in the Amphi-Atlantic tropical genus *Guibourtia* (Fabaceae, Detarioideae), combining NGS phylogeny and morphology. *Molecular Phylogenetics and Evolution* **120**: 83–93.
- Trovó M, Andrade MJG, Sano PT, Ribeiro PL, van den Berg C. 2013.** Molecular phylogenetics and biogeography of Neotropical Paepalanthoideae with emphasis on Brazilian *Paepalanthus* (Eriocaulaceae). *Botanical Journal of the Linnean Society* **171**: 225–243.
- Van der Maesen LJG. 2003.** Cajaninae of Australia (Leguminosae: Papilionoideae). *Australian Systematic Botany* **16**: 219–227.
- Van der Pijl L. 1982.** *Principles of dispersal in higher plants*. Berlin, Heidelberg: Springer.
- Van der Werf GR, Randerson JT, Giglio L, Collatz GJ, Mu M, Kasibhatla PS, Morton DC, DeFries RS, Jin Y, van Leeuwen TT. 2010.** Global fire emissions and the contribution of deforestation, savanna, forest, agricultural,

- and peat fires (1997–2009). *Atmospheric Chemistry and Physics* **10**: 11707–11735.
- Vargas W. 2019.** *Filogenia, biogeografia e evolução de estruturas secretoras de representantes da subtribo Cajaninae (Leguminosae, Papilionoideae, Phaseoleae)*. Unpublished D. Phil. Thesis, São Paulo State University.
- Vargas W, Fortuna-Perez AP, Lewis GP, Piva TC, Vatanparast M, Machado SR. 2019.** Ultrastructure and secretion of glandular trichomes in species of subtribe Cajaninae Benth. (Leguminosae, Phaseoleae). *Protoplasma* **256**: 431.
- Vargas W, Machado SR, Lewis GP, Cândido ES, Vatanparast M, Fortuna-Perez AP. 2018.** Revisiting the leaflet secretory structures in subtribe Cajaninae Benth. (Leguminosae, Phaseoleae). *International Journal of Plant Sciences* **179**: 697–711.
- Vasconcelos TNC, Alcantara S, Andrino CO, Forest F, Reginato M, Simon MF, Pirani JR. 2020.** Fast diversification through a mosaic of evolutionary histories characterizes the endemic flora of ancient Neotropical mountains. *Proceedings of the Royal Society B* **287**: 20192933.
- Vatanparast M, Klitgaard BB, Adema FACB, Pennington RT, Yahara T, Kajita T. 2013.** First molecular phylogeny of the pantropical genus *Dalbergia*: implications for infrageneric circumscription and biogeography. *South African Journal of Botany* **89**: 143–149.
- Veldman JW, Buisson E, Durigan G, Fernandes GW, Le Stradic S, Mahy G, Negreiros D, Overbeck GE, Veldman RG, Zaloumis NP, Putz FE, Bond WJ. 2015.** Toward an old-growth concept for grasslands, savannas, and woodlands. *Frontiers in Ecology and the Environment* **13**: 154–162.
- White F. 1983.** *The vegetation of Africa. A descriptive memoir to accompany the UNESCO/AETFAT/UNSO vegetation map of Africa*. Paris: UNESCO.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

Appendix S1. Nucleotide sequence database accession numbers of taxa used in the phylogenetic analysis.

Appendix S2. List of molecular markers and primers used in this study.

Appendix S3. Information relating to the separate DNA datasets.

Appendix S4. Phylogenetic tree for *Eriosema* from maximum likelihood analysis of the ITS/5.8S and plastid DNA dataset (*rpl32-trnL* and *trnQ-5'rps16*). Numbers along branches are bootstrap values and posterior probabilities.

Appendix S5. BEAST chronogram based on combined dataset (ITS/5.8S, *rpl32-trnL* and *trnQ-5'rps16*). Numbers at nodes refer to mean age of the nodes (My). Blue bars represent 95% confidence intervals (CI) for the estimated mean dates. Posterior probabilities are indicated at nodes.